A&A 436, 503–526 (2005)
DOI: 10.1051/0004-6361:20042520

Astronomy & Astrophysics

Relativistic MHD simulations of extragalactic jets*

T. Leismann[1], L. Antón[2], M. A. Aloy[1], E. Müller[1], J. M. Martí[2], J. A. Miralles[3], and J. M. Ibáñez[2]

[1] Max-Planck-Institut für Astrophysik, Postfach 1312, 85741 Garching, Germany
e-mail: maa@mpa-garching.mpg.de
[2] Departamento de Astronomía y Astrofísica, Universidad de Valencia, 46100 Burjassot, Spain
[3] Departamento de Física Aplicada, Universidad de Alicante, Ap. Correus 99, 03080 Alacant, Spain

Received 10 December 2004 / Accepted 5 March 2005

Abstract. We have performed a comprehensive parameter study of the morphology and dynamics of axisymmetric, magnetized, relativistic jets by means of numerical simulations. The simulations have been performed with an upgraded version of the GENESIS code which is based on a second-order accurate finite volume method involving an approximate Riemann solver suitable for relativistic ideal magnetohydrodynamic flows, and a method of lines. Starting from pure hydrodynamic models we consider the effect of a magnetic field of increasing strength (up to $\beta \equiv |b|^2/2p \approx 3.3$ times the equipartition value) and different topology (purely toroidal or poloidal). We computed several series of models investigating the dependence of the dynamics on the magnetic field in jets of different beam Lorentz factor and adiabatic index.

We find that the inclusion of the magnetic field leads to diverse effects which contrary to Newtonian magnetohydrodynamics models do not always scale linearly with the (relative) strength of the magnetic field. The relativistic models show, however, some clear trends. Axisymmetric jets with toroidal magnetic fields produce a cavity which consists of two parts: an inner one surrounding the beam which is compressed by magnetic forces, and an adjacent outer part which is inflated due to the action of the magnetic field. The outer border of the outer part of the cavity is given by the bow-shock where its interaction with the external medium takes place. Toroidal magnetic fields well below equipartition ($\beta = 0.05$) combined with a value of the adiabatic index of 4/3 yield extremely smooth jet cavities and stable beams.

Prominent nose cones form when jets are confined by toroidal fields and carry a high Poynting flux ($\sigma \equiv |b|^2/\rho > 0.01$ and $\beta \geq 1$). In contrast, none of our models possessing a poloidal field develops such a nose cone. The size of the nose cone is correlated with the propagation speed of the Mach disc (the smaller the speed the larger is the size). If two models differ only by the adiabatic index, jets having smaller adiabatic indices tend to develop smaller nose cones.

Key words. magnetohydrodynamics (MHD) – methods: numerical – relativity – galaxies: active – galaxies: jets

1. Introduction

The presence of magnetic fields in relativistic fluids is ubiquitous among the most luminous objects in the universe like, e.g., active galactic nuclei (AGNs), quasars, gamma-ray bursts, X-ray binaries, core-collapse supernovae, etc. In the specific case of jets originating from AGNs, the observation of non-thermal synchrotron radiation (see, e.g., Ferrari 1998, for a review) requires the presence of a magnetic field in which relativistic electrons gyroradiate. The high degree of polarization observed in many AGN sources (e.g., Gabuzda et al. 2000) indicates that magnetic fields are not randomly oriented but posses some large scale structure (at least at parsec scales).

Disentangling the intrinsic structure of the magnetic field is still an observational challenge because of the finite spatial resolution of the interferometric beams used in observations of AGN jets. At parsec scales a variety of magnetic field configurations are observed including predominantly toroidal ones (e.g., 0954 + 658: Gabuzda & Cawthorne 1996; 1803 + 784: 0823 + 033 and 1749 + 701: Gabuzda & Pushkarev 2001), configurations with alternating orthogonal, alternating aligned and predominantly poloidal fields along the jet (e.g., OJ 287: Gabuzda & Gómez 2001; 1418 + 546: Gabuzda 2003), and even helical field configurations (1055 + 018: Attridge et al. 1999; 0820 + 225: Gabuzda et al. 2001; 0745 + 241: Pushkarev & Gabuzda 2001; 1652 + 398: Gabuzda 2003). Different magnetic field topologies are also observed at kiloparsec scales ranging from fields which are mostly aligned with the jet (e.g., 3C 120: Walker et al. 1987; NGC 4258: Krause & Löhr 2004) to fields which are oriented preferentially perpendicular (toroidal) to the jet axis (e.g., 3C 449: Kigure et al. 2004). Other more complex structures are observed, too (e.g., Laing & Bridle 2002). Furthermore, there are indications that the intergalactic and interstellar medium the AGN jets are propagating

* Appendices are only available in electronic form at
http://www.edpsciences.org

into can be threaded with large scale magnetic fields of some microgauss (e.g., Kim et al. 1990; Crusius-Waetzel et al. 1990; Taylor & Perley 1993). This variety of ordered magnetic field structures most likely reflects different formation and evolutionary processes. For example, magnetized jets with a dominant toroidal field component can be produced by winding up an initial field which has a significant longitudinal component due to the rotation of the central AGN engine (e.g., Nakamura et al. 2001; Lovelace et al. 2002; Tsinganos & Bogovalov 2002; Lynden-Bell 2003).

Relativistic magnetized jets from AGNs can be described in the framework of ideal relativistic magnetohydrodynamics (RMHD). Among the first attempts to simulate the corresponding nonlinear, time dependent and multidimensional RMHD equations we mention here that of Evans & Hawley (1988), who using a two-dimensional finite difference code investigated several problems involving general relativistic magnetohydrodynamic accretion flows onto a black hole. More importantly, these authors also proposed a new numerical technique called constrained transport (CT) for evolving the induction equation while maintaining vanishing divergence of the magnetic field down to machine roundoff error. Later on van Putten (1993, 1996) performed the first axisymmetric simulations of RMHD jets having moderately small Lorentz factors ($\lesssim$3). Koide & coworkers (Koide et al. 1996; Koide 1997) simulated 2D RMHD slab jets being restricted, however, to rather low spatial resolution, short evolution times, and small Lorentz factors ($\lesssim$4.6). This study was extended to 3D jets by Nishikawa & coworkers (Nishikawa et al. 1997, 1998). Komissarov developed a high-resolution, shock-capturing, second-order accurate RMHD code (Komissarov 1999a) which he used to simulate 2D jets possessing toroidal fields (Komissarov 1999b) and the jet-torus structure observed in the Crab nebula (Komissarov & Lyubarsky 2003, 2004, see also Del Zanna et al. 2004). Jet formation has been studied using general relativistic magnetohydrodynamic (GRMHD) simulations based on a simplified Total Variation Diminishing scheme ignoring the evolution of the constraint $\nabla \cdot \boldsymbol{B} = 0$. Koide et al. (1998, 1999, 2000, 2002) and Koide (2003) modeled the formation of axisymmetric jets in a system consisting of an accretion disk and a rotating black hole. The simulations, which cover a few rotational periods of the black hole, show the formation of a relativistic outflow with a Lorentz factor $W \approx 2$. A similar study but without any symmetry restrictions was carried by Nishikawa et al. (2002, 2003). Jet formation in the context of gamma-ray burst progenitors was considered by Mizuno et al. (2004a,b) using a GRMHD code (and no specific measures to maintain $\nabla\cdot\boldsymbol{B} = 0$). The evolution of magnetized accretion tori around rotating black holes was investigated in axisymmetry by Gammie et al. (2003) and De Villiers & Hawley (2003); De Villiers & Hawley (2003) including a CT algorithm to keep the magnetic field divergence-free. While Gammie et al. (2003) used a Godunov-type conservative scheme, De Villiers & Hawley (2003) employed a ZEUS-like (i.e., non conservative) scheme to integrate the GRMHD equations.

In previous 2D RMHD simulations relativistic jets only served the purpose of demonstrating the capabilities of the RMHD code (e.g., Komissarov 1999a, in slab geometry; Del Zanna et al. 2003, assuming axial symmetry and using cylindrical coordinates), or the simulations only covered a very limited range of jet parameters (e.g., in Komissarov 1999b, only two models were considered both involving only toroidal fields). In the following we present the first comprehensive parameter study of the morphology and dynamics of magnetized relativistic axisymmetric jets including both toroidal and poloidal field configurations of different strength, and beam plasmas of different adiabatic index. Preliminary results of our research can be found in Leismann et al. (2004).

The paper is organized as follows. The basic equations and the numerical algorithm used in the code are discussed in Sect. 2. As we assume cylindrical symmetry in our simulations of relativistic magnetized jets, we give the explicit form of the corresponding equations in the Appendix A. Various tests our code has passed successfully are described in Appendix B. In Sect. 3 we discuss the simulation setup and the model parameters, and in Sect. 4 we present the results of our study. Finally, in Sect. 5 we discuss and interpret our findings, and point out some limitations of our approach.

2. Numerical method

In this section we describe the details of the numerical algorithm which we used to integrate the equations of ideal relativistic magnetohydrodynamics (Sect. 2.1). The numerical method is implemented on an upgraded version of GENESIS (Aloy et al. 1999b) and relies on the modular structure of its predecessor. As GENESIS, it is based on a directional-splitting, Godunov-type, finite-volume method, a series of intercell reconstruction routines and a method of lines for the time advance. The new code is second order accurate both in space and time, and relies on a constrained transport method (mainly based on the developments of Ryu et al. 1998) in order to keep (to machine precision) the magnetic field divergence-free throughout a simulation. Unlike GENESIS, the numerical fluxes at the cell interfaces are computed employing an HLL-type solver (Sect. 2.3.1) which uses as the maximum and minimum signal speeds some accurate estimates which are given in Sect. 2.2.1 together with the generic spectral decomposition and properties of the RMHD system of conservation laws. The combination of an HLL-like solver along with a high order spatial reconstruction and a consistently high order method to keep the divergence-free property of the magnetic field was proven to be useful to build RMHD algorithm by Del Zanna et al. (2003). Finally, a new algorithm to recover the primitive variables from the conserved ones has been implemented (Sect. 2.3.3).

2.1. Equations of ideal relativistic MHD

The equations that describe the evolution of an ideal relativistic magneto-fluid can be written in the form of conservation laws (see, e.g., Anile 1989, for a full derivation of the equations), the conservation of mass,

$$\nabla_\alpha(\rho u^\alpha) = 0, \tag{1}$$

and the conservation of total energy-momentum,

$$\nabla_\beta T^{\alpha\beta} = 0, \tag{2}$$

where ∇_α is the covariant derivative with respect to the metric tensor $g_{\alpha\beta}$. The energy-momentum tensor of a magnetic perfect fluid is given by[1].

$$T^{\alpha\beta} = \rho h^* u^\alpha u^\beta + p^* g^{\alpha\beta} - b^\alpha b^\beta. \tag{3}$$

Here ρ is the rest mass density of the fluid, u^α the 4-velocity, b^α the magnetic field measured by a comoving observer, h^* and p^* are the hydromagnetic specific enthalpy and the hydromagnetic total pressure, respectively, given by

$$h^* = h + \frac{b^\alpha b_\alpha}{\rho} = 1 + \varepsilon + \frac{p}{\rho} + \frac{|b|^2}{\rho} \quad \text{and} \quad p^* = p + p_{\rm mag}, \tag{4}$$

where p is the fluid pressure, ε denotes the specific internal energy, $h = 1 + \varepsilon + p/\rho$ is the specific enthalpy,

$$p_{\rm mag} = \frac{|b|^2}{2} \tag{5}$$

is the magnetic pressure, and $|b|^2 = b^\alpha b_\alpha$ is the magnetic energy density.

Introducing the 3-velocity vector $v^i = u^i/u^0$ (Latin indices run from 1 to 3, or x, y, z in Cartesian coordinates), the 4-velocity u^α can be written as

$$u^\alpha = W(1, v^1, v^2, v^3), \tag{6}$$

where $W = 1/\sqrt{1 - \boldsymbol{v}^2}$ is the Lorentz factor and $\boldsymbol{v}^2 = v^i v_i$. Using the infinite conductivity condition, which states that the electric field in the comoving frame must be zero, the magnetic field b^α can be written in terms of the magnetic field measured in the laboratory frame, which is given by the 3-vector $\boldsymbol{B}$, as follows

$$b^0 = W(\boldsymbol{v} \cdot \boldsymbol{B}), \tag{7}$$

$$b^i = \frac{B^i}{W} + v^i b^0. \tag{8}$$

In terms of $\boldsymbol{v}$ and $\boldsymbol{B}$, we have $|b|^2 = B^2 W^{-2} + (\boldsymbol{v} \cdot \boldsymbol{B})^2$.

The evolution of the magnetic field components is described by the homogeneous Maxwell equation,

$$\nabla_\alpha (u^\alpha b^\beta - u^\beta b^\alpha) = 0, \tag{9}$$

whose spatial part (in terms of the magnetic field in the laboratory frame) leads to the induction equation

$$\frac{\partial \boldsymbol{B}}{\partial t} - \nabla \times (\boldsymbol{v} \times \boldsymbol{B}) = 0, \tag{10}$$

in complete analogy with the Newtonian case.

The time component of Eq. (9) becomes the usual divergence constraint

$$\nabla \cdot \boldsymbol{B} = 0, \tag{11}$$

which has to be fulfilled at all times.

[1] In the following we will use a system of units where the speed of light in vacuum, $c = 1$, and a factor of $1/\sqrt{4\pi}$ is absorbed into the definition of the magnetic field. Greek subscripts in 4-vectors run from 0 to 3. We assume a metric signature $(-, +, +, +)$.

Equations (1), (2) and (10), together with the constraint (11) and the equation of state (EOS)

$$p = p(\varepsilon, \rho) \tag{12}$$

provide the complete set of RMHD equations.

In the following we will use an ideal EOS with a constant adiabatic index, γ:

$$p = (\gamma - 1)\rho\varepsilon. \tag{13}$$

To characterize the importance of the magnetic field, it is useful to introduce the *magnetization parameter*, β, which is the ratio of magnetic to gas pressure (defined as the reciprocal of the standard β-parameter of plasma physics):

$$\beta \equiv \frac{p_{\rm mag}}{p}. \tag{14}$$

Following Appl & Camenzind (1988), we introduce a further parameter that will prove useful in parameterizing RMHD flows of astrophysical interest: the ratio of magnetic energy density to that of the rest mass given by

$$\sigma \equiv \frac{|b|^2}{\rho} \quad \left(= \frac{2p_{\rm mag}}{\rho} \right). \tag{15}$$

Then $h^* = h + \sigma$, and for $\sigma \gg h$ the Poynting flux $\rho\sigma W^2 \boldsymbol{v}$ will be much larger than the material energy flux, $\rho h W^2 \boldsymbol{v}$. Note that we define the parameter σ differently from Appl & Camenzind (1988) or Komissarov (1999b), who defined it to be the inverse ratio. According to our definition both β and σ become large for large magnetic fields and zero for non-magnetized plasma.

The speed of sound, $c_{\rm s}$, can then be calculated from (e.g. Landau & Lifshitz 1966)

$$h c_{\rm s}^2 = \left.\frac{\partial p}{\partial \rho}\right|_\varepsilon + \frac{p}{\rho^2} \left.\frac{\partial p}{\partial \varepsilon}\right|_\rho, \tag{16}$$

i.e.,

$$c_{\rm s} = \sqrt{\frac{\gamma p}{\rho h}}. \tag{17}$$

The Alfvén speed, $c_{\rm a}$, is defined through (Anile 1989)

$$c_{\rm a}^2 = \frac{|b|^2}{\rho h^*} = \frac{\sigma}{h + \sigma}. \tag{18}$$

2.2. Primitive and conserved variables

The RMHD system of partial differential equations in a flat space-time and in Cartesian coordinates can be cast in conservation form as follows

$$\partial_t \boldsymbol{U} + \partial_x \boldsymbol{F}^x + \partial_y \boldsymbol{F}^y + \partial_z \boldsymbol{F}^z = \boldsymbol{0}, \tag{19}$$

where $\boldsymbol{U}$ is the vector of *conserved quantities* and $\boldsymbol{F}^i$ are the vectors of *fluxes*. Written in terms of *primitive variables*,

$$\boldsymbol{V} = (\rho, v^x, v^y, v^z, p, B^x, B^y, B^z)^T, \tag{20}$$

and using the symbols D, $\boldsymbol{S}$ and τ to denote the rest mass density, momentum density vector and energy density measured in the laboratory frame, respectively, the conserved quantities, $\boldsymbol{U}$, read

$$\boldsymbol{U} = \begin{pmatrix} D \\ S^x \\ S^y \\ S^z \\ \tau \\ B^x \\ B^y \\ B^z \end{pmatrix} \equiv \begin{pmatrix} \rho W \\ \rho h^* W^2 v^x - b^0 b^x \\ \rho h^* W^2 v^y - b^0 b^y \\ \rho h^* W^2 v^z - b^0 b^z \\ \rho h^* W^2 - p^* - b^0 b^0 - \rho W \\ B^x \\ B^y \\ B^z \end{pmatrix}. \quad (21)$$

The fluxes, $\boldsymbol{F}^i$, are then

$$\boldsymbol{F}^i = \begin{pmatrix} \rho W v^i \\ \rho h^* W^2 v^i v^x + p^* \delta^i_x - b^i b^x \\ \rho h^* W^2 v^i v^y + p^* \delta^i_y - b^i b^y \\ \rho h^* W^2 v^i v^z + p^* \delta^i_z - b^i b^z \\ \rho h^* W^2 v^i - b^0 b^i - \rho W v^i \\ v^i B^x - B^i v^x \\ v^i B^y - B^i v^y \\ v^i B^z - B^i v^z \end{pmatrix}. \quad (22)$$

Note that while system (19) consists of eight conservation equations, only seven components of the fluxes, F^i, are non-trivial. Owing to the antisymmetric character of the induction Eq. (10) the flux of B^i in i-direction is always zero. For cylindrical coordinates, the equations are given in Appendix A.

2.2.1. Spectral decomposition

Our method exploits a directional splitting algorithm to compute the numerical fluxes across cell interfaces, i.e., in each directional sweep the changes of the variables due to fluxes in the orthogonal directions are assumed to be zero. The fluxes along the sweep direction are computed by solving a one dimensional system, e.g., in the x-sweep the system reads

$$\partial_t \boldsymbol{U} + \partial_x \boldsymbol{F}^x = 0, \quad (23)$$

where $\boldsymbol{U}$ and $\boldsymbol{F}^x$ are given by Eqs. (21) and (22), respectively. Note that in the one dimensional system the flux of B^x in x-direction is zero. Thus the number of independent variables reduces from 8 to 7. As explained in detail below, the numerical integration of system (23) requires knowledge of the corresponding eigenvalues λ, which are the solutions of the characteristic polynomial

$$\det\left(\mathcal{A}^x - \lambda \mathcal{A}^t\right) = 0, \quad (24)$$

where $\mathcal{A}^x$ and $\mathcal{A}^t$ are Jacobian matrices defined as

$$\mathcal{A}^x_{ij} = \frac{\partial \boldsymbol{F}^x_i}{\partial \boldsymbol{V}_j} \quad \text{and} \quad \mathcal{A}^t_{ij} = \frac{\partial \boldsymbol{U}_i}{\partial \boldsymbol{V}_j}, \quad (25)$$

respectively. Here $i, j = 1 \ldots 7$ denote the seven (independent) components of the vectors (20), (21) and (22).

Equation (24) yields a polynomial of degree seven, the solution of which is non-trivial. Anile (1989) tackled the problem using a fully covariant formulation of the RMHD system. The new system, formed by ten evolution equations instead of seven, leads to the appearance of three additional non-physical waves that have to be discarded. The remaining seven waves are physical. The corresponding eigenvalues can be found in, e.g., Anile (1989). The speeds (eigenvalues) of the four magnetosonic (ms) waves (two slow and two fast ones, sms and fms, respectively) are given by the roots of a quartic polynomial in $\lambda_{\rm ms}$ (C.1) which does not have a simple analytic solution, i.e., it has to be solved numerically (Sect. 2.3.1).

In RMHD there exist degeneracies where system (23) is not strictly hyperbolic (for an extended discussion see, e.g., Komissarov 1999a). The two possible degenerate cases can be characterized in terms of the components of the magnetic field parallel ($b_{\rm p}$) and normal ($b_{\rm n}$) to the Alfvén-wavefronts in the fluid rest frame. Particularly, when $b_{\rm n} = 0$, the slow magnetosonic and the Alfvén waves propagate at the same speed as the entropy wave, it is possible to find an analytic solution to the eigenvalue problem Eq. (24) taking into account that, applied to the one dimensional system (23), $b_{\rm n} = 0$ in the fluid rest frame implies $B^x = 0$ in the laboratory frame, i.e., the quartic equation factors into two parts: a quadratic equation in λ whose solutions are the two fast magnetosonic waves, and a trivial factor $(v^x - \lambda)^2$. The complete set of eigenvalues is hence given by

$$\lambda_0 = v^x \quad \text{(five times degenerate)} \quad (26)$$

and

$$\lambda^{\pm}_{\rm fms} = \frac{v^x(1-\omega^2)}{1 - \boldsymbol{v}^2\omega^2 - R} \pm \frac{\sqrt{\Big((\boldsymbol{v}^2 - 1)\omega^2 + R\Big)\Big((\boldsymbol{v}^2 - (v^x)^2)\omega^2 + (v^x)^2 - 1 + R\Big)}}{1 - \boldsymbol{v}^2\omega^2 - R} \quad (27)$$

with $R = c_{\rm s}^2(\boldsymbol{v}\cdot\boldsymbol{B})^2/(\rho h^* W^2)$ and $\omega^2 = c_{\rm s}^2 + c_{\rm a}^2 - c_{\rm s}^2 c_{\rm a}^2$. In the second degenerate case the characteristic polynomial (24) does not have such a simple solution.

2.3. Numerical implementation

In this section, we will describe the main ingredients of our numerical algorithm. We apply a method of lines (e.g., LeVeque 1991) to the equations written in conservation form. This method consists of a spatial discretization step during which the numerical fluxes (Sect. 2.3.1) and the source terms are computed using different types of inter cell reconstructions of the primitive variables (Sect. 2.3.2). Subsequently, the remaining system of semi-discrete ordinary differential equations is integrated in time using a multi-step Runge-Kutta algorithm developed by Shu & Osher (1988) which provides up to third order accuracy. Our implementation also includes a second order time accurate version of the algorithm. A set of test problems which have been successfully passed by the code are presented in Appendix B.

2.3.1. Flux formula

Numerical fluxes across cell interfaces are computed using the HLL flux formula (Harten et al. 1983) similar as in

Del Zanna et al. (2003) and Gammie et al. (2003). The flux formula is based on the calculation of the maximum speed of perturbations propagating into the states left and right of the cell interface:

$$\widehat{\boldsymbol{F}}(\boldsymbol{U}_L, \boldsymbol{U}_R) = \frac{\psi_+ \boldsymbol{F}(\boldsymbol{U}_L) - \psi_- \boldsymbol{F}(\boldsymbol{U}_R) + \psi_+\psi_-(\boldsymbol{U}_R - \boldsymbol{U}_L)}{\psi_+ - \psi_-}, \quad (28)$$

where

$$\psi_+ = \max\left(0, \lambda^+_{\mathrm{fms},L}, \lambda^+_{\mathrm{fms},R}\right), \quad (29)$$
$$\psi_- = \min\left(0, \lambda^-_{\mathrm{fms},L}, \lambda^-_{\mathrm{fms},R}\right).$$

These speeds are computed from the characteristic quartic polynomial (C.1, and can effectively be obtained with a Newton-Raphson iteration scheme starting from $\lambda = \pm 1$.

Although they are presently not implemented in our code, we have explored several numerical and analytical methods (including the method described in Abramowitz & Stegun 1965 and used by Del Zanna et al. 2003) to find the four roots of the quartic Eq. (C.1). According to our tests the best procedure is (1) to compute the two fast magnetosonic wave speeds using a Newton-Raphson iteration scheme; then (2) to reduce the quartic to a quadratic equation by polynomial division; and finally (3) to obtain the two slow magnetosonic wave speeds by a combination of Newton-Raphson iteration and bisection schemes (Press et al. 1992). However, even this best procedure leads to severe numerical problems for high Lorentz factors ($W \gtrsim 40$).

In the code we use the analytical solution for the left- and right-propagating waves as given by Eq. (27) (corresponding to the first degenerate case) for $\boldsymbol{v} \cdot \boldsymbol{B} = 0$. The resulting speeds are always smaller than the speed of light and provide very good (i.e., within 1%) lower and upper bounds to the actual slowest and fastest magnetosonic wave speeds (as given by the numerical solution of the quartic), respectively. Because of the latter property (proved in Appendix C), the estimates are ideally suited to be used in the HLL flux formula (28). Our analytic estimates are more precise than those reported by Gammie et al. (2003) obtained from the solution of an approximate RMHD dispersion relation. The estimates of the maximum and minimum eigenvalue are also used to compute the Courant time step condition.

The resulting HLL algorithm (as those of Del Zanna et al. 2003; and Gammie et al. 2003) is robust and very efficient computationally. The implementation of more refined Riemann solvers or flux formulas requiring the complete set of eigenvalues and eigenvectors (like, e.g., Balsara 2001; Komissarov 1999b; or Koldoba et al. 2002) is very difficult to realize because of the presence of the RMHD degeneracies discussed above. This may change once a consistent set of eigenvectors is found which can handle both non-degenerate and degenerate states.

2.3.2. Higher order of accuracy

In order to increase the spatial order of the accuracy of the code, we have implemented two algorithms for reconstructing variables within numerical cells.

Piecewise linear method (PLM) We use a modified version of the *minmod* linear interpolation algorithm (e.g., LeVeque 1991) for the one dimensional reconstruction of the variables within cells. If a is one of the variables to be reconstructed and a_k is its value in zone k, we construct the slopes Δ_k:

$$\Delta_k = 0.5\ (\mathrm{sign}(s_+) + \mathrm{sign}(s_-))\ \min(|s_+|, |s_-|) \quad (30)$$

where

$$s_+ = \frac{a_{k+1} - a_k}{x_{k+1} - x_k}, \quad s_- = \frac{a_k - a_{k-1}}{x_k - x_{k-1}}. \quad (31)$$

Values of a at the left and right interface of the zone, $a_{L,k}$ and $a_{R,k}$, are then computed according to

$$a_{L,k} = a_k + \Delta_k(x_{k-1/2} - x_k), \quad (32)$$
$$a_{R,k} = a_k + \Delta_k(x_{k+1/2} - x_k). \quad (33)$$

The monotonicity of the data is preserved when taking the minimum of $|s_+|$ and $|s_-|$ at every numerical cell, whereas the TVD property is accomplished by switching to first order (i.e., piecewise constant data) at local extrema (where $\Delta_k = 0$).

In the x sweep the reconstruction procedure just described is applied to the variables $\{\ln\rho, \ln p, Wv^x, Wv^y, Wv^z, B^y, B^z\}$. Interpolating the logarithms of density and pressure reduces the magnitude of the jumps at cell interfaces with large gradients of density and/or pressure. Finally, and here is where the modification with respect to the standard minmod algorithm takes place, we limit the absolute value of the slope to 2 when interpolating $\ln\rho$ and $\ln p$ in those zones where the magnetization parameter β (14) is larger than a certain threshold (between 4 and 10 in our applications). Reconstruction of B^x is unnecessary in the x sweep, as the magnetic field is originally defined on a staggered grid (see Sect. 2.3.4). In the y and z sweep the primitive variables are reconstructed in a similar way, but permuting the indices (x, y, z) cyclically.

Piecewise parabolic method (PPM). From the original relativistic version of the PPM algorithm of Martí & Müller (1996) we only adopt the cell reconstruction in order to construct monotonic parabola. For each zone k the quartic polynomial is obtained, which has zone-averaged values a_{k-2}, a_{k-1}, a_k, a_{k+1} and a_{k+2}, where a is the variable to be reconstructed (the same as in the PLM algorithm). The polynomial interpolates the structure inside the zone, and provides the values at the left and right interface of the zone, $a_{L,k}$ and $a_{R,k}$. These reconstructed values are then modified such that the parabolic profile defined by $a_{L,k}$, $a_{R,k}$ and a_k is monotonic inside the zone. The modified interpolated values at the zone interfaces define local Riemann problems which are solved by Eq. (28). Near contact discontinuities the interpolation procedure is slightly modified to produce narrower jumps. In the vicinity of shocks the scheme switches (locally) to a piecewise constant approximation in order to avoid spurious post shock oscillations (Appendix I in Martí & Müller 1996). The original relativistic hydrodynamic algorithms for the detection of contact discontinuities and shocks can also be used in RMHD provided the character of the discontinuity does not change in the presence of a magnetic field: in a shock there is a jump of the thermal pressure accompanied by a negative value of the divergence of the velocity field and, in a contact discontinuity there is a jump in density without change of normal velocity.

2.3.3. Recovery of primitive variables

The code evolves the conserved quantities $\{D, \boldsymbol{S}, \tau, \boldsymbol{B}\}$, but not the primitive variables $\{\rho, \boldsymbol{v}, p, \boldsymbol{B}\}$. Therefore, an algorithm is required which computes the latter from the zone-averaged values of the former set. Since the conserved quantities can be written in terms of the primitives in an analytic closed form, but not the other way around, one has to employ iterative algorithms to compute the primitive variables.

We use Eqs. (21) and (13) to construct two functions $\mathcal{F}_1$ and $\mathcal{F}_2$ depending on the Lorentz factor W and the variable $Z \equiv \rho h W^2$ (the inertial rest mass density for an ideal gas),

$$\mathcal{F}_1(Z,W) = \frac{\tau + D}{1+\boldsymbol{B}^2} - \frac{Z+\boldsymbol{B}^2}{1+\boldsymbol{B}^2} + \frac{(\boldsymbol{B}\cdot\boldsymbol{S})^2}{2Z^2(1+\boldsymbol{B}^2)} + \frac{\gamma-1}{\gamma}\frac{Z-DW}{W^2(1+\boldsymbol{B}^2)} + \frac{\boldsymbol{B}^2}{2W^2(1+\boldsymbol{B}^2)} \tag{34}$$

and

$$\mathcal{F}_2(Z,W) = 1 - \frac{1}{W^2} - \frac{1}{(Z+\boldsymbol{B}^2)^2}\left(\boldsymbol{S}^2 + 2\frac{(\boldsymbol{B}\cdot\boldsymbol{S})^2}{Z} + \boldsymbol{B}^2\left(\frac{\boldsymbol{B}\cdot\boldsymbol{S}}{Z}\right)^2\right). \tag{35}$$

The common zero of both functions – computed with a two dimensional Newton-Raphson iteration (function `mnewt` in Press et al. 1992) – yields values for Z and W, which are used to calculate the primitive variables via

$$v^i = \frac{S^i + \frac{b^0}{W}B^i}{Z+\boldsymbol{B}^2} \quad \text{where} \quad \frac{b^0}{W} = \frac{\boldsymbol{B}\cdot\boldsymbol{S}}{Z}, \tag{36}$$

$$\rho = \frac{D}{W} \quad \text{and} \quad p = \frac{\gamma-1}{\gamma}\frac{Z-DW}{W^2}. \tag{37}$$

The internal energy ε follows then from Eq. (13).

This recovery scheme yields physical results for a wide range of parameters. Using in FORTRAN double-precision arithmetics, for an initial guess which differs from the solution by less than about 20%, a tolerance of 10^{-8}, and magnetic fields of the order of 10^{-5}, the recovery provides physical solutions (expressed in code units) for $10^{-10} < p < 10^8$, $10^{-9} < \rho < 10^8$, and $1 < W < 10^4$. For stronger fields these intervals get smaller, while for smaller fields they become larger. If no magnetic field is present, the recovery works for density and pressure values as low as 10^{-17} up to $W = 10^4$. It also works for strong magnetic fields, if $\beta < 1$, or in general if the magnetic energy density and the Poynting flux do not much exceed the internal plus kinetic energy density and the hydrodynamic momentum fluxes, respectively. The parameter range where the recovery algorithm works properly widens when better initial values are available for the iterative procedure.

2.3.4. Conservation of $\nabla \cdot B = 0$

The equations for the rest mass, momentum and energy are integrated using the numerical fluxes given in Eq. (28), while the equations for the magnetic field are evolved in a different way in order to keep $\nabla \cdot \boldsymbol{B} = 0$ during the simulation. The latter constraint is not trivially fulfilled although it is implicitly encoded in the RMHD Eqs. (19). To guarantee the constraint also numerically, we use the constrained transport (CT) method developed by Evans & Hawley (1988) and adapted to Godunov-type schemes by Ryu et al. (1998), which keeps $\nabla \cdot \boldsymbol{B} = 0$ up to machine precision, but which has the disadvantage that it is at most second order accurate in space (see, Londrillo & Del Zanna (2004) for higher order implementations of the divergence free condition).

We define two sets of magnetic field vectors (for the sake of clarity we will restrict ourselves here to the algorithm for two dimensional problems):

1. the zone centered vector $\widetilde{\boldsymbol{B}}_{k,l}$ defined at coordinates (x_k, y_l); and
2. the *staggered* interface magnetic field $\widehat{\boldsymbol{B}}_{k,l}$, where $\widehat{B}^x_{k,l}$ is defined at $(x_{k-1/2}, y_l)$ and $\widehat{B}^y_{k,l}$ at $(x_k, y_{l-1/2})$.

Note that we introduced a slight inconsistency in the notation in order to avoid formulae cluttered with indices.

The zone centered vector $\widetilde{\boldsymbol{B}}$, which is required for setting the boundary conditions, and for calculating the source terms and the fluxes of the other variables, is computed in every sweep by a simple interpolation from the staggered field components

$$\widetilde{B}^x_{k,l} = \frac{1}{2}\left(\widehat{B}^x_{k,l} + \widehat{B}^x_{k+1,l}\right). \tag{38}$$

and

$$\widetilde{B}^y_{k,l} = \frac{1}{2}\left(\widehat{B}^y_{k,l} + \widehat{B}^y_{k,l+1}\right). \tag{39}$$

The field components are updated by applying the Runge-Kutta time integration algorithm to the set of equations

$$\frac{\mathrm{d}\widehat{B}^x_{k,l}}{\mathrm{d}t} = -\frac{1}{\Delta y}(\Omega_{k,l+1} - \Omega_{k,l}) \tag{40}$$

and

$$\frac{\mathrm{d}\widehat{B}^y_{k,l}}{\mathrm{d}t} = +\frac{1}{\Delta x}(\Omega_{k+1,l} - \Omega_{k,l}), \tag{41}$$

where

$$\Omega_{k,l} = \frac{1}{2}\left(\widehat{F}^y(B^x)_{k-1,l} + \widehat{F}^y(B^x)_{k,l} - \widehat{F}^x(B^y)_{k,l-1} - \widehat{F}^x(B^y)_{k,l}\right) \tag{42}$$

corresponds to the z-component of $\boldsymbol{B}\times\boldsymbol{v}$ computed at the lower left corner of the zone centered at (x_k, y_l), while $\widehat{F}^y(B^x)_{k,l}$ and $\widehat{F}^x(B^y)_{k,l}$ are the numerical advective fluxes in y and x-direction across the interface at $(x_{k-1/2}, y_l)$ and $(x_k, y_{l-1/2})$ corresponding to B^x and B^y, respectively, i.e.,

$$\widehat{F}^y(B^x)_{k,l} = (\widehat{v^y B^x})_{k,l}$$

$$\widehat{F}^x(B^y)_{k,l} = (\widehat{v^x B^y})_{k,l}.$$

It is easy to show that Eqs. (40)–(42) conserve the following discretized version of the divergence of the magnetic field:

$$(\nabla \cdot \boldsymbol{B})_{k,l} \equiv \frac{\widehat{B}^x_{k+1,j} - \widehat{B}^x_{k,l}}{\Delta x} + \frac{\widehat{B}^y_{k,l+1} - \widehat{B}^y_{k,l}}{\Delta y}. \quad (43)$$

2.3.5. Boundary conditions

At the boundaries the computational grid is extended by up to four so-called *ghost zones* which serve to enforce the boundary conditions for the primitive variables before the spatial interpolation is performed. The number of ghost zones employed in this process depends on the order of the spatial interpolation algorithm (one for PLM, four in case for our implementation of PPM).

A second boundary routine is called after the dimensional sweeps to set the boundary values for the fluxes associated to the magnetic field components. These fluxes only need to be fixed in the first ghost zone (see Eq. (42)), e.g., when computing $\Omega_{1,1}$ one has to prescribe both $\widehat{F^y}(B^x)_{0,1}$ and $\widehat{F^x}(B^y)_{1,0}$. Usually fluxes at the ghost zone are set in such a way that zero flux gradients are enforced.

3. Simulations

3.1. Simulation setup

3.1.1. The numerical grid

All jet simulations in this work are performed on a 2D equidistant grid in cylindrical coordinates, (r, z), assuming axisymmetry. The simulations themselves are *2.5-dimensional* as we evolve all three spatial components of vectors using Eqs. (A.2)–(A.9), and thus include a toroidal magnetic field and flow velocity.

The simulated jets are produced by injecting beam matter into the grid along the z axis through a nozzle of radius r_b (the *beam radius*) at $z = 0$. Outside the nozzle, i.e., for $r > r_\mathrm{b}$ and $z = 0$ we impose special reflecting boundary conditions assuming that the axial and toroidal velocity and field components are mirrored. This is justified by the presence of a twin counterjet in actual radio galaxies, and eliminates the Lorentz force in the equatorial plane. The assumed axisymmetry implies reflecting boundary conditions on the symmetry axis at $r = 0$, where the radial and toroidal velocity and magnetic field components are mirrored, i.e., they are zero at zero radius. At the two outer edges of the grid at $r = r_\mathrm{max}$ and $z = z_\mathrm{max}$, respectively, we impose zero gradient boundary conditions. Material is allowed to freely leave the grid there.

Initially the computational domain is filled with a uniform medium at rest having the same thermal pressure and adiabatic index as the jet. The initial magnetic field configuration depends on the type of simulation.

3.1.2. Jet parameterization

Every jet simulation is fully specified by setting the following independent parameters:

- the flow speed of the beam at the nozzle, v_b (associated Lorentz factor, W_b);
- the ratio of the rest mass densities of the beam fluid and the external medium, η;
- the adiabatic index, γ, used in the equation of state (13);
- the internal (classical) Mach number of the beam at the nozzle $M_\mathrm{b} = v_\mathrm{b}/c_\mathrm{s,b}$ (which controls the thermodynamic relativistic effects of the model) or its relativistic analogue, the proper Mach number (Königl 1980), $\mathcal{M}_\mathrm{b} = M_\mathrm{b} W_\mathrm{b}/W_\mathrm{s,b}$, where $W_\mathrm{s,b} \equiv (1 - c^2_\mathrm{s,b})^{-1/2}$ is the Lorentz factor associated with the local sound speed in the beam;
- the magnetization parameter, β (Eq. (14)), both of the beam fluid and the external medium.

The other parameter, σ, defined by Eq. (15), will prove useful when discussing the simulations. However, as it is not independent of the other parameters, it need not to be specified for a jet model.

We point out here that there are three ways for a magnetized jet to be relativistic. First, when the flow velocity is close to c, and therefore the Lorentz factor of the flow is much larger than one. Second, when the beam plasma is hot, i.e., $\varepsilon_\mathrm{b} \gg c^2$ such that $h_\mathrm{b} \gg 1$ and the sound speed (17) are relativistic approaching the maximum value $\sqrt{\gamma - 1}$ ($\approx 0.58c$ for $\gamma = 4/3$, and $\approx 0.83c$ for $\gamma = 5/3$). In this case, the internal beam Mach number, M_b, approaches the minimum value for a given v_b ($1.73 v_\mathrm{b}$ for $\gamma = 4/3$, and $1.22 v_\mathrm{b}$ for $\gamma = 5/3$). Finally, when $\sigma_\mathrm{b} \gg 1$ such that $h^*_\mathrm{b} \gg 1$, and when $h^*_\mathrm{b} \gg h_\mathrm{b}$, the Alfvén speed (18) is close to c.

3.1.3. Magnetization parameter

When simulating a jet with a non-uniform magnetic field, the magnetization parameter of the beam is given by the average value, $\overline{\beta}$, across the beam. In our simulations of jets with purely toroidal magnetic fields presented below we have assumed the same profile for the magnetic field as Lind et al. (1989) and Komissarov (1999b), corresponding to a beam in transverse hydromagnetic equilibrium with a core of uniform electric current with radius r_m – the magnetization radius – and a shell without any current. The (toroidal) magnetic field in the beam, b^ϕ_b, is then given by

$$b^\phi_\mathrm{b} = \begin{cases} b_\mathrm{m} \dfrac{r}{r_\mathrm{m}} & \text{if} \quad r < r_\mathrm{m}, \\[2ex] b_\mathrm{m} \dfrac{r_\mathrm{m}}{r} & \text{if} \quad r \geq r_\mathrm{m}, \\[2ex] 0 & \text{if} \quad r > r_\mathrm{b}, \end{cases} \quad (44)$$

where b_m is the magnetic field strength at the magnetization radius r_m.

Averaging (44) over the beam cross section at the nozzle and dividing by the uniform thermal beam pressure, p_b, yields

$$\overline{\beta} = \frac{b^2_\mathrm{m} r^2_\mathrm{m} (0.25 - \ln{(r_\mathrm{m}/r_\mathrm{b})})}{p_\mathrm{b} r^2_\mathrm{b}}. \quad (45)$$

From p_b, $\overline{\beta}$ and r_m one obtains b_m through (45). Finally, the toroidal magnetic field in the beam as seen in the laboratory frame is $B^\phi_\mathrm{b} = W_\mathrm{b} b^\phi_\mathrm{b}$.

In the simulations with purely poloidal magnetic fields we have chosen a simpler initial setup. The grid is initially filled with a uniform axial magnetic field, B^z_a, which according to the definitions of $|\boldsymbol{b}|^2$ and β is given by

$$B^z_\mathrm{a} = \sqrt{2\beta p_\mathrm{b}}. \tag{46}$$

Through the nozzle the same field is injected with the flow, i.e., $B^z_\mathrm{b} = B^z_\mathrm{a}$.

3.1.4. Analysis tools

Besides the analysis and comparison of the distributions of rest-mass density, flow velocity, magnetic field lines, etc. for the different models, we introduce a number of global quantities that are used to characterize some morphodynamical aspects of the models. These quantities are:

- the *head position* of the jet is the z-coordinate of the leading bow shock, and hence the length of the cavity, l_c;
- the associated jet velocity, v_h, i.e., the change of the length of the cavity as a function of time;
- the (mean) velocity of the terminal (Mach) shock, v_M;
- the *cylindrical aspect ratio*, $A_\mathrm{c} \equiv l_\mathrm{c}/r_\mathrm{c}$, is the aspect ratio of a cylinder which has the same volume and length, l_c, as the cavity, i.e., $r_\mathrm{c} = \sqrt{V_\mathrm{c}/(\pi l_\mathrm{c})}$;
- various mean values along the axial direction defined as $\langle A\rangle = \sum_{z_k=1}^{z_k=n_z} A(r_i, z_k)/n_z$, where A is a physical quantity whose zone average value is $A(r_i, z_k)$ in zone (r_i, z_k).

As in Aloy et al. (1999a, 2000); Aloy et al. (2003), the code also evolves a variable, f, representing the beam mass fraction. In order to trace the beam material across the computational domain, this tracer variable is set to $f = 1$ for the material injected through the nozzle, and to $f = 0$ for the ambient medium. In the following, the beam mass fraction will be used to identify different parts of the jet: zones where $f \geq 0.9$ belong to the beam, and zones where $0.1 < f < 0.9$ belong to the cocoon. The cavity is defined as the region where $f > 0$. Note that cavity and cocoon are different structures according to these definitions, whereas in the literature the terms cocoon and cavity are sometimes used interchangeably.

A one dimensional estimate for the head advance speed of a non-magnetized, relativistic jet can be obtained by equating the momentum flux of the beam and the ambient gas in the frame of the working surface (Martí et al. 1997):

$$v_\mathrm{h}^{1D} = \frac{\sqrt{\eta_R^*}}{1+\sqrt{\eta_R^*}} v_\mathrm{b}, \tag{47}$$

with $\eta_R^* = \eta_R W_\mathrm{b}^2$, where $\eta_R = (\rho_\mathrm{b} h_\mathrm{b})/(\rho_\mathrm{a} h_\mathrm{a})$ is the ratio of the enthalpies of the beam and the ambient medium (subscript a denotes the ambient medium). Even though this expression does not hold for magnetized jets, it will prove to be a very good estimate, as discussed below.

The evolution of 2D jets can be divided into two phases (Scheck et al. 2002): (1) a 1D phase where the velocity is constant and approximated by (47); and (2) a 2D phase where 2D effects become important and the jet decelerates. The 2D phase begins when the first major vortex shedding occurs at the terminal Mach disk.

Table 1. Overview of the simulated models: the columns give from left to right the name of the model, the type of the initial field configuration, the adiabatic index γ, the beam speed v_b, the proper beam Mach number $\mathcal{M}_\mathrm{b}$, the average magnetization parameter $\overline{\beta}$, and the ratio of the magnetic energy density and the rest mass density σ.

Model	Field	γ	v_b	$\mathcal{M}_\mathrm{b}$	$\overline{\beta}$	σ
C2–0	none	5/3	0.99	41.95	0	0
C2–1/20	tor	5/3	0.99	41.95	1/20	0.0017
C2–1	tor	5/3	0.99	41.95	1	0.0341
C2–10/3	tor	5/3	0.99	41.95	10/3	0.1135
C2–pol-1	pol	5/3	0.99	41.95	1	0.0341
B1–0	none	4/3	0.99	41.95	0	0
B1–1/20	tor	4/3	0.99	41.95	1/20	0.0022
B1–1	tor	4/3	0.99	41.95	1	0.0445
B1–10/3	tor	4/3	0.99	41.95	10/3	0.1482
B1–pol-1	pol	4/3	0.99	41.95	1	0.0445
C1–0	none	5/3	0.9	13.61	0	0
C1–1	tor	5/3	0.9	13.61	1	0.0279
C1–10/3	tor	5/3	0.9	13.61	10/3	0.0931
C1–pol-1	pol	5/3	0.9	13.61	1	0.0279

3.2. Model parameters

We have selected three of the jet models of Martí et al. (1997) to study the effects of magnetic fields on the dynamics and morphology of relativistic jets. All three jets are light ($\eta = 0.01$), supersonic, cold ($M_\mathrm{b} = 6$), and the gas pressure is matched with that of the ambient medium. In the toroidal field models the magnetization radius is $r_\mathrm{m} = 0.6 r_\mathrm{b}$, and the magnetic field (44) is injected with the beam into a uniform, non-magnetized medium. In the poloidal field simulations, the whole grid is filled with a uniform, purely axial magnetic field of the same strength as that injected with the beam according to Eq. (46). The other parameters of the external medium are identical to those of the toroidal field models. The simulations are performed on a uniform grid of $(n_r \times n_z) = (420 \times 2000)$ zones covering a domain of 10.5×50 beam radii, i.e., the resolution is 40 zones$/r_\mathrm{b}$. We point out that this numerical resolution is equivalent to that of Martí et al. (1997), who used a third order algorithm (but only 20 zones$/r_\mathrm{b}$), while we only use a second order accurate one. Thus, as in Martí et al. (1997), the most prominent supersonic structures are properly resolved. Of course, even better numerical resolution would be required to study problems like the mass entrainment in the jet or the mixing of beam and ambient plasma in the cavity of the generated jets. The purely hydrodynamic and the toroidal field models are simulated with PLM spatial interpolation, while the poloidal field runs employ PPM interpolation (although in both cases the two spatial interpolations could have been used indistinctly; see Appendix B). All models are evolved until the head of the jet reaches the opposite edge of the computational domain.

For our models (see Table 1) we use the same model names as Martí et al. (1997) extended by suffixes describing the magnetic field strength. For example, the toroidal field model of series C2 with an average magnetization of 1 (i.e., the equipartition model) is called C2−1, and the corresponding poloidal



Fig. 1. Snapshots of the logarithm of the rest mass density of models C2−0, C2−1/20, C2−1 and C2−10/3 at $t \sim 126\, r_\mathrm{b}/c$.

field model is called C2−pol-1. The model corresponding to the pure hydrodynamic case is named C2−0. All models have $\sigma < 1$, i.e., they have a low Poynting flux. The models of series B1 and C2 only differ in the adiabatic index γ, and those of series C1 and C2 only in the beam velocity v_b. The toroidal velocity at injection is zero in all models, which together with the chosen initial magnetic field configurations prevent the models from generating additional components of magnetic field and flow velocity in the course of evolution.

Our boundary condition at $z = 0$ (reflecting outside the nozzle, see previous section) differs from that of Martí et al. (1997) (zero gradients outside the jet nozzle), as does the Riemann solver and the reconstruction procedure. Therefore, the results of our purely hydrodynamic reference simulations are slightly different from their results. We choose the reflecting boundary conditions in order to avoid that a large part of the energy contained in the back flow leaves the grid. This boundary condition is also consistent with that of more recent jet simulations, like e.g., Scheck et al. (2002) and Krause (2003).

Besides the field topology the setup of the toroidal and the poloidal field models differs in other aspects, too. In the toroidal field models B^ϕ is injected together with the beam flow into a domain filled with a non-magnetized medium. In the poloidal case, however, the computational domain initially already contains a magnetic field of the same strength as that injected with the beam. Hence, in the poloidal field models both the total pressure p^* (as in the toroidal field models) and the thermal pressure p are matched at the beam surface, p being uniform and the same inside the beam and in the ambient medium. In contrast, the toroidal field models have a negative radial gradient in p^*, which is compensated by the magnetic tension.

All jet simulations were carried out on an IBM p690 Regatta computer and required between 8 and 30 h on a shared memory node with 32 Power4 1300 MHz processors.

4. Results

The most remarkable result we find is that different from classical MHD simulations (e.g., Lind et al. 1989) the inclusion of a magnetic field leads to diverse effects which do not always scale linearly with the relative strength of the field. There are, however, some clear trends which we will discuss in detail below.

4.1. Series C2

4.1.1. C2–0 to C2–10/3: Purely hydrodynamic and toroidal field models

Figures 1 and 2 show the rest mass density and the Lorentz factor of the purely hydrodynamic model C2−0, and those of the toroidal field models C2−1/20, C2−1 and C2−10/3 (from top to bottom), respectively. Apart from differences in details, one morphological feature is particularly noticeable: with growing toroidal field strength, the supersonic part of the beam ends further behind the leading edge of the bow shock (note the steep decline of the Lorentz factor near the head of the jet in Fig. 2), i.e., the jet becomes transonic much closer to the origin. The high density and pressure structure between the termination of the supersonic beam at the Mach disk and the head of the jet is called *nose cone* (Clarke et al. 1986) or *plug* (Lind et al. 1989) (e.g., the region around the axis between 28 and 46 beam radii in the bottom panel of Fig. 1). The nose cone is made up of beam material that it is not deflected backwards into the



Fig. 2. Same as Fig. 1 but showing snapshots of the Lorentz factor and the flow field.

cocoon, and of entrained ambient material near its rear end. Both the density and the pressure in the nose cone are very high, and show indications of turbulence. Moreover, the pressure in the nose cone is relatively smooth and particularly high on the axis (due to magnetic confinement) where it can reach values $\sim 10^3$ times larger than that of the external medium. The nose cone makes the head of the jet appear narrower for larger values of $\overline{\beta}$.

In the cocoon, close to the head of the jet, a number of organized structures appear as a result of the dynamic effects of the toroidal field. The magnetic field confines and suppresses the development of Kelvin-Helmholtz instabilities at the interface cocoon/shocked ambient medium, and the material deflected into the cocoon after passing the Mach disk starts to fill a low density bubble. With increasing toroidal field strength the bubble grows larger in size. In model C2−10/3 it roughly stretches from $z = 20\, r_\mathrm{b}$ up to $z = 35\, r_\mathrm{b}$, and is confined by a shell of highly magnetized material (Fig. 3). The Mach disk and the terminal shock of the beam reside in the middle of the bubble. The beam flow is terminated at the Mach disk, and matter has to flow around it to reach the head of the jet (Fig. 2). In model C2−1 the shocked flow around the Mach disk reaches relativistic speeds with Lorentz factors of 3 and more. The average beam Lorentz factor decreases with increasing magnetization of the beam.

In models C2−1 and C2−10/3 the magnetization β (Fig. 3) decreases drastically across the first cross shock due to an increase in the thermal pressure (Fig. 4). It remains low inside the beam up to the terminal shock, increases downstream of this shock, and becomes even larger when the shocked plasma begins to flow away radially from the jet axis. The pinching force provided by the toroidal field in that region suppresses any further sideways expansion of the jet, and pushes a large fraction of the plasma into the nose cone. In model C2−10/3 this pinching is strongest at $z = 38\, r_\mathrm{b}$ (see the bottom panel of Fig. 2). The nose cone contains primarily highly magnetized plasma which leads to a strong pinching force confining the flow close to the axis. The magnetic field accumulates at the inside of the high density shell (the *shroud* of Lind et al. 1989) of the jet, where the thermal pressure is low and β is large (Fig. 3). This magnetized shell will influence the emission properties of the radio lobes in actual sources, as it contains the largest and most ordered return current (Fig. 4, top panel), which occurs because there is no confining field in the ambient medium, and because this environment behaves as a perfect conductor (Begelmann et al. 1984). However, the structure of the return current becomes highly anisotropic throughout the cocoon and at the surface of the jet. The magnetic field itself is largest in the beam and in the nose cone (see the two bottom panels of Fig. 3).

4.1.2. C2−pol-1: Poloidal field model

Figure 5 shows the rest mass density, the Lorentz factor with the flow field, the logarithm of the modulus of the magnetic field with the field lines superimposed, and the magnetization parameter β for the poloidal field model C2−pol-1. Although the overall morphology and structure of the jet and the cocoon of this model are very similar to those of other models of series C2, there are a number of striking differences. In particular, we will compare model C2−pol-1 both with the hydrodynamic model C2−0 and the toroidal field model having the same magnetization (C2−1).

First of all, instead of forming a narrow nose cone structure, the beam broadens close to its termination point giving the head of the jet a hammer-like appearance caused by the emission of vortices from the head of the jet. The beam plasma flows along



Fig. 3. Snapshots of the magnetization parameter β (*upper two panels*) and the toroidal field component B^{ϕ} (*lower two panels*) of the two most strongly magnetized models of series C2 at $t \sim 126\, r_{\mathrm{b}}/c$.



Fig. 4. Snapshots of the logarithm of the pressure for all the models having a toroidal field with $\overline{\beta} = 10/3$. The overlaid vector field shows the structure of the currents ($\nabla \times \boldsymbol{B}/|\nabla \times \boldsymbol{B}|$).

the field lines up to the head of the jet, where it is deflected to form a cocoon (second panel of Fig. 5) dragging along the magnetic lines.

Model C2−pol-1 also differs from all the other models in the strength of the cross shocks on the axis. While the first cross shock in model C2−pol-1 occurs much nearer to the nozzle (at $z = 8\, r_{\mathrm{b}}$) than in model C2−0, it is also much weaker (the density contrast is ~8 and ~3 times smaller than in models C2−10/3 and C2−1, respectively) and has the least extended post-shock region (in axial direction) of all models. Downstream of this shock, the beam pressure and density remain low (below $\sim 5 p_{\mathrm{b}}$) until the flow reaches the second cross shock, which is weak compared to those of the other models, too. It is especially weak compared to the toroidal field model with the same average β. This behavior results from the increase of the radially (outward) directed magnetic tension force which straightens the poloidal field lines, which are bent towards the axis due to external radial pressure forces and thereby repel waves driven into the beam. Thus, during the whole evolution the beam remains much less affected by shocks than in the models with toroidal magnetic fields. The terminal shock, however, is as strong as in model C2−1 or



Fig. 5. Snapshots of the poloidal field model C2−pol-1 at $t \sim 126\, r_b/c$. *From top to bottom*: rest mass density, Lorentz factor with flow field, logarithm of the modulus of the magnetic field with field lines superimposed, and magnetization parameter β. The magnetic field lines shown in the third panel are placed randomly, but weighted with the local field strength, such that on average their density reflects the field strength in the plane where the integration is started ($z = 50\, r_b$).

C2−0. Furthermore, the value of β decreases with increasing axial distance along the beam in model C2−pol-1 (Fig. 5) while it is rather uniform in model C2−1.

The cocoon is also much more homogeneous in model C2−pol-1 than in the other models of series C2, which is especially apparent when comparing the rest mass density snapshots in Figs. 1 and 5. The poloidal field model shows much less turbulent structure, and the back flow of beam material through the cocoon is almost straight (Fig. 5, third panel) indicating that the poloidal magnetic field reduces or completely suppresses the development of Kelvin-Helmholtz instabilities at the surface between the cocoon and the shroud. This is very different in the toroidal field and purely hydrodynamic cases, where the back flow is less ordered.

In model C2−pol-1 the magnetic field is almost completely expelled from the cocoon. The third panel of Fig. 5 shows that the absolute value of the field strength remains very low. The narrow dark filaments in the cavity interior of model C2−pol-1 (Fig. 5, third panel from top) trace the magnetopauses, i.e., the surfaces where the magnetic field changes direction and hence is zero. The magnetopauses are surrounded by areas of high magnetic field strength giving rise to the extended, twisted filaments. The magnetization parameter (Fig. 5, bottom panel) is nearly zero everywhere within the jet, because the magnetic field is advected out the cavity by the fluid flow, and accumulates in the shroud with its large thermal pressure yielding a low value of β (compared to e.g., model C2−1). This differs from the behavior of models with toroidal magnetic fields where the magnetic tension prevents a strong reduction of the magnetic field in the cavity.

4.1.3. C2: Temporal evolution

The top panel of Fig. 7 displays the motion of the jet's head for all models of series C2 and a straight line corresponding to the 1D velocity estimate given by Eq. (47). Apart from model C2−10/3 the jet propagates at essentially the same speed in all models. The 1D estimate gives a very good approximation to the speed in the 1D phase, and becomes an upper bound in the 2D phase. The small differences between the analytic estimate for non-magnetized jets and the actual simulation values are due to the fact that the jets are cold. Even the equipartition magnetic energy density (model C2−1) is still very small compared to the ram pressure at the head of the jet. Remarkably, there is no simple dependence of the propagation speed on the magnetic field strength: in model C2−10/3 the jet propagates fastest, in model C2−1 slowest, in model C2−1/20 second fastest, and the remaining models fall in between. The propagation speed of magnetized jets is determined by a number of competing effects related to both the strength and the topology of the magnetic field. However, our simulations do not show a clear trend with these two parameters, i.e., the speed depends on both parameters nonlinearly. The high speed of model C2−10/3 can be explained by its large nose cone.

The speed of propagation of the Mach disk, v_M (which corresponds, approximately, to the speed of propagation of the jet without considering the nose cone) tends to decrease with increasing magnetization (Table 2). In the hydrodynamic model C2−0, v_M is slightly smaller than in the weakly magnetized model. This trend also holds for the models of series B1, but not for those of series C1.



Fig. 6. Average thermal pressure $\langle p \rangle \equiv \sum_{z_k=1}^{z_k=n_z} p(r_i, z_k)/n_z$ measured in code units (*top row*) and average of the radial component of the net force $\langle F_r \rangle \equiv \sum_{z_k=1}^{z_k=n_z} [-\nabla p(r_i, z_k) + (\nabla \times B) \times B]_r / n_z$ (*bottom row*) versus radius for the models of all series. The snapshots are computed at $t \approx 126\, r_b/c$, $\approx 80\, r_b/c$ and $\approx 249\, r_b/c$ for series C2, B1 and C1 (*from left to right*), respectively. For models with zero magnetic field and with poloidal fields, the net force is almost zero, and hence the corresponding lines show almost no variation (around zero) at the scale used in the plots in the bottom row.

Table 2. Propagation velocity of the Mach disc (v_M) for models with a toroidal magnetic field.

Model	v_M
C2−0	0.32
C2−1/20	0.33
C2−1	0.29
C2−10/3	0.29
B1−0	0.59
B1−1/20	0.61
B1−1	0.49
B1−10/3	0.42
C1−0	0.10
C1−1	0.10
C1−10/3	0.10

Figure 8 shows the cylindrical aspect ratio of all models as a function of time. As the jet reaches the radial grid boundary in models C1 and C2 before the end of the simulation, the aspect ratio approaches that of the grid, which is about 5. This and the fact that the differences among the models are smaller than those for a computational domain containing the whole cavity, makes one to interpret these plots with caution. Nevertheless, our explanation for the high propagation speed of model C2−10/3 is further supported by its large aspect ratio. The other models do not show any trend.

The average thermal pressure $\langle p \rangle$ (defined as $\langle p \rangle \equiv \sum_{z_k=1}^{z_k=n_z} p(r_i, z_k)/n_z$) of the more strongly magnetized models (poloidal or toroidal) of series C2 is up to 4 times larger inside the beam at $t \approx 126\, r_b/c$ than in the corresponding hydrodynamic model C2−0 (Fig. 6). The pressure distributions show a distinct maximum at the beam axis instead of being flat there and leveling off at larger radii as predicted by Begelmann et al. (1984). This difference arises because in our simulations the magnetization radius r_m is smaller than the jet radius ($r_m = 0.37\, r_b$), while Begelmann et al. (1984) assume a jet with a uniform current along the whole beam, i.e., $r_m = r_b$. Our simulations further show that the average thermal pressure $\langle p \rangle$ can be dominated by the contributions of the hotspot and/or the nose cone, while Begelmann et al. (1984) exclude these regions



Fig. 7. Motion of the head of the jet for all computed models. The solid lines correspond to the 1D velocity estimate given by Eq. (47).



Fig. 8. Temporal change of the cylindric aspect ratio A_c for all models.

from their analysis. Just outside the beam ($1 \lesssim r \lesssim 2$) the radial distribution of $\langle p \rangle$ is quite flat in all models of series C2 (except of model C2−10/3), and declines at larger radii ($r \gtrsim 2$). Thus, the jets have a core-sheath structure in radial direction. The radial pressure gradient is larger in model C2−10/3 than in any other model of series C2 and seems to decrease monotonically with decreasing magnetization. Beyond some typical radius ($\approx 4\, r_b$) $\langle p \rangle$ as a function of r becomes very similar (almost flat) for all the models. Also the thermal pressure gradients (∇p) approach to zero beyond $\approx 4\, r_b$. The behavior of model C2−pol-1 is different from that of the models transporting toroidal fields. Its average thermal pressure is smaller than that in the hydrodynamic model C2−0 at all radii.

The high compression of the beam in models carrying a toroidal magnetic field is the result of the imposed magnetic field structure in the beam (Eq. (44)). Up to the magnetization radius r_m, b^{ϕ} grows linearly with radius, i.e., the pressure necessary to keep hydromagnetic equilibrium for $r < r_m$ is (e.g. Lind et al. 1989)

$$p(r) = \left[\alpha + 2\beta_m \left(1 - \frac{r^2}{r_m^2}\right)\right] p_e, \tag{48}$$

where α is a constant, and $\beta_m \equiv (b_m W_m)^2/(8\pi p_e) = (1-\alpha)(r_b/r_m)^2$. Hence, the pressure distribution has its maximum at $r = 0$.

Models C2−1/20, C2−1 and C2−10/3 which transport toroidal magnetic fields are confined by the toroidal fields in the beam and in the cocoon. In the cocoon the density profile rearranges itself in such a way that the toroidal field decreases nearly linearly with radius (note that this scaling does not hold inside the magnetization radius r_m). The radial distribution of the axial average of the net radial force $\langle F_r \rangle \equiv \sum_{z_k=1}^{z_k=n_z} [-\nabla p(r_i, z_k) + (\nabla \times B) \times B]_r / n_z$ (Fig. 6) displays two distinct regions in case of models with toroidal fields (for model C2−pol-1 $\langle F_r \rangle \approx 0$): (1) an internal (*contractive region* up to $\approx 4\, r_b$) where $\langle F_r \rangle < 0$ and where the gas in the beam and its



Fig. 9. Snapshots of the logarithm of the rest mass density of models B1−0, B1−1/20, B1−1 and B1−10/3 at $t \sim 80\, r_{\rm b}/c$. For model B1−1/20 the jet is about to leave the computational grid because of its large propagation speed.

neighborhood is pinched; and (2) an external (*expansive region*), where $\langle F_r \rangle > 0$ because the thermal pressure gradient dominates the magnetic tension force.

4.2. Series B1

4.2.1. B1–0 to B1–10/3: Purely hydrodynamic and toroidal field models

The results of the non-magnetized and toroidal field models of the series B1 are displayed in Figs. 9 and 10. The rest mass density (Fig. 9) shows some of the trends of series C2, however the effects caused by the toroidal magnetic field do not scale linearly with the magnetic field strength. The weakly magnetized model B1−1/20 behaves quite differently both from the equipartition model B1−1 and the strongly magnetized model B1−10/3. The more magnetized models develop nose cones preceded in the case of model B1−10/3 by a magnetically confined bubble. The rest mass density (Fig. 9) and the pressure (not shown) distributions show much more structure for these two more strongly magnetized models than that of the non-magnetized model. The corresponding distributions of model B1−1/20 are very smooth suggesting that its weak magnetic field inhibits the formation of structures (see Sect. 5).

The first recollimation shock shows the same trend as in series C2. Due to magnetic pinching the shock is stronger and is located closer to the nozzle for a stronger toroidal field. The strong pinching of the beam in models B1−1 and B1−10/3 leads to a widening of the beam downstream of the recollimation shocks and to smaller Lorentz factors ($W \sim 5$; Fig. 10). In model B1−1/20 the pinching is very weak, and no difference to the non-magnetized model can be recognized. Instead, it is even better collimated than model B1−0, and the flow remains highly relativistic up to the terminal shock of the beam, where the beam plasma is deflected and flows backwards smoothly in a very thin layer around the beam. This behavior explains why the jet of model B1−1/20 has propagated further than those of the other models in the same time. These results imply, that with the right combination of parameters, a small toroidal field helps to collimate the beam without producing too much pinching, which helps to keep the jet stable.

Figure 11 shows the magnetization β and B^ϕ for the two models B1−1 and B1−10/3, respectively. As for the corresponding C2 models the initial magnetization of the beam is reduced very much in the first cross shock. It is strongest near the beam's terminal shock, where the plasma flows radially away from the axis leading to the formation of the nose cone. Apart from the fact that the structures are much narrower than in the models of series C2, the overall distributions of β and B^ϕ are very similar everywhere in the cavity. β is lowest near the axis and then increases until it reaches its maximum value in the magnetic shell around the beam. B^ϕ remains largest inside the beam and the nose cone. Different from series C2, an increase of the magnetic field strength yields a more turbulent cocoon in case of the models of series B1.

4.2.2. B1–pol-1: Poloidal field model

Figure 12 shows snapshots of the rest mass density, the Lorentz factor, the magnetic field strength and the magnetization parameter of model B1−pol-1. Most of the features found in model C2−pol-1 are also present here: the jet as a whole is very smooth both in density and pressure, the cocoon showing only an indication of an eddy. The beam is almost undisturbed,



Fig. 10. Same as Fig. 9 but showing snapshots of the Lorentz factor and the flow field.



Fig. 11. Snapshots of the magnetization parameter β (*upper two panels*) and the toroidal field component B^{ϕ} (*lower two panels*) of the two most strongly magnetized models of series B1 at $t \sim 80\, r_{\mathrm{b}}/c$.

and the cross shocks are very weak. The back flow is confined to a narrow region directly around the beam and is absolutely straight. The hammer-like structure at the head of the jet seen in model C2−pol-1, which is associated with the emission of vortices, is not present in model B1−pol-1.

The smoothness of the cocoon results from the presence of the poloidal field (see the top panel of Fig. 12), which is smooth throughout the cavity. The field is only twisted in the narrow low density sheet around the beam, and even there, the twist is mostly aligned with the axis. This does not favor the formation of eddies, which could drive strong shocks into the beam. As in model C2−pol-1, $\beta \approx 10^{-6}$ in a large part of the cavity, which however, does not diminish the effect of the poloidal field on the morphology of the jet. Model B1−pol-1 shows a smoother