

# PROGENITOR-EXPLOSION CONNECTION AND REMNANT BIRTH MASSES FOR NEUTRINO-DRIVEN SUPERNOVAE OF IRON-CORE PROGENITORS

MARCELLA UGLIANO<sup>1</sup>, HANS-THOMAS JANKA<sup>1</sup>, ANDREAS MAREK<sup>1</sup>, AND ALMUDENA ARCONES<sup>2,3</sup>

*Draft version May 17, 2012*

## ABSTRACT

We perform hydrodynamic supernova simulations in spherical symmetry for over 100 single stars of solar metallicity to explore the progenitor-explosion and progenitor-remnant connections established by the neutrino-driven mechanism. We use an approximative treatment of neutrino transport and replace the high-density interior of the neutron star (NS) by an inner boundary condition based on an analytic proto-NS core-cooling model, whose free parameters are chosen such that explosion energy, nickel production, and energy release by the compact remnant of progenitors around  $20 M_{\odot}$  are compatible with Supernova 1987A. Thus we are able to simulate the accretion phase, initiation of the explosion, subsequent neutrino-driven wind phase for 15–20 s, and the further evolution of the blast wave for hours to days until fallback is completed. Our results challenge long-standing paradigms. We find that remnant mass, launch time, and properties of the explosion depend strongly on the stellar structure and exhibit large variability even in narrow intervals of the progenitors’ zero-age-main-sequence mass. While all progenitors with masses below  $\sim 15 M_{\odot}$  yield NSs, black hole (BH) as well as NS formation is possible for more massive stars, where partial loss of the hydrogen envelope leads to weak reverse shocks and weak fallback. Our NS masses of  $\sim 1.2$ – $2.0 M_{\odot}$  and BH masses  $> 6 M_{\odot}$  are compatible with a possible lack of low-mass BHs in the empirical distribution. Neutrino heating accounts for SN energies between some  $10^{50}$  erg and  $\sim 2 \times 10^{51}$  erg, but can hardly explain more energetic explosions and nickel masses higher than 0.1–0.2  $M_{\odot}$ . These seem to require an alternative SN mechanism.

*Subject headings:* supernovae: general — stars: neutron — stars: massive — stars: evolution

## 1. INTRODUCTION

One of the main goals of the long-standing quest for a better understanding of the supernova (SN) explosion mechanism is the establishment of a theoretical connection between the properties of progenitor stars and those of SNe and their remnants. Understanding the progenitor-remnant systematics would not only mean a better definition of SNe as the end point of massive-star evolution, it could also serve as input for continuative theoretical studies like binary population synthesis calculations. Moreover, it would provide indispensable information and the theoretical basis for deducing constraints from observations on the physical processes that trigger and power the onset of the explosion at the center of the SN. In particular, the explosion mechanism plays a crucial role for determining the energetic and dynamical properties of the SN blast wave. It thus controls the conditions for explosive nucleosynthesis and is of central importance for answering long-standing astrophysical questions: Which progenitors give birth to neutron stars (NSs), which ones to black holes (BHs)? How do SN properties depend on the progenitor metallicity? Are the observed masses of compact remnants compatible with modeling results? Do observed SNe provide evidence for more than one explosion mechanism? In particular, considering neutrino heating as the most widely favored process to trigger the explosions, the specific question arises, which range of SN energies and nickel masses can be explained by the neutrino-driven mechanism.

A popular way to model the blast wave initiation for study-

ing SN nucleosynthesis and determining compact remnant masses are piston-driven explosions (e.g., Woosley & Weaver 1995; Zhang et al. 2008, and references therein), in which a piston is placed at a chosen position in the pre- or post-collapse stellar core and moved with a prescribed, time-dependent velocity to mimic the infall of the selected mass shell and the explosive expansion of the newly formed SN shock. Besides the initial mass cut—fallback of nonescaping material may shift the final mass cut—also the energy deposited by the piston is prescribed as a free parameter. Alternatively, internal energy “bombs” have been used, in which a defined amount of thermal energy is released by an artificial increase of the temperature in a certain volume (Aufderheide et al. 1991).

Investigating spherically symmetric stellar core collapse by hydrodynamic simulations with simplified neutrino treatment, O’Connor & Ott (2011) identified a single parameter, the “progenitor bounce compactness”  $\xi_{2.5}$  of the innermost  $2.5 M_{\odot}$  of a star, whose value above a certain threshold they considered to be indicative for an unlikely explosion and thus BH formation. Progenitors with  $\xi_{2.5} \gtrsim 0.45$  were found to reach the BH formation limit typically within  $\lesssim 0.8$  s after bounce because of their tremendous mass accretion rates. Such stars also turned out to require a particularly high time-averaged neutrino-heating efficiency to develop an explosion against the huge ram pressure of the infalling core material. Nevertheless, the exact value of  $\xi_{2.5}$  where the bifurcation of the behavior occurs, appears somewhat arbitrarily chosen and not supported by strong theoretical arguments. In particular, the BH formation limit could already take place at a lower value of  $\xi_{2.5}$ .

Belczynski et al. (2011) and Fryer et al. (2012) made theoretical predictions of the NS/BH mass distribution by applying a simple analytic concept for estimating the explo-

<sup>1</sup> Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str. 1, D-85748 Garching, Germany

<sup>2</sup> Institut für Kernphysik, Technische Universität Darmstadt, Schlossgartenstr. 2, D-64289 Darmstadt, Germany

<sup>3</sup> GSI Helmholtzzentrum für Schwerionenforschung GmbH, Planckstr. 1, D-64291 Darmstadt, Germany

sion time, explosion energy, and fallback mass on the basis of structural properties of the progenitor stars (Fryer 2006). Fryer & Kalogera (2001) derived theoretical BH mass distributions by employing the explosion energy as function of the progenitor mass from a fit to results of a small set of four 2D models of neutrino-driven explosions (Fryer 1999). They considered different fallback masses in dependence on the (parametrized) fraction of the energy released by the SN mechanism that is used to unbind the outer layers of the progenitor stars.

In this work we adopt a different approach. We perform hydrodynamical simulations in spherical symmetry for a large set of 101 progenitor models, following core collapse and bounce, postbounce accretion, the possible onset of an explosion and subsequent neutrino-driven wind phase of the cooling proto-neutron star (PNS), and the development of the SN blast wave over hours to days until the fallback mass is determined. In order to power the explosions, we refer to the neutrino-driven mechanism, whose viability seems to be supported by recent 2D simulations with detailed energy-dependent neutrino transport (Marek & Janka 2009) and general relativity (Müller et al. 2012, Müller et al., in preparation), although an ultimate robustness of the model predictions may require 3D simulations and may hinge on remaining uncertainties of the high-density neutrino opacities and equation of state (EoS). Since in our spherical simulations explosions by neutrino heating must be triggered artificially, we excise the high-density core of the nascent NS and replace it by a time-dependent, analytic cooling model. This core model yields the time evolution of the neutrino emission from the high-density interior of the PNS. Its free parameters are adjusted such that observational characteristics of SN 1987A, i.e., the explosion energy,  $E_{87A} \sim 1.3 \times 10^{51}$  erg, and ejected  $^{56}\text{Ni}$  mass,  $M_{\text{Ni},87A} \sim 0.07 M_{\odot}$ , are reproduced for progenitor stars around  $20 M_{\odot}$  (e.g., Tanaka et al. 1999; Utrobin & Chugai 2011). This implies the assumption that SN 1987A was an ordinary case of a neutrino-driven explosion in which, e.g., exceptionally rapid core rotation did not play a crucial role for the collapse dynamics. The NS core model with the same parameter values is then also applied to all other collapsing stars.

Provided SN 1987A was such a normal case and neutrino-energy deposition was responsible for its explosion, our approach thus allows us to explore the response of other progenitor stars to the input of neutrino energy released from their forming compact remnants, and in the case of successful explosions to determine the blast wave and remnant properties of the stars. In the following we will give more information on our modeling strategy and tools, present our results, and discuss the implications.

## 2. NUMERICAL SETUP AND INPUT

We employ the explicit, finite-volume, Eulerian multi-fluid code PROMETHEUS (Fryxell et al. 1989) to solve the 1D hydrodynamics equations on a spherical grid. We use typically 1000 geometrically spaced radial zones with the outer boundary at a radius  $R_{\text{ob}}$ . The latter is placed at a large distance (initially at 150,000 km) that is not directly affected by the simulated stellar collapse and SN explosion. In particular it is sufficiently large to prevent the SN shock from leaving the computational domain during the simulated time. The advection of nuclear species is treated by the Consistent Multi-fluid Advection (CMA) scheme of Plewa & Müller (1999), and self-gravity of the stellar plasma includes corrections of the New-

tonian gravitational potential for general relativistic effects as described in detail in Scheck et al. (2006) and Arcones et al. (2007).

The inner core of the PNS with a mass of  $M_c = 1.1 M_{\odot}$  and densities well above those of the neutrinospheric layer is excised and replaced by a point mass at the coordinate origin. The shrinking of the PNS is mimicked by a retraction of the closed inner boundary at  $R_{\text{ib}}$  together with the radial grid, applying the function defined in Arcones et al. (2007). For the simulations discussed in this paper we use the values  $R_{\text{ib}}^i = R(1.1 M_{\odot} \text{ at } t \sim 10 \text{ ms p.b.}) \sim 60 \text{ km}$ ,  $R_{\text{ib}}^f = 20 \text{ km}$ , and  $t_{\text{ib}} = 0.4 \text{ s}$  for the initial and final boundary radii and the exponential contraction timescale, respectively. The chosen values yield a boundary contraction similar to the “standard case” of Scheck et al. (2006). Hydrostatic equilibrium is assumed at the boundary radius  $R_{\text{ib}}$  between excised core and accretion envelope of the nascent NS.

The cooling of the PNS interior and corresponding boundary luminosities are described by the neutrino core-cooling model presented in Sect. 2.1. The temperatures of the spectra of inflowing neutrinos (assumed to have Fermi-Dirac shape) are set equal to the gas temperature  $T_i$  in the innermost grid zone,  $T_{\nu}(R_{\text{ib}}) = T_i$ , consistent with the fact that neutrinos in highly optically thick regions are close to local thermal equilibrium. The spectral degeneracy parameters are chosen to be  $\eta_{\nu_e} = \eta_{\bar{\nu}_e} = \eta_{\bar{\nu}_x} = 0$  for electron neutrinos  $\nu_e$ , electron antineutrinos  $\bar{\nu}_e$ , and heavy-lepton neutrinos  $\nu_x$  ( $= \nu_{\mu}, \bar{\nu}_{\mu}, \nu_{\tau}, \bar{\nu}_{\tau}$ ), respectively. (Using the values for local chemical equilibrium for  $\eta_{\nu_e}$  and  $\eta_{\bar{\nu}_e}$  does not cause any significant differences.) Neutrino transport and neutrino-matter interactions in the computational domain are approximated as in Scheck et al. (2006) by radial integration of the one-dimensional (spherical), grey transport equations for neutrino number and energy, assuming the neutrino spectra to have Fermi-Dirac shape with a calculated local temperature  $T_{\nu}$ .

The high-density plasma is described by the tabulated microphysical equation of state (EoS) of Janka & Müller (1996) including arbitrarily degenerate and arbitrarily relativistic electrons and positrons, photons, and four predefined nuclear species (neutrons, protons, alpha particles, and a representative Fe-group nucleus) in nuclear statistical equilibrium (NSE). At temperatures and densities below NSE the thermodynamics are treated by the Helmholtz-EoS of Timmes & Swesty (2000).

The NS cooling and its neutrino emission are followed for 8–25 s until the shock has crossed  $10^{10}$  cm. At that time the power of the neutrino-driven wind, which pushes the dense ejecta shell enclosed by the expanding SN shock and the wind-termination shock, has decayed to a dynamically insignificant level. The models are then mapped to a grid extending to larger radii, and the inner boundary is moved from below the neutrinosphere to  $10^9$  cm in order to ease the time step constraint. The gravity potential of the central object is thereby corrected for the gas mass in the larger excised region, and free-outflow conditions are applied at the new inner boundary. When the SN blast is followed beyond shock breakout for several days, the star is embedded in a wind-like medium with a density  $\rho \propto r^{-2}$  and a constant temperature of  $T = 1100 \text{ K}$ , and the grid is shifted outward once again to cover radii from  $R_{\text{ib}} = 10^{10}$  cm to  $R_{\text{ob}}$  up to  $10^{15}$  cm.

During the long-time simulations of the SN dynamics we follow approximately the explosive nucleosynthesis by solving a small alpha-reactions network, similar to the network

described in Kifonidis et al. (2003). It consists of the 13  $\alpha$ -nuclei:  $^4\text{He}$ ,  $^{12}\text{C}$ ,  $^{16}\text{O}$ ,  $^{20}\text{Ne}$ ,  $^{24}\text{Mg}$ ,  $^{28}\text{Si}$ ,  $^{32}\text{S}$ ,  $^{36}\text{Ar}$ ,  $^{40}\text{Ca}$ ,  $^{44}\text{Ti}$ ,  $^{48}\text{Cr}$ ,  $^{52}\text{Fe}$ ,  $^{56}\text{Ni}$ , and an additional tracer nucleus, which is produced via the reaction rate for  $^{52}\text{Fe}(\alpha,\gamma)^{56}\text{Ni}$  within grid cells whose electron fraction  $Y_e$  is below 0.49. This allows us to keep track of element formation in regions with neutron excess. The network is solved in grid cells whose temperature is between  $10^8$  K and  $7 \times 10^9$  K. We assume that all nuclei are photo-disintegrated to  $\alpha$ -particles at  $T > 7 \times 10^9$  K. We also neglect feedback from the network composition to the EoS and thus to the hydrodynamic evolution.

### 2.1. Proto-neutron star core model

The cooling of the PNS core is described by an analytic model that couples the excised core region to the surrounding accretion layer, whose evolution is followed on the computational grid. We assume the dense PNS core with mass  $M_c$  and radius  $R_c$  to be approximately homogeneous and its EoS to be an ideal  $\Gamma$  law,  $P = (\Gamma - 1)e$  (with  $P$  being the pressure,  $e$  the internal energy density). Combining the total core energy,  $E_c = E_g + E_i$ , and the virial theorem,  $E_g + 3(\Gamma - 1)E_i + S = 0$ , we can replace the integrated internal energy,  $E_i$ , and express  $E_c$  in terms of the Newtonian gravitational energy,  $E_g = -\frac{2}{5}GM_c^2/R_c$ , and the surface term,  $S = -4\pi R_c^3 P_s$ , for pressure  $P_s$  at  $R_c$ :

$$E_c = \frac{3\Gamma - 4}{3(\Gamma - 1)} E_g - \frac{S}{3(\Gamma - 1)}. \quad (1)$$

The energetic evolution of the core is given by its loss of neutrino energy and compression work done on its surface as

$$\dot{E}_c \equiv \frac{dE_c}{dt} = -L_{\nu,c} - 4\pi P_s R_c^2 \dot{R}_c, \quad (2)$$

where  $\dot{E}_c$  can be computed as time derivative of Eq. (1),  $L_{\nu,c}$  is the total neutrino luminosity, and the second term results from the time derivative of the core volume. Instead of setting  $P_s$  equal to the boundary pressure on the hydro grid, we prefer to link it to overall properties of the accretion layer. This prescription is intended to capture the nature of the core-mantle coupling but not to constrain the freedom to tune the parameters of the simple PNS core model. We therefore consider hydrostatic equilibrium in terms of the mass coordinate  $m(r)$ ,  $dP/dm = -GM/(4\pi r^4)$ , and linearize both sides to obtain

$$P_s = \zeta \frac{GM_c m_{\text{acc}}}{4\pi R_c^4}, \quad (3)$$

where  $m_{\text{acc}}$  is the mass of the accretion layer that surrounds the PNS core,  $\zeta > 0$  a numerical factor of order unity, and we assumed  $P_0 \ll P_s$  for the pressure  $P_0$  outside of the accretion layer. Moreover, in performing the time derivative of Eq. (1) we assumed  $M_c$  and  $\Gamma$  to be constant. Combining Eqs. (1)–(3) we arrive at

$$L_{\nu,c} = \frac{3\Gamma - 4}{3(\Gamma - 1)} (E_g + S) \frac{\dot{R}_c}{R_c} - \frac{\zeta}{3(\Gamma - 1)} \frac{\delta E_{\text{acc}}}{\delta t} \quad (4)$$

with  $S = -\zeta GM_c m_{\text{acc}}/R_c$  and  $\delta E_{\text{acc}}/\delta t \equiv GM_c \dot{m}_{\text{acc}}/R_c$ . While the first term on the rhs of Eq. (4) describes the luminosity increase due to the deepening of the gravity potential and surface work in the case of PNS compression, the second term accounts for the higher core pressure (and internal energy) needed when the accretion layer grows in mass.

Equation (4) is used to prescribe the boundary luminosities,  $L_{\nu,c} \equiv \ell_{\nu_i} L_{\nu,c}$ , of neutrinos of all kinds with  $\ell_{\nu_e} = 0.20$ ,  $\ell_{\bar{\nu}_e} = 0.15$ ,  $\ell_{\nu_x} = 0.1625$ . (This choice corresponds to a certain loss of lepton number from the PNS core, see Scheck et al. (2006), and ensures a reasonable evolution of  $Y_e$  in the PNS mantle and surface layers, but does not have much relevance for the dynamical evolution.)

In our simulations  $m_{\text{acc}}$  is taken to be the mass between the inner grid boundary and a density of  $\rho_0 = 10^{10}$  g cm $^{-3}$  at radius  $r_0$ , where we define  $\dot{m}_{\text{acc}} = -4\pi r_0^2 v_0 \rho_0$  (accretion means a velocity  $v_0 < 0$  and  $\dot{m}_{\text{acc}} > 0$ ). The core radius is assumed to contract according to  $R_c(t) = R_{c,f} + (R_{c,i} - R_{c,f})/(1+t)^n$  with  $R_{c,i} = R_{\text{ib}}^i$  and  $R_{c,f}$  being the initial and final radius, respectively, and  $t$  is measured in seconds. With  $\Gamma = 3$ ,  $n = 3$ , and  $\zeta = 0.6$  a choice of  $R_{c,f} = 6$  km allows us to reproduce  $E_{87A}$  and  $M_{\text{Ni},87A}$  of SN 1987A for progenitors in the  $20 M_\odot$  range. For the simulations discussed below the  $19.8 M_\odot$  progenitor serves for the calibration of the PNS core model, but the overall results are similar when neighboring stars or a SN 1987A blue supergiant progenitor (Woosley et al. 1988) are used for the calibration.

The chosen parameter values lead to typical PNS neutrino-cooling times ( $t_{90}$  for 90% of the total neutrino-energy release) of 3.5–5.5 s (Fig. 1). This is shorter than the  $\sim 10$  s of emission inferred from the SN 1987A neutrino events of Kamiokande II. However, this detector reported a 7 s gap after 8 events in the first two seconds, and the last 3 events were very close to the detection threshold (Hirata et al. 1987). It is interesting to note that the neutrino signal in all three experiments (Kamiokande II, Irvine-Michigan-Brookhaven, and Baksan) is compatible with a PNS cooling period (exponential cooling timescale) of only 4–5 s (Raffelt 1996; Loredo & Lamb 2002; Pagliaroli et al. 2009). We emphasize that our 1D simulations, which ignore postshock convection and yield standard values for the mean energies of radiated  $\nu_e$  and  $\bar{\nu}_e$  ( $\langle \epsilon \rangle \approx 10$ –17 MeV), naturally require overestimated postbounce neutrino fluxes to trigger explosions. Since the neutrino emission of our PNS model is roughly compatible with the SN 1987A data, we expect that our calibration by SN 1987A explosion properties reflects trends of neutrino-driven explosions whose validity for the investigated progenitors in relative comparison holds beyond the considered 1D setup. We also stress that our description of the core behavior accounts for the presence of an accretion layer surrounding the (excised) high-density core. The growth of this layer depends on the progenitor-specific stellar structure and corresponding mass-infall rate. Its accretion luminosity, which adds to the analytically modeled core-neutrino luminosity, is also included in our simulations by means of a simple, grey transport approximation.

### 2.2. Progenitor stars

Density profiles versus radius and mass for a subset of the investigated 101 solar-metallicity progenitors with iron cores (Woosley et al. 2002) are displayed in Fig. 2. The models are given in  $0.2 M_\odot$  steps between  $10.8 M_\odot$  and  $28 M_\odot$  and further up to  $40 M_\odot$  in  $1.0 M_\odot$  steps. We also include two stars with  $10.0 M_\odot$  and  $10.2 M_\odot$ . While the density profiles are labeled with the zero-age-main-sequence (ZAMS) masses, the corresponding pre-collapse masses are visible in Fig. 3. The images show that mass loss is moderate up to  $19.6 M_\odot$  (except for the  $10.0 M_\odot$  case) but grows considerably beyond that value.

The nonmonotonic star-to-star scatter is a consequence of




FIG. 1.— Timescale of 90% of the neutrino-energy loss of the forming compact remnant as function of the progenitor ZAMS mass. Vertical ticks in the upper part of the diagram indicate BH formation cases. These correspond to “cooling times” in excess of 6 s, because the compact object in our simulations remains radiating neutrinos even when its mass nominally exceeds the BH formation limit. Red histogram bars indicate successful explosions, grey ones correspond to cases where BHs form without a SN explosion, and the green bar marks the  $19.8 M_{\odot}$  progenitor used for the calibration with SN 1987A observations (see text).




FIG. 2.— Density profiles vs. radius (*top*) and vs. enclosed mass (*bottom*) at the onset of core collapse for a selection of models from the considered set of solar-metallicity progenitors with iron cores. Solid lines correspond to ZAMS masses less than  $19 M_{\odot}$ , dashed lines to higher values. Note that the stellar shell structure and also the high-density core ( $\rho \gtrsim 10^5 \text{ g cm}^{-3}$ ) exhibit considerable variations with the progenitor mass (see also Fig. 4).

the turn-on and -off of convective shells during the late evolution stages. The scatter can be seen in Fig. 4 for different structural parameters as functions of the ZAMS mass: The iron-core mass,  $M_{\text{Fe}}$ , is identified with the volume where the electron fraction  $Y_e < 0.495$ , the binding energy of the




FIG. 3.— Stellar masses at the onset of core collapse as function of ZAMS mass. Mass loss is particularly strong beyond  $\sim 20 M_{\odot}$ . The red histogram bars indicate successful explosions, the grey shaded ones correspond to cases where BHs form without a SN explosion, and the green bar marks the  $19.8 M_{\odot}$  progenitor used for the calibration with SN 1987A observations (see text).

overlying shells,  $E_b$ , is evaluated at the onset of core collapse, and the enclosed mass at the inner edge of the oxygen-burning shell,  $M_{\text{O}_\zeta}$ , is defined by the point where the dimensionless entropy per nucleon reaches a value of 4. Finally, the compactness parameter  $\xi_{2.5}$  is defined as in Eq. (10) of O’Connor & Ott (2011) by the ratio of mass  $M = 2.5 M_{\odot}$  and radius  $R(M)$  that encloses this mass:

$$\xi_{2.5} \equiv \frac{M/M_{\odot}}{R(M)/1000 \text{ km}}, \quad (5)$$

where we consider  $M$  as the baryonic mass. For our solar-metallicity progenitors it makes no noticeable difference whether we compute  $\xi_{2.5}$  at the onset of collapse or at the time of bounce (as suggested by O’Connor & Ott 2011), in contrast to the situation for low-metallicity stars, which experience less mass loss during their evolution and which develop more compact cores.

The grey histogram bars mark the cases where we found BH formation without a SN explosion. No clear correlation with extrema of any of the characteristic parameters can be observed. (We note that the binding energy at bounce, though different in detail, does not reveal any better correlation.) While many of the BH formation cases are located in a local peak of  $M_{\text{Fe}}$ ,  $E_b$ ,  $M_{\text{O}_\zeta}$ , and  $\xi_{2.5}$  between  $\sim 22 M_{\odot}$  and  $\sim 26 M_{\odot}$ , there are exceptions in this interval. Moreover, there are BH formation cases below and above this ZAMS mass window with clusterings around  $15\text{--}16 M_{\odot}$ ,  $\sim 18 M_{\odot}$ , and  $21\text{--}22 M_{\odot}$ . The clearest indication of possible BH formation seem to be high values of  $\xi_{2.5}$ , at least local maxima (as visible between  $15$  and  $16 M_{\odot}$  and around  $18 M_{\odot}$ ). But there are exceptions like progenitors near  $21 M_{\odot}$ , which have relatively high values of  $\xi_{2.5}$  and  $E_b$ . Despite this fact they explode, correlated with possessing lower values of  $M_{\text{O}_\zeta}$  than BH formation cases in the close neighborhood. We therefore conclude that the dynamics leading to a successful SN blast does not depend on a single property of the progenitor star but on different ones or even a combination of them.

Although of all quantities displayed in Fig. 4 the compactness parameter  $\xi_{2.5}$  in the form of local maxima exhibits the tightest correlation with BH formation cases, the corresponding threshold value considered by O’Connor & Ott (2011),




FIG. 4.— Quantities characterizing the core structure of the progenitors at the onset of collapse vs. ZAMS mass: iron-core mass (*top left*), binding energy of the matter outside of the iron core in units of  $1 \text{ B} = 1 \text{ bethe} = 10^{51} \text{ erg}$  (*top right*) at the onset of collapse, enclosed mass at the bottom of the oxygen-burning shell (*bottom left*), and compactness parameter  $\xi_{2.5}$  of the innermost  $2.5 M_{\odot}$  as defined in Eq. (5) (*bottom right*). Red, grey, and green histogram bars have the same meaning as in Fig. 3.

i.e.,  $\xi_{2.5} > 0.45$ , significantly underestimates the BH formation probability compared to our results. The progenitor variability seen in Fig. 4, however, defies the specification of a better and reliable number for the BH formation limit. Instead, there is a wide range of  $\xi_{2.5}$  values between  $\sim 0.15$  on the one side and nearly 0.35 on the other side, for which NS and BH formation cases occur. Of course, any such conclusions can be affected by the 1D nature of our simulations. It cannot be excluded that multi-dimensional effects will shift the  $\xi_{2.5}$  limit for BH formation to larger values.

### 3. RESULTS FOR EXPLOSIONS AND REMNANTS

The stellar core collapse until  $\sim 10$ – $15$  ms after bounce is computed with the PROMETHEUS-VERTEX code including sophisticated neutrino transport (Rampp & Janka 2002; Buras et al. 2006a). At that time the bounce shock has converted to a slowly expanding accretion shock. We then continue the simulations of the subsequent accretion phase and the possible explosion with the PROMETHEUS version described in Sect. 2. The mapping, excision of the NS core, and approximate neutrino treatment does not cause any worrisome transients.

Explosions can develop in the case of a favorable interplay of mass accretion rate and neutrino luminosities (e.g., Burrows & Goshy 1993; Janka 2001; Fernández 2012). In all successful cases compared to failed explosions of neighboring progenitors, the mass accretion rate is either lower during a long postbounce period or decreases rapidly when a composition-shell interface arrives at the shock. Shock revival occurs when the neutrino luminosity is still sufficiently high (and thus neutrino heating strong enough) at this

time. In a large number of successful and unsuccessful models the decreasing mass-accretion rate triggers shock oscillations, which indicate the proximity to runaway conditions (Buras et al. 2006b; Murphy & Burrows 2008; Fernández 2012) and whose amplification also leads to large-amplitude pulses of the accretion component of the driving neutrino luminosity (see Buras et al. 2006b). In some stars the explosion is fostered by the Si/O interface reaching the shock relatively soon after bounce, either due to its location at a smaller mass coordinate or because of higher mass accretion rates at earlier times, corresponding to a more compact Si-layer. In this case the high accretion luminosity seems to be supportive. (More information on the time evolution, dynamics, and the neutrino emission of our models will be provided in a separate paper). In summary, the destiny of a collapsing star does not hinge on a single parameter but depends on the overall structure of the stellar core.

Figure 5 gives an overview of the results of our whole model set. All displayed quantities exhibit considerable scatter even in narrow mass windows, which is a consequence of the nonmonotonocities of the progenitor structure. Failed explosions with BH formation seem to be possible for progenitors below  $20 M_{\odot}$ , and successful SNe with NS formation are found also between  $20$  and  $40 M_{\odot}$ . While below  $15 M_{\odot}$  all core collapses produce NSs, the investigated progenitor set yields several “islands” with preferred BH creation above  $15 M_{\odot}$ .

The energies of the neutrino-driven explosions do not exceed  $2 \times 10^{51} \text{ erg}$  and  $^{56}\text{Ni}$  production up to  $0.1$ – $0.15 M_{\odot}$  can be expected. Note that our determination of nickel yields




Fig. 5.— Explosion and remnant properties resulting from our parametrized 1D neutrino-driven SN simulations: explosion energy (*top left*), time of the onset of the explosion (*top right*), baryonic mass of the compact remnant (*middle left*), total release of gravitational binding energy by the compact remnant in neutrinos (*middle right*), and ejected  $^{56}\text{Ni}$  mass (*bottom left*) as functions of stellar birth (ZAMS) mass. The lower right panel shows the compact remnant mass vs. the enclosed mass at the bottom of the oxygen-enriched silicon shell of the progenitor where the stars possess an entropy jump of varying size. The green histogram bar indicates the  $19.8 M_{\odot}$  calibration model (see text). While vertical ticks in some panels mark masses where computed models did not explode, grey histogram bars reaching to the upper panel edge and arrows in the bottom right panel signal the formation of a BH containing the whole mass of the progenitor at collapse. The only exception here is the  $37 M_{\odot}$  star, where the explosion expulses  $\sim 3.2 M_{\odot}$  while  $4.5 M_{\odot}$  of fallback give birth to a BH with  $6.5 M_{\odot}$ . Blue histogram segments indicate fallback masses and orange segments the uncertainties of the  $^{56}\text{Ni}$  ejecta masses. The latter uncertainties are associated with inaccuracies in the  $Y_e$  determination of the neutrino-heated ejecta because of our approximative treatment of neutrino transport.

is uncertain because the neutrino transport approximation does not provide very accurate information of the  $Y_e$  in the neutrino-heated ejecta and in the neutrino-driven wind, where a sizable contribution to the ejected nickel may come from (orange histogram bars in the bottom left panel of Fig. 5 denote the corresponding production masses of the neutron-rich tracer nucleus in the network). The onset of the SN blast (roughly defined by the moment the shock passes 500 km) varies in time between  $\sim 0.1$ – $1.1$  s after bounce, thus including “early” as well as “late” explosions. Later explosions tend to be less energetic, because less mass is available in the more dilute gain layer for absorbing energy from neutrinos. This simple relation for each progenitor is partly masked by the large differences of the mass infall rates in the different stars. Moreover, the neutrino-driven wind, which consists of mass shed off the PNS surface by neutrino heating, adds a contribution to the explosion power that does not depend on the density structure of the silicon and oxygen layers.

The NS baryonic masses are in the range of  $\sim 1.2$ – $2 M_{\odot}$ . For the observable gravitational masses these numbers need to be reduced by the mass defect associated with the total neutrino-energy loss during NS formation, corresponding to  $\Delta E_{\text{tot}}/c^2$ . Our model estimates of  $\Delta E_{\text{tot}}$  are given in the middle right panel of Fig. 5, implying corrections roughly between  $0.11 M_{\odot}$  and  $0.23 M_{\odot}$  for the least massive up to the most massive NSs. However, these numbers are approximative —though in a reasonable range—, and exact binding energies depend on the NS EoS and have to be determined from detailed general relativistic NS structure models. The smallest BH contains  $6.5 M_{\odot}$  and forms in the  $37 M_{\odot}$  star by late fallback of  $4.5 M_{\odot}$ . Since the progenitor has a mass of  $\sim 9.7 M_{\odot}$  at collapse, the SN explosion ejects roughly  $3.2 M_{\odot}$ . All other BHs originate from failed explosions and absorb the total mass of the progenitor at collapse ( $> 8.5 M_{\odot}$ ). Note that we do not terminate the neutrino emission from the excised core when the mass of the compact remnant exceeds the BH




FIG. 6.— Normalized distribution of (baryonic) remnant masses originating from the neutrino-driven explosion models of our investigated set of solar-metallicity progenitors, weighted by Salpeter’s (1955) stellar birth rate function. While the main panel shows the fraction of remnants per  $0.25 M_{\odot}$  bin, the insert provides an enlarged and refined display of the NS mass range with histogram bars giving the remnant fractions per  $0.10 M_{\odot}$  bin.

formation limit. Instead, the accreting central object is considered to remain gravitationally stable, for which reason its total energy loss can reach  $\Delta E_{\text{tot}} > 5 \times 10^{53}$  erg (middle right panel of Fig. 5) and its cooling time can become  $t_{90} \geq 6$  s (Fig. 1). However, none of our successful explosions was triggered by this unphysical, long-time release of neutrino energy, because all NSs, including the one temporarily present in the  $37 M_{\odot}$  progenitor, have a baryonic mass of  $\leq 2 M_{\odot}$ . This is safely below the gravitational mass limit of  $1.97 \pm 0.04 M_{\odot}$  set by the recent precision measurement of Demorest et al. (2010), according to which objects up to at least this mass do not collapse to BHs.

A gap that seems to be present in the observed remnant distribution (e.g., Özel et al. 2010; Valentim et al. 2011; Farr et al. 2011) is compatible with our results. This is visible in Fig. 6, which displays the normalized (baryonic) remnant mass distribution originating from our neutrino-driven explosion models, weighted by Salpeter’s (1955) stellar birth rate function (with power-law index  $-2.35$ ). We obtain a relatively wide distribution of NS masses (see insert). Note that these are birth masses resulting from single, nonrotating (or at most slowly rotating) stars, for which effects like mass transfer and mass loss associated with possible binary interaction during the progenitor evolution have been ignored. Accretion of the NSs in binaries will, of course, also modify this NS birth-mass distribution.

The fallback mass that adds to the initial NS mass is shown in blue in the middle left panel of Fig. 5. Fallback is bigger (up to  $\sim 0.2 M_{\odot}$ ) for the lower-mass progenitors where an extended hydrogen envelope leads to a significant deceleration of the explosion shock and a stronger reverse shock. We point out that in the considered sample of 101 solar-metallicity progenitors only the  $37 M_{\odot}$  star produces a fallback SN with delayed BH formation. In this case the black hole swallows a baryonic mass of  $\sim 6.5 M_{\odot}$  while the explosion ejects the remaining  $\sim 3.2 M_{\odot}$  of the pre-collapse mass of the star. The distribution of Fig. 6 implies that 22.5% of the solar-metallicity progenitors produce BHs at the end of their lives.

Although the remnant mass follows a trend of growth with the enclosed mass at the bottom of the oxygen-burning shell, this location is no reliable indicator for the fate of the star

because some models with relatively small silicon core do not explode. However, BH formation clusters in regions where local maxima of the NS mass occur.

#### 4. CONCLUSIONS

We performed simulations of neutrino-powered explosions in spherical symmetry for a large set of solar-metallicity, nonrotating stars with Fe-cores. We calibrated an analytic neutrino-cooling model of the PNS core such that the observed explosion properties of SN 1987A were reproduced for progenitors around  $20 M_{\odot}$ . This yields numbers for the total energy release and cooling timescale of the newly formed NS in rough agreement with the SN 1987A neutrino measurement, thus confirming the overall consistency of our approach.

The neutrino-driven explosions turned out to be fostered by “steps” in the stellar density and entropy profiles, which reduce the mass-infall rate and ram pressure of the accretion flow and thus allow the shock to expand. Explosions set in either by a continuous acceleration of the shock or after a phase of shock oscillations with growing amplitude. The onset times range from only 0.1 s to more than 1 s after core bounce, meaning that depending on the progenitor the explosion can occur relatively early or with a considerable delay. This conflicts with the claim that neutrino-driven explosions should generally be launched soon after bounce in order to account for the observed gap between highest NS masses and smallest BH masses (Belczynski et al. 2011).

Reflecting the “erratic” structure differences of the progenitor models, the explosion properties and remnant masses exhibit highly nonmonotonic variability even in narrow ranges of the ZAMS mass. This might explain the considerable differences of the explosion energies and nickel masses inferred for observed SNe from lightcurve and spectral analyses as function of the ZAMS mass of the progenitors (e.g., Utrobin & Chugai 2011; Smartt et al. 2009; Smartt 2009) without invoking additional pre-collapse degrees of freedom like rapid core rotation. Neutrino heating seems hardly able to provide explosion energies significantly higher than  $2 \times 10^{51}$  erg and  $^{56}\text{Ni}$  yields in excess of  $0.1\text{--}0.2 M_{\odot}$ . Observed SNe with higher explosion energies and nickel production (e.g., Tanaka et al. 1999) are therefore likely to be driven by a mechanism that is different from neutrino heating and could involve magnetohydrodynamic processes.

The distribution of NS birth masses (including fallback) spans from  $\sim 1.2 M_{\odot}$  to  $\sim 2.0 M_{\odot}$  in baryonic mass with a broad maximum between  $1.4 M_{\odot}$  and  $1.7 M_{\odot}$ . BH formation occurs in certain mass intervals and seems to be possible not only for progenitors with ZAMS masses beyond  $20 M_{\odot}$  but also between  $15 M_{\odot}$  and  $20 M_{\odot}$ . With a single exception (the  $37 M_{\odot}$  progenitor) all BHs swallow the progenitor completely. This suggests that fallback SNe could be rare cases in solar-metallicity environments although roughly 23% of the progenitor population end as BHs. Because of the large mass loss of stars with  $M_{\text{ZAMS}} > 15 M_{\odot}$ , the fallback is strongest (up to  $\sim 0.2 M_{\odot}$ ) for stars below  $\sim 20 M_{\odot}$ . Correspondingly, we find a wide gap between the maximum NS birth mass (around  $1.8 M_{\odot}$  gravitational mass) and the minimum BH mass (roughly  $6 M_{\odot}$ ), compatible with conclusions from observed compact remnant masses (e.g., Özel et al. 2010; Valentim et al. 2011; Farr et al. 2011). It should, however, be noted that our predictions do not account for binary effects prior and after the SN explosion, nor do we include NSs formed from ONeMg-core progenitors. A direct

comparison with observations therefore requires caution.

The compactness parameter  $\xi_{2.5}$  introduced by O’Connor & Ott (2011) provides the best indicative quantity based on the pre-collapse structure for the final fate of a star, i.e., whether it is likely to end as NS or BH. However, our models fail to explode significantly below the discriminating value of  $\xi_{2.5} = 0.45$  considered by O’Connor & Ott (2011). If this result were valid more generally, in particular also for lower-metallicity progenitors, it will have implications for the discussion of GRB progenitors (O’Connor & Ott 2011; Dessart et al. 2012). Instead of depending on an exact bifurcation value of  $\xi_{2.5}$  we see BH formation occurring preferentially (but not exclusively) in regions with local maxima of  $\xi_{2.5}$  and NS as well as BH formation to be possible for  $\xi_{2.5}$  between  $\sim 0.15$  and  $\sim 0.35$ . The success or failure of the explosion mechanism obviously depends on the progenitor structure (determining the time-dependent mass-accretion rate, shock radius, and accretion luminosity) in a complex way and cannot be predicted exactly on grounds of a single parameter.

So far we have investigated only solar metallicity progenitors from one modeling group, but we plan to extend our studies also on sets of progenitors from other groups and for different metallicities. Naturally, the spherical symmetry of our simulations is a major constraint, since the explosion mechanism has been recognized to be intrinsically multi-dimensional (e.g., Janka et al. 2007; Burrows et al. 2007). Nonradial fluid flows in the gain layer have a supportive influence and thus reduce the critical luminosity for driving the explosion (e.g., Janka & Müller 1996; Murphy et al. 2009; Nordhaus et al. 2010; Hanke et al. 2011). With a higher neutrino-heating efficiency lower values of the early postbounce luminosities from our analytic PNS core-cooling model would be needed to reproduce the explosion properties of SN 1987A. This, in turn, would stretch the PNS cooling timescale, which in our simulations tends to be on the lower side of the range compatible with the SN 1987A neutrino data.

Nevertheless, this would only mean a recalibration of the time decay of the core-neutrino source adopted for our modeling, but it is unlikely to fundamentally alter qualitative aspects of the behavior of the different progenitors in relative comparison to each other. Therefore we are hopeful that our basic conclusions will retain their validity also in the multi-dimensional case even if quantitative changes (for example lower explosion energies by less powerful neutrino-driven winds in low-mass progenitors) could be the consequence. It also cannot be excluded that some of the extreme sensitivity of 1D explosions to details of the progenitor structure might be moderated or disappear in multi-dimensional models. It is interesting to note, however, that the relatively early explosions of the  $11.2 M_{\odot}$  and  $27 M_{\odot}$  progenitors of our model set and the

later explosion of progenitors around  $15 M_{\odot}$  are fully compatible with recent results of 2D models with sophisticated neutrino transport (cf. Marek & Janka 2009; Müller et al. 2012, Müller et al., in preparation).

Our findings challenge canonical thinking, e.g. of the progenitor masses leading to BH formation (e.g., Zhang et al. 2008; O’Connor & Ott 2011), and they raise doubts about claims in the literature that the possible gap between NS and BH masses requires explosions that set in relatively early after core bounce (Belczynski et al. 2011; Fryer et al. 2012). We have shown that a highly variable onset time of the explosion and little fallback for massive progenitors (i.e., a low occurrence rate of fallback SNe) naturally leads to the mass gap. On the other hand, a long delay of the beginning of the explosion is questioned by the recent work of Pejcha et al. (2012). While we cannot comment on the statistical approach presented in this paper, which involves a lot of assumptions and is based on a constrained set of data, we do not see how the SN explosion mechanism could achieve to single out a specific value of the gas entropy for success. The core profiles are highly variable for the different progenitors, they are modified by the nuclear burning during the collapse phase, and the neutrino-driven mechanism in spherical symmetry as well as in the multi-dimensional case is sensitive to the mass-infall rate rather than the pre-collapse entropy structure. To be more explicit, the mass-accretion rate and the accretion luminosity exhibit large differences between different progenitors for the specific entropy  $S/(N_A k_B) \approx 2.8$  close to the edge of the iron core, and it is hard to understand how the explosion mechanism could dynamically couple to this specific pre-collapse value of the entropy.

Although our simulations can mean only a very first step of systematically exploring the consequences of neutrino-powered explosions for a wide variety of progenitors, our results already shed new light on a number of important astrophysical questions connected to how the properties of SN explosions and their remnants connect to those of the progenitor stars.

We thank Lorenz Hüpdepohl and Annap Wongwatharat for support. At Garching, this work was supported by the Deutsche Forschungsgemeinschaft through Sonderforschungsbereich/Transregio 27 “Neutrinos and Beyond”, Sonderforschungsbereich/Transregio 7 “Gravitational-Wave Astronomy”, and the Cluster of Excellence EXC 153 “Origin and Structure of the Universe”. A.A. acknowledges funding by the Helmholtz-University Young Investigator Grant No. VH-NG-825. The computations were partially performed at the Rechenzentrum Garching.

## REFERENCES

- Arcones, A., Janka, H.-T., & L.Scheck. 2007, *A&A*, 467, 1227  
 Aufderheide, M. B., Baron, E., & Thielemann, F. K. 1991, *ApJ*, 370, 630  
 Belczynski, K., Wiktorowicz, G., Fryer, C. L., Holz, D. E., & Kalogera, V. 2011, arXiv:1110.1635  
 Buras, R., Rampp, M., Janka, H.-T., & Kifonidis, K. 2006a, *A&A*, 447, 1049  
 —. 2006b, *A&A*, 457, 281  
 Burrows, A., Dessart, L., Ott, C. D., & Livne, E. 2007, *PhR*, 442, 23  
 Burrows, A., & Goshy, J. 1993, *ApJL*, 416, L75  
 Demorest, P. B., Pennucci, T., Ransom, S. M., Roberts, M. S. E., & Hessels, J. W. T. 2010, *Nature*, 467, 1081  
 Dessart, L., O’Connor, E., & Ott, C. D. 2012, *ApJ*, submitted (arXiv:1203.1926)  
 Farr, W. M., Sravan, N., Cantrell, A., et al. 2011, *ApJ*, 741, 103  
 Fernández, R. 2012, *ApJ*, 749, 142  
 Fryer, C. L. 1999, *ApJ*, 522, 413  
 —. 2006, *NewAR*, 50, 492  
 Fryer, C. L., Belczynski, K., Wiktorowicz, G., et al. 2012, *ApJ*, 749, 91  
 Fryer, C. L., & Kalogera, V. 2001, *ApJ*, 554, 548  
 Fryxell, B. A., Müller, E., & Arnett, W. D. 1989, MPA-Preprint 449 (Max-Planck Institut für Astrophysik, Garching)  
 Hanke, F., Marek, A., Müller, B., & Janka, H.-T. 2011, *ApJ*, submitted (arXiv:1108.4355)  
 Hirata, K., Kajita, T., Koshiba, M., Nakahata, M., & Oyama, Y. 1987, *PhRvL*, 58, 1490

- Janka, H.-T. 2001, *A&A*, 368, 527
- Janka, H.-T., Langanke, K., Marek, A., Martínez-Pinedo, G., & Müller, B. 2007, *PhR*, 442, 38
- Janka, H.-T., & Müller, E. 1996, *A&A*, 306, 167
- Kifonidis, K., Plewa, T., Janka, H.-T., & Müller, E. 2003, *A&A*, 408, 621
- Loredo, T. J., & Lamb, D. Q. 2002, *PhRvD*, 65, 063002
- Marek, A., & Janka, H.-T. 2009, *ApJ*, 694, 664
- Müller, B., Janka, H.-T., & Marek, A. 2012, *ApJ*, submitted (arXiv:1202.0815)
- Murphy, J., Ott, C. D., & Burrows, A. 2009, *ApJ*, 707, 1173
- Murphy, J. W., & Burrows, A. 2008, *ApJ*, 688, 1159
- Nordhaus, J., Burrows, A., Almgren, A., & Bell, J. 2010, *ApJ*, 720, 694
- O'Connor, E., & Ott, C. D. 2011, *ApJ*, 730, 70
- Özel, F., Psaltis, D., Narayan, R., & McClintock, J. E. 2010, *ApJ*, 725, 1918
- Pagliaroli, G., Vissani, F., Costantini, M. L., & Ianni, A. 2009, *Aph*, 31, 163
- Pejcha, O., Thompson, T. A., & Kochanek, C. S. 2012, arXiv:1204.5478
- Plewa, T., & Müller, E. 1999, *A&A*, 342, 179
- Raffelt, G. G. 1996, *Stars as Laboratories for Fundamental Physics. The Astrophysics of Neutrinos, Axions, and Other Weakly Interacting Particles.* (University of Chicago Press)
- Rampp, M., & Janka, H.-T. 2002, *A&A*, 396, 361
- Salpeter, E. E. 1955, *ApJ*, 121, 161
- Scheck, L., Kifonidis, K., Janka, H.-T., & Müller, E. 2006, *A&A*, 457, 963
- Smartt, S., Eldridge, J. J., Crockett, R. M., & Maund, J. R. 2009, *MNRAS*, 395, 1409
- Smartt, S. J. 2009, *ARA&A*, 47, 63
- Tanaka, M., Tominaga, N., Nomoto, K., et al. 1999, *ApJ*, 692, 1131
- Timmes, F. X., & Swesty, F. D. 2000, *ApJS*, 126, 501
- Utrobin, V. P., & Chugai, N. N. 2011, *A&A*, 532, A100
- Valentim, R., Rangel, E., & Horvath, J. E. 2011, *MNRAS*, 414, 1427
- Woosley, S. E., Heger, A., & Weaver, T. A. 2002, *RevMPh*, 74, 1015
- Woosley, S. E., Pinto, P. A., & Ensmann, L. 1988, *ApJ*, 324, 466
- Woosley, S. E., & Weaver, T. A. 1995, *ApJS*, 101, 181
- Zhang, W., Woosley, S. E., & Heger, A. 2008, *ApJ*, 679, 639