Kinematics of Simulated Galaxies I: Connecting Dynamical and Morphological Properties of Early-Type Galaxies at Different Redshifts

Felix Schulze^{1,2*}, Rhea-Silvia Remus¹, Klaus Dolag^{1,3}, Andreas Burkert^{1,2}, Eric Emsellem^{4,5}, and Glenn van de Ven^{5,6}

¹ Universitäts-Sternwarte München, Scheinerstr. 1, D-81679 München, Germany

² Max Planck Institute for Extraterrestrial Physics, Giessenbachstrae 1, D-85748 Garching, Germany

³ Max Planck Institut for Astrophysics, D-85748 Garching, Germany

⁴ Université Lyon 1, Observatoire de Lyon, Centre de Recherche Astrophysique de Lyon and Ecole Normale Supérieure de Lyon,

9 avenue Charles André, F-69230 Saint-Genis Laval, France

⁵ European Southern Observatory, Karl-Schwarzschild-Str. 2, 85748 Garching, Germany

⁶ Max Planck Institute for Astronomy, Königstuhl 17, 69117 Heidelberg, Germany

Accepted ... Received ...; in original form ...

ABSTRACT

State-of-the-art integral field surveys like ATLAS^{3D}, SLUGGS, CALIFA, SAMI, and MaNGA provide large data sets of kinematical observations of early-type galaxies (ETGs), yielding constraints on the formation of ETGs. Using the cosmological hydrodynamical Magneticum Pathfinder simulations, we investigate the paradigm of fast and slow rotating ETGs in a fully cosmological context. We show that the ETGs within the *Magneticum* simulation are in remarkable agreement with the observations, revealing fast and slow rotators quantified by the angular momentum proxy $\lambda_{\rm R}$ and the flattening ϵ with the observed prevalence. Taking full advantage of the three-dimensional data, we demonstrate that the dichotomy between fast and slow rotating galaxies gets enhanced, showing an upper and lower population separated by an underpopulated region in the edge-on $\lambda_{\rm R}$ - ϵ plane. Following the evolution of the $\lambda_{\rm R}$ - ϵ plane through cosmic time, we find that, while the upper population is already in place at z = 2, the lower population gets statistically significant below z = 1 with a gradual increase. At least 50% of the galaxies transition from fast to slow rotators on a short timescale, in most cases associated to a significant merger event. Furthermore, we connect the M_{*} - j_{*} plane, quantified by the b-value, with the $\lambda_{\rm R}$ - ϵ plane, revealing a strong correlation between the position of a galaxy in the $\lambda_{\rm R}$ - ϵ plane and the b-value. Going one step further, we classify our sample based on features in their velocity map, finding all five common kinematic groups, also including the recently observed group of prolate rotators, populating distinct regions in the $\lambda_{\rm R}$ -b plane.

Key words: galaxies: evolution – galaxies: formation – galaxies: kinematics and dynamics – cosmology: dark matter – methods: numerical

1 INTRODUCTION

During the last decade results from cosmological simulations within the Λ CDM framework revealed a two-phase picture of galaxy formation (Oser et al. 2010). Subsequent to the hierarchical merging of dark matter halos, a rapid dissipative phase of smooth gas accretion triggering in-situ star formation proceeds the galaxy assembly. Below $z \approx 1$ the main driver of galaxy growth is minor and major merging

of larger structures from outside the halo, hence most new stars are accreted onto the galaxy. Due to the complex interplay of baryonic processes formulating a predictive theory of galaxy formation is still highly debated.

Starting with the famous Hubble-sequence proposed in 1926 (Hubble 1926) many classification schemes attempted to capture the diversity of galactic structures. Especially within the category of early-type galaxies (ETGs), the improvement of observational techniques revealed a more complex picture of the internal structure. The discovery, that the isophotal shape of ETGs differs from ellipses and can be ei-

^{*} E-mail:fschulze@usm.lmu.de

ther 'discy' or 'boxy' as measured by the a_4 parameter led to a revision of the picture of ETGs (Bender et al. 1988). Furthermore, the result that discy ETGs seemed to rotate more rapidly than boxy ETGs and the correlation between the isophotal shape and the central surface density slope led to the proposal of a new classification scheme for ETGs capturing those properties (Kormendy & Bender 1996; Faber et al. 1997).

Since the discovery and classification of ETGs within the Hubble-sequence the general formation theory has converged to a picture in which, due to the hierarchical assembly of dark matter halos, disc galaxies get morphologically transformed into ETGs in major or multiple minor mergers of disk galaxies (Toomre & Toomre 1972; Toomre 1977; Barnes 1988, 1992; Hernquist 1992; Mihos & Hernquist 1996; Naab & Burkert 2003; Bournaud et al. 2005, 2007). Processes like dynamical friction and violent relaxation during galaxy mergers are capable of drastically modifying the statistical distribution of orbits, while triggered star formation and accretion of fresh gas builds up new cold components (Lynden-Bell 1967; Thomas et al. 2009; Hoffman et al. 2010). Since the formation of ETGs is predominantly driven by mergers they preserve a richness of information about their assembly encoded in the orbital structure and therefore also in the velocity field.

The emerging picture of ETGs, acquired from photometry, was advanced by the development of integral-field spectroscopy allowing to measure detailed spatial maps of several galaxy properties including stellar kinematics. The SAURON survey was the first project to map the twodimensional kinematics, ionised gas and stellar population of a statistically significant sample of 48 nearby ETGs (de Zeeuw et al. 2002; Bacon et al. 2001). Already a simple visual investigation of stellar line-of-sight velocity maps within one effective radius revealed two types of kinematics: The class of fast rotators shows a regular rotating velocity pattern consistent with an inclined rotating disc, whereas the class of slow rotators features complex kinematical patterns like kinematically distinct and counter-rotating cores (Sarzi et al. 2006; Emsellem et al. 2004; Davies et al. 2001). These results provoked a paradigm shift towards a kinematically motivated classification scheme closer related to the formation history, since stellar kinematics are believed to encode the detailed accretion history of a galaxy (Cox et al. 2006).

By extending the SAURON sample to 260 ETGs in a volume- and luminosity-limited survey the ATLAS^{3D} project substantiated the dichotomy of fast and slow rotators based on a statistically more meaningful sample (Emsellem et al. 2011). Comprising 86% of the total sample, fast rotators are significantly more frequent and seem to be the dominant kinematical final stage of ETGs. Subsequent to these pioneering studies, Cappellari (2016) combined results from ATLAS^{3D} and the SAMI Pilot survey (Fogarty et al. 2015) to strengthen the existence of the two classes of fast and slow rotators.

Aiming to gain insight into the general formation pathway of these two types of ETGs, studies were conducted in an attempt to link the cosmological environment of ETGs to their kinematical state, i.e. into a 'kinematic morphologydensity relation'. Interestingly, the global fraction of slow rotators within relaxed clusters has proven to be remarkably constant across a large range of clusters with varying galaxy number density (D'Eugenio et al. 2013; Houghton et al. 2013; Scott et al. 2014; Fogarty et al. 2015; Veale et al. 2017). Only when investigating the local density dependence within the cluster these studies found a rising slow rotator fraction in higher density environments. Interestingly, Veale et al. (2017) found that for a given absolute K-band magnitude, which is strongly correlated with the stellar mass of a galaxy, the slow rotator fraction is independent of the local and global environment (see also Greene et al. (2018)).

Attempting to capture the variety of kinematical features, Krajnović et al. (2011) prior introduced a more refined classification scheme: The velocity map of a galaxy is found to (a) not show any sign of rotation, (b) exhibit a complex velocity pattern without any specific feature, (c) feature a kinematically distinct core (KDC, including counterrotating cores CRCs), (d) show a double peak in the dispersion map or (e) show a regular rotation pattern. Classifying those groups into fast and slow rotators revealed that group (a)-(d) are in general slow rotating, while members of group (e) are fast rotating (Emsellem et al. 2011). Observational results from McDermid et al. (2006) revealed that young and very compact KDCs can also be found in fast rotating ETGs. Furthermore, Schulze et al. (2017) showed that KDCs can dissolve on a timescale of 1.5Gyr and feature a complex global motion with respect to the surrounding galaxy.

To depict the different formation pathways responsible for the dichotomy of fast and slow rotators in ETGs, several hydrodynamical simulations using isolated galaxy mergers have been performed. Since this kind of simulations allows to follow the evolution of galaxy properties through time they represent a diagnostically conclusive method to probe the formation of fast and slow rotators, and even more detailed kinematical features, depending on the respective merger configuration. These studies confirmed the notion that galaxy mergers are capable of transforming kinematically cold disks into spheroidal hot objects, resembling the kinematics of fast and slow rotators (Jesseit et al. 2009; Bois et al. 2011; Moody et al. 2014; Bois et al. 2010). The kinematical properties of the remnants crucially depends on the orbital parameters of the merger, the detailed merger sequence (binary or multiple mergers) and the intrinsic properties of the progenitors like mass-ratio and gas content.

Although idealised disk merger simulations had major success in resembling kinematic and photometric properties of galaxies, this method is limited to artificially determined initial conditions and hence does not represent a natural formation pathway in a fully cosmological environment. Due to a significant improvement of computational and numerical techniques, recent state-of-the-art hydrodynamical cosmological simulations reach resolutions that allow to investigate the kinematic properties of ETGs in a statistical meaningful manner. Recent studies using cosmological zoom-in simulations showed that fast and slow rotators can be formed in various scenarios through minor and major mergers (Naab et al. 2014; Choi & Yi 2017). However, investigating the influence of different physical processes on the spin evolution, Choi & Yi (2017) found that, in contrast to the general expectation, the combined impact of effects other than mergers like extremely minor mergers (of ratios smaller than 1/50), secular evolution, fly-by encounters, harassment and dynamical friction is dominating the spin evolution. Therefore, the simple picture of relating the kinematic evolution of ETGs

solely to mergers seems to be insufficient. In a first attempt to comprehend the formation of fast and slow rotators in a fully cosmological context Penoyre et al. (2017) followed the evolution of ETGs in the Illustris simulation. They find that at z = 1 the respective progenitors of fast and slow rotators are indistinguishable. At lower redshift they find major mergers to be the driving mechanism engendering the kinematical dichotomy. As a key difference in the evolution of fast and slow rotators they conclude that fast rotators, in contrast to slow rotators, get spun up by accretion of fresh gas and stars from the cosmic environment. In line with these results Lagos et al. (2018) show that, within the EA-GLE simulation, gas-rich mergers can significantly increase the angular momentum content of galaxies.

The aim of this work is to study the kinematic properties of ETGs in a statistical manner using the cosmological hydrodynamical *Magneticum Pathfinder* simulations. Rather than focusing on individual formation pathways we follow the formation of fast and slow rotators as populations and relate their intrinsic kinematics to global and morphological properties. In a second step we apply a refined classification based on kinematical features in the velocity maps similar to Krajnović et al. (2011) and (Emsellem et al. 2011), including the recently observed prolate rotators (Tsatsi et al. 2017; Ebrová & Lokas 2017), to further disentangle the kinematical variety of ETGs.

2 SIMULATION AND ANALYSIS

2.1 The Magneticum Pathfinder Simulation

The galaxies investigated in this study are extracted from the *Magneticum Pathfinder* simulations¹, which are a set of cosmological hydrodynamical simulations performed with the Tree/SPH code GADGET-3. GADGET-3 is an extended version of GADGET-2 (Springel 2005; Springel et al. 2001a) including improvements concerning the treatment of viscosity and the used kernels (Dolag et al. 2005; Donnert et al. 2013; Beck et al. 2015).

Furthermore, the simulations include a wide variety of baryonic physics such as gas cooling and star formation (Springel & Hernquist 2003), black hole seeding, evolution and AGN feedback (Springel et al. 2005; Fabjan et al. 2010; Hirschmann et al. 2014; Steinborn et al. 2015) as well as stellar evolution and metal enrichment (Tornatore et al. 2007).

The Magneticum Pathfinder simulations implement a standard Λ CDM cosmology with parameters adapted from the seven-year results of the Wilkinson Microwave Anisotropy Probe (WMAP7) (Komatsu et al. 2011). The density parameters are $\Omega_{\rm b} = 0.0451$, $\Omega_{\rm M} = 0.272$ and $\Omega_{\Lambda} =$ 0.728 for baryons, matter and dark energy, respectively. The Hubble parameter is h = 0.704 and the normalisation of the fluctuation amplitude at 8Mpc is given by $\sigma_8 = 0.809$. The Magneticum Pathfinder simulation set includes boxes of different sizes and resolutions. Sizes range from 2688Mpc/h box length to 18Mpc/h box length, while resolutions cover a particle mass range of $10^{10} > m_{\rm dm} > 10^7 M_{\odot}/h$ for the dark matter and $10^9 > m_{\rm gas} > 10^6 M_{\odot}/h$ for gas particles. The implemented star formation scheme allows a gas particle to form up to four star particles (Springel & Hernquist 2003).

For this study we choose the medium-sized cosmological Box4, with a side length of 48Mpc/h at the ultra high resolution level. With masses for the dark matter and gas particles of $m_{\rm DM} = 3.6 \cdot 10^7 M_{\odot}/h$ and $m_{\rm gas} = 7.3 \cdot 10^6 M_{\odot}/h$, respectively, with a gravitational softening length of 1.4kpc/h for dark matter and gas particles, and 0.7 kpc/h for star particles. This box is chosen to ensures a high enough resolution for galaxy kinematic studies as well as a large sample size to examine the kinematics of galaxies in a statistically meaningful manner.

The galaxies are identified using SUBFIND (Springel et al. 2001b) which utilises a standard Friends-of-Friends algorithm, and is adapted for the treatment of the baryonic component (Dolag et al. 2009), which allows to identify both satellite and central galaxies inside the main haloes.

2.2 Sample Selection

Since we aim to conduct a statistical comparison to results which are based on various observational samples, the galaxy selection should ensure a meaningful comparability. The observational classification scheme which essentially all authors converged on is purely based on the morphological properties of a galaxy (de Vaucouleurs 1959, 1963; Sandage 1961). Adopting the separation criterion between late-type galaxies (LTGs) and ETGs outlined by Sandage (1961), they select ETGs from their complete sample by visually inspecting multi-colour images. Therefore, the separation between LTGs and ETGs is strongly based on the presence of spiral arms, while other galaxy characteristics, which vary with morphology, are neglected (Sandage 1975). This classification scheme was adopted to create the frequently used RC2 and RC2 galaxy catalogue (de Vaucouleurs et al. 1976, 1991).

Due to the potentially large sample size and the challenging task of generating realistic multi-colour images from simulated data, a visual inspection is not convenient for our study. Therefore, we first select all main- and subhaloes identified by SUBFIND with a total stellar mass $M_* > 2 \cdot 10^{10} M_{\odot}$ to ensure a proper mass-resolution (i.e. number of stellar particles). In comparison to the ATLAS^{3D} observations, which represents the largest comparison sample, this lower mass limit introduces a minor bias since they comprise ETGs down to $M_* \gtrsim 6 \cdot 10^9 M_{\odot}$ in their sample. In total, we find 1191 galaxies satisfying this condition in *Magneticum* Box4.

In order to assure an adequate spatial resolution the selection proceeds by constraining the stellar half-mass radius to be larger than twice the stellar softening length:

$$R_{1/2} > \frac{1.4 \text{ kpc}}{h (z+1)}.$$
 (1)

Since the softening length is defined to be constant in comoving coordinates, the threshold involves a redshift dependence in physical units. The half-mass radius is determined to be the radius of a three dimensional sphere containing half of the total stellar mass of the halo.

Due to the star-forming nature of spiral arms their presence is tightly coupled to the amount of cold gas available in

¹ www.magneticum.org

4 Schulze et al.

Table 1. Characteristics of the	Magneticum	ETGs at	z = 0.
---------------------------------	------------	---------	--------

Total number of ETG's:	784
Total halo masses:	$2.7 \cdot 10^{10} - 1.2 \cdot 10^{14} \ M_{\odot}$
Galaxy stellar masses:	$2 \cdot 10^{10} - 1.6 \cdot 10^{12} M_{\odot}$
Galaxy cold gas fractions:	$1 \cdot 10^{-4} - 0.099$
Galaxy effective radii:	$2.0 - 30.5 \; \mathrm{kpc}$

the galaxy. Therefore, we impose a upper limit on the coldgas fraction f_{gas} rather than visually identifying the absence of spiral arms to select a galaxy to be an ETG:

$$f_{\rm gas} = \frac{M_{\rm coldgas}}{M_*} \leqslant 0.1. \tag{2}$$

Here, $M_{\rm coldgas}$ is the total mass of all gas particles possessing a temperature below $10^5 K$ within $3R_{1/2}$. Serra et al. (2014) investigated the HI fraction for a subsample of the ATLAS^{3D} ETGs, observing $M_{\rm HI}/M_*$ -fractions ranging from 0.04% to 10%. To ensure a proper comparability we checked the maximum HI fractions within our sample, yielding a value of 5%. Hence, Eq. 2 imposes a conservative limit on the HI fractions in comparison to the observations.

Our final sample of ETGs at z = 0 includes 784 galaxies, which we will refer to as *Magneticum* ETGs in the following. Some basic characteristics of our sample are summarised in Tab. 1.

To select our sample at higher redshifts we maintain the lower mass cut and the gas fraction threshold, whereas the lower limit for $R_{1/2}$ is adapted according to Eq. 1. This leaves us with a sample size of 157, 670 and 591 at z = 2, z = 1 and z = 0.5, respectively. Therefore already at z = 2we find a significant number of ETGs with a low fraction of cold gas and stellar masses above $2 \cdot 10^{10} M_{\odot}$.

2.3 Mass-Size Relation

Fig. 1 shows the mass-size relation for all galaxies in the simulation satisfying the resolution criteria, separated into selected (left column, black circles) and rejected (right column, grey circles). In each panel the horizontal dashed lines correspond to the resolution limit (Eq. 1). We include observational results from the GAMA (Lange et al. 2015; Baldry et al. 2012), CALIFA, SLUGGS (Forbes et al. 2017) and ATLAS^{3D} (Cappellari et al. 2013b) surveys as given in the legend.

The green and blue solid lines depict the most recent GAMA results by Lange et al. (2015) representing the statistically most significant observational sample in our comparison. From the various relations presented in Lange et al. (2015) we adopt the relations obtained by fitting a double power-law function to the visually identified ETGs and LTGs. It is important to note that Lange et al. (2015) uses actual major-axis effective radii taking the elliptical shape of the isophotes into account (Kelvin et al. 2012). Hence, especially for highly elongated galaxies, this can lead to a bias towards larger radii with respect to the half-mass radii determined for the *Magneticum* sample.

An interesting general feature among the observational samples is the considerable scatter between them. For a given mass, the SLUGGS, CALIFA and ATLAS^{3D} ETGs seem to be systematically more compact than the GAMA galaxies. Since the half-light radii of those samples are obtained by different methods, this suggests a significant impact of the observational approach on the result (Cappellari et al. 2013a).

At the stellar mass range of $2 \cdot 10^{10} M_{\odot} < M_* < 10^{11} M_{\odot}$ the observations show that LTGs have larger effective radii at a given mass than the ETGs. This difference is also present between the *Magneticum* ETGs and rejected galaxies, albeit with a significant overlap, similar to the results from CALIFA and GAMA. However, the rejected galaxies in the right panel of Fig. 1 include disc-like galaxies as well as peculiar galaxies, potentially influenced by interactions. Hence, a rigorous comparison to observations is not reasonable due to the contamination by disturbed objects.

The Magneticum ETGs are in substantial agreement with GAMA observations by Baldry et al. (2012) showing a clustering in the vicinity of the red curve with a weak trend towards larger radii (see right panel of Fig. 1). The more recent results by Lange et al. (2015) exhibit slightly smaller radii in the considered mass range up to $10^{11} M_{\odot}$, however still larger than those found for the ATLAS^{3D} galaxies. For galaxies more massive than $10^{11} M_{\odot}$, the Magneticum ETGs follow the almost linear trend found for the GAMA ETGs, albeit tending towards slightly smaller radii. It is important to note that, due to the limited size of the simulated box, the high mass end of the Magneticum ETGs is underpopulated and therefore statistically unreliable.

Overall, the mass-size relation found for the *Magneticum* ETGs is consistent with the observed trend. Furthermore, the galaxies considered in this study are in agreement with present day scaling relations since they represent an extension of the sample investigated in Remus & Dolag (2016) and Remus et al. (2017).

2.4 Angular Momentum Proxy: λ_R

For many years the V/σ -parameter has been used to quantify the relative amount of stellar rotation in a system, where V and σ denote the projected stellar velocity and velocity dispersion, respectively. This parameter is a useful tool to investigate the dynamical state of ETGs. However, it fails to distinguish between small scale rotation (like kinematically distinct cores) and large scale rotation (Emsellem et al. 2007). Therefore, a new parameter was introduced by Emsellem et al. (2007) within the SAURON project, which takes the global velocity structure into account and in addition maintains the information about the dynamical state, i.e. ordered vs. random motion.

For observed two-dimensional velocity and dispersion maps it is defined as

$$\lambda_R = \frac{\langle R \ |V| \rangle}{\langle R \ \sqrt{V^2 + \sigma^2} \rangle},\tag{3}$$

where R is the projected radius, V the line-of-sight velocity and σ the projected velocity dispersion. $\langle \cdot \rangle$ denotes the luminosity-weighted average over the full two-dimensional kinematic field. Inserting the velocity weighting it reads

$$\lambda_R = \frac{\sum_{i=1}^{N_p} F_i R_i |\overline{V}_i|}{\sum_{i=1}^{N_p} F_i R_i \sqrt{\overline{V}_i^2 + \sigma_i^2}},\tag{4}$$

with the sum running over all pixels in the considered field



Figure 1. The mass-size relation at z = 0 for the *Magneticum Pathfinder* simulation in direct comparison to recent observations. Left panel: Black circles show the distribution for *Magneticum* ETGs. Diamonds correspond to observations by the ATLAS^{3D}, SLUGGS and CALIFA survey with colours as given in the legend. The red and blue solid lines represent observations for ETGs from the GAMA survey. Right panel: *Magneticum* galaxies which are selected to not resemble ETGs in grey, while blue diamonds show observations by the CALIFA survey. The blue dashed line marks observations for LTGs from the GAMA survey (Baldry et al. 2012). In both panels the 1σ range for the GAMA observations are marked by the shaded areas.

of view. F_i , R_i , $|\overline{V}_i|$ and σ_i are the flux, projected distance to the galaxy centre, mean stellar velocity and velocity dispersion of the ith photometric bin, respectively.

For a purely rotational supported system, $\lambda_{\rm R}$ tends to unity. The lower limit of $\lambda_{\rm R}$ \rightarrow 0 corresponds to either a purely dispersion-dominated system with no ordered rotation, or a rotating system where the total angular momentum vector is along the line of sight. The usage of $\langle R | V | \rangle$ as a surrogate for the angular momentum ensures the distinction between large-scale or small-scale rotation. Investigating remnants of simulated binary disc mergers, Jesseit et al. (2009) found that $\lambda_{\rm R}$ is a robust indicator of the true intrinsic angular momentum content of these objects. It is important to mention that $\lambda_{\rm R}$ obviously depends on the spatial size, over which the sum in Eq. 4 is taken. Moreover, it is sensitive to the tessellation method used for the velocity and dispersion maps. $\lambda_{\rm R}$ is specifically customised to fit the needs and constraints of the current observational methods. It solely uses projected quantities and fluxes, which can be observed by multi-wavelength surveys.

For simulations the fluxes are replaced by stellar masses, assuming a constant mass-to-light ratio within each galaxy. Following previous theoretical studies (Jesseit et al. 2009; Naab et al. 2014; Bois et al. 2011; Wu et al. 2014), the expression for $\lambda_{\rm R}$ is transformed into

$$\lambda_R = \frac{\sum_{i=1}^{N_p} M_i R_i |\overline{V}_i|}{\sum_{i=1}^{N_p} M_i R_i \sqrt{\overline{V}_i^2 + \sigma_i^2}}.$$
(5)

Eq. 5 is the final formula we will use in this study to calculate $\lambda_{\rm R}$ from the kinematical maps.

In order to calculate λ_R given in Eq. 5, we need to generate two types of kinematical maps. The mean projected

velocity \overline{V}_i within a pixel is given by

$$\overline{V}_i = \frac{\sum_{j=1}^{N_c} V_j}{N_c},\tag{6}$$

where V_j is the particle velocity, and the sum runs over all N_c particles within a cell. Similarly, the projected velocity dispersion is defined as

$$\sigma_i = \sqrt{\frac{\sum_{j=1}^{N_c} V_j^2}{N_c} - \left(\frac{\sum_{j=1}^{N_c} V_j}{N_c}\right)^2}.$$
 (7)

To be comparable to observations we have to use calculation methods which take the properties of numerical simulations into account. For example, the limited mass resolution in SPH leads to low particle numbers when assigning particles onto a grid, which causes statistical errors. Another issue is the limited spatial resolution which makes the maps sensitive to numerical small-scale fluctuations. For a detailed resolution study on this subject see Bois et al. (2010). In order to avoid artificial statistical noise we adopt the following approach for each galaxy: The stellar particles within the considered spatial domain and projection are sampled onto a rectangular grid with a spatial resolution comparable to state-of-the-art IFU surveys. Subsequently, a Centroidal Voronoi Tessellation (CVT) is performed, which combines pixels into cells containing a minimum of 100 particles, while preserving a proper spatial resolution (Cappellari & Copin 2003). To determine $\lambda_{\rm R}$ we then calculate the lineof-sight mean velocity, velocity dispersion and the radius of the centre-of-mass within each CVT cell.

The calculation of $\lambda_{\rm R}$ proceeds by determining the spatial area over which the sum in Eq. 5 is taken. Given the ellipticity of the galaxy the calculation domain is specified to be an ellipse with the corresponding ellipticity enclosing an area of $A_{\rm ellipse} = \pi R_{1/2}^2$. We indicate this by assigning a subscript '1/2' to the parameter name $\lambda_{R_{1/2}}$. This procedure of calculating $\lambda_{R_{1/2}}$ ensures maximal comparability to observational results.

It is crucial to emphasise that the smallest values obtained from this procedure have to be interpreted with caution due to the functional definition of $\lambda_{R_{1/2}}$. As $\lambda_{R_{1/2}}$ is a cumulative parameter of absolute values, the statistical noise of Voronoi cells with nearly zero velocity adds up, creating a lower limit for $\lambda_{R_{1/2}}$ (Naab et al. 2014; Bois et al. 2010). We verify this notion by calculating $\lambda_{R_{1/2}}$ for all galaxies assuming zero velocity, only considering the statistical noise. It reveals a mean value of 0.07, which corresponds to the lowest values found for the simulated *Magneticum* galaxies.

2.5 Morphological Parameters: Ellipticity and Sérsic-Index

During the course of this paper we use of the ellipticity or flattening ϵ of galaxies. Following the definition by Cappellari et al. (2007) and adapting it to simulated data yields

$$\epsilon = 1 - \sqrt{\frac{\langle y^2 \rangle}{\langle x^2 \rangle}} = 1 - \sqrt{\frac{\sum_{j=1}^N M_i \ y^2}{\sum_{j=1}^N M_i \ x^2}},\tag{8}$$

where the coordinates (x, y) are centred on the galaxy nucleus and x is measured along the morphological major axis. We calculate ϵ for a given projection of a simulated galaxy within an iso-density contour containing the same amount of stellar mass as a circle with a radius of $1.5R_{1/2}$ by diagonalising the two dimensional moments-of-inertia-tensor. This procedure yields an estimate of the average global ellipticity mostly independent of central substructures like bars or disks.

Since the Sérsic-index describes the curvature of the Sérsic-profile fitted to the stellar radial surface density distribution, it is a purely morphological parameter. The Sérsicprofile is given by

$$\Sigma(R) = \Sigma_e \, \exp\left(-b_n \left[\left(\frac{R}{R_e}\right)^{\frac{1}{n}} - 1\right]\right) \tag{9}$$

(Sérsic 1963), where R_e is the effective radius, Σ_e is the surface density at the effective radius, and n is the Sérsicindex. The dimensionless parameter b_n is defined according to the definition of the effective radius. There is no analytic solution to the defining equation for b_n . Hence, we use a numeric approximation given in Ciotti & Bertin (1999).

The Sérsic fits are performed on the edge-on projection of the stellar component which is subdivided into elliptical annuli of fixed ellipticity and position angle. Hence we neglect effects of isophotal twists or variations of ϵ with radius. The border of the annuli are determined by demanding a constant particle number of 200, ensuring a proper weighting of all data points. Excluding one effective radius in the centre the moving window approach is extended out to 5 R_e . In this manner we exclude the influence of central subcomponents like cusps or cores on the fit.

3 THE λ_{R} - ϵ PLANE

Based on a sample of 48 ETG's, Emsellem et al. (2007) used λ_{Re} to separate ETG's into fast and slow rotating objects,

depending on whether the galaxy exhibits a $\lambda_{\rm R_e}$ larger or smaller than 0.1.

Having access to a statistically more complete sample ATLAS^{3D} redefined the criterion to disentangle fast and slow rotators based on 260 ETGs, including the galaxies' shapes:

$$\lambda_{\rm R_e} > 0.31 \cdot \sqrt{\epsilon_{\rm R_e}} \to \text{fast rotator}$$

$$\lambda_{\rm R_e} \leqslant 0.31 \cdot \sqrt{\epsilon_{\rm R_e}} \to \text{slow rotator}$$
(10)

It is based on qualitative theoretical considerations and has proven to properly distinguish the two kinematical families (Emsellem et al. 2011). For our simulated sample of galaxies, we will use an analogue, approximating λ_{R_e} by $\lambda_{R_{1/2}}$ and ϵ_{R_e} by $\epsilon_{R_{1/2}}$

3.1 Statistical Comparison to IFU Observations

We want to test if our simulated sample of ETG's reproduces the observed distribution in the $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ plane. The comparison sample combines observations from the ATLAS^{3D}, CALIFA, SAMI Pilot and SLUGGS survey extracted from Emsellem et al. (2011), Querejeta et al. (2015), Fogarty et al. (2015) and Arnold et al. (2014), respectively. Since the observations are limited to one random projection for each galaxy, we choose an arbitrary viewing angle for each simulated galaxy to determine $\lambda_{R_{1/2}}$ and $\epsilon_{1/2}$.

Fig. 2 shows the $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ plane for the simulated sample (red circles) in direct comparison to the IFU observations (blue and grey symbols). The threshold between fast and slow rotators given in Eq. 10 is shown as solid green line. Furthermore, Fig. 2 comprises cumulative distributions for $\lambda_{R_{1/2}}$ on the right and for $\epsilon_{1/2}$ on top of the main panel each split up into the various samples.

The Magneticum simulation reproduces both, fast and slow rotating ETG's with $\lambda_{\mathrm{R}_{1/2}}$ and $\epsilon_{1/2}$ in the range 0.06 < $\lambda_{R_{1/2}} < 0.72 \text{ and } 0.016 < \epsilon_{1/2} < 0.7, respectively. With 75\%$ (587/784) the vast majority is classified as fast rotators, accordingly only 25% (197/784) are slowly rotating. This is in good agreement with the results from the ATLAS^{3D} survey, where 86% (224/260) of the ETGs are fast rotators and 14%(36/260) are slow rotators (Emsellem et al. 2011). Due to the lower mass cut for the simulation, the lower percentage of fast rotators in the Magneticum sample is an expected behaviour as the fast rotating regime is preferentially occupied by low mass galaxies. The frequencies of slow and fast rotators in the SAMI Pilot sample are with 15% slow rotators and 85% fast rotators very similar to $\rm ATLAS^{3D}.$ Within the CALIFA and SLUGGS sample the frequencies differ significantly from ATLAS^{3D}: 28% of the CALIFA sample are slow rotating, while 72% are fast rotating. Similarly, 27% of the SLUGGS sample are slow rotators and 73% are fast rotators. The discrepancies among the observational samples are most probably due to the underlying galaxy selection and the associated environmental bias. Especially the SAMI Pilot sample is designed to investigate the kinematical state of ETGs within Clusters. Although the SLUGGS sample covers the full range of environments it only contains 22 ETGs. Even the volume-limited ATLAS^{3D} sample, which covers a volume of 42Mpc, is not representative for the total ETG population in the universe. ATLAS^{3D}, however is a volume and magnitude limited sample aiming to be representative

of the low redshift galaxy population. Hence, for a abundance comparison of fast and slow rotators the ATLAS^{3D} sample is the most sensible choice.

The fast rotating regime is in excellent agreement with the different observational samples. Similar to the observations, we see a well defined upper envelope: as $\epsilon_{1/2}$ increases, the maximum $\lambda_{\mathrm{R}_{1/2}}$ increases accordingly. However, the $ATLAS^{3D}$ sample reveals an envelope reaching higher $\lambda_{\mathbf{R}_{1/2}}$ values for a given $\epsilon_{1/2}$. This can be explained by the applied mass selection criterion, an issue that was already reported in several former studies using isolated merger simulations (Bois et al. 2011; Jesseit et al. 2009) as well as in cosmological zoom-in simulations (Naab et al. 2014; Wu et al. 2014). A similar behaviour is visible when comparing $\mathrm{ATLAS}^{\mathrm{3D}}$ to the remaining observations, where only one ETG exhibits a $\lambda_{R_{1/2}}$ larger than 0.65. Therefore, the discrepancies in the extremely fast rotating range might be due to the different definitions of an ETG applied to classify the galaxies within the different samples, or driven by the different environments included in these studies.

In the slow rotating regime we clearly see the aforementioned resolution limit of $\lambda_{\rm R_{1/2}}$ > 0.07, as the simulation does not reach values lower than 0.07. We find a significant fraction of slow rotators with $\epsilon_{1/2} > 0.4$ which is not present in the observations. Although there are three galaxies observed in this regime, we find a distinctly larger fraction of 3.5% of the total sample in the simulation. This tension with observations was already reported by Naab et al. (2014) based on a sample of 44 zoom-in simulations of individual galaxies (Oser et al. 2010). In addition, high resolution isolated galaxy merger simulations by Bois et al. (2011) and Moody et al. (2014) also produced highly elongated slow rotators, albeit their higher resolution in the simulations, and the different implementation of physics as well as the very different nature of such isolated merger simulations. Thus, the origin of these elongated slow rotators remains to be explored in future work.

3.2 Edge-on Projections and Environmental Dependence

For galaxies that rotate around their minor axis, $\lambda_{R_{1/2}}$ is largest if the galaxy is seen edge-on. The edge-on projection is the plane spanned by the minor and major axis of the stellar distribution. Therefore, also the maximum ellipticity should be obtained under this projection.

We rotate all our galaxies² into edge-on projections, and the result is shown in Fig. 3. The red symbols again mark the *Magneticum* ETGs, while the grey filled circles mark all other galaxies.

As expected, we find that the non-ETGs are much more common at the high- $\lambda_{\rm R}$ end than the ETGs, and that flat ellipticities are also much more frequent for the non-ETGs than for the ETGs. However, we also clearly see that the population of non-ETGs is contaminated by objects currently influenced by interactions: A certain portion of these galaxies do not follow the expected behaviour of high $\lambda_{R_{1/2}}$ and ϵ_e values characteristic for late-type galaxies (Cappellari 2016; Bois et al. 2011). In addition, recent observations revealed that early-type galaxies can posses a significant amount of cold gas, with f_{gas} distinctly larger than 0.1 (Davis et al. 2015), and hence these ETGs would be included in the non-ETG population by our selection criterion. Therefore, the non-ETGs cover the full range of $0.07 \leq \lambda_{R_{1/2}} \leq 0.83$ and $0.08 \leq \epsilon_{1/2} \leq 0.79$. For very low $\lambda_{\rm R}$, nevertheless, the non-ETGs become rare. This is in good agreement with expectations, as the very round and slowrotating galaxies do usually not have that much cold gas. Nevertheless, a substantial amount of the non-ETGs populate the extremely high $\lambda_{R_{1/2}}/\epsilon_{1/2}$ domain, showing the expected trend for late-type galaxies, which can be seen even better in the sideways cumulative distributions of Fig. 3.

Additionally, we separate the *Magneticum* ETGs into centrals (i.e. galaxies that reside at the centre of the potential, filled red circles in Fig. 3) and subhalos (i.e. galaxies that are satellite galaxies within the halo of another galaxy). Furthermore, we split the subhalo ETGs according to the environment they are living in, into cluster satellites (filled red triangles), group satellites (filled red squares), and field satellites (filled red diamonds).

We do not find any difference between the ETGs residing in different environments, and only a slight difference between centrals and satellite ETGs, with a tendency for the extremely slow rotating galaxies to be preferentially central galaxies and not satellites. This is in good agreement with recent studies of environmental dependencies of $\lambda_{R_{1/2}}$, which also could not find any substantial influence of the large scale environment on the kinematical properties of ETGs, as discussed above.

The shaded red areas in Fig. 3 illustrate the number density of the *Magneticum*-ETGs in the $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ plane. They reveal two distinct populations of ETGs, fairly well separated by the magenta line and the observational fast-slow rotator threshold, with only very few galaxies in the transition region.

The population of slow rotators is clearly separated from the upper population, indicating that the different behaviour of slow- and fast rotators in the $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ plane is not just a line drawn visually but actually contains physical meaning. The clustering of galaxies in the vicinity of the black dashed line is caused by the resolution limit³ described in Sec. 2.4 and thus does not contain physical meaning other than that the global kinematic properties of these ETGs are slowly- or even non-rotating.

The fast rotator population splits up into the galaxies in the transition region and those above the magenta line, with the majority of the fast rotators being above that line. Most of these "real" fast rotators are actually residing within the two magenta lines shown in Fig. 3, with only very few ETGs above the upper curve. The nature of these two curves will be discussed in detail in the following subsection.

 $^{^2\,}$ All galaxies that satisfy the resolution criterion are treated like this, independent of further classifications

 $^{^3}$ Estimating the effect of an increased mass resolution is not trivial: In general an increase in resolution reduces the statistical noise allowing to reach lower $\lambda_{\rm R_{1/2}}$ values. It is however not possible to predict the impact of a more resolved velocity distribution on $\lambda_{\rm R_{1/2}}$ due to the increased number of particles, which could for example reveal small kinematical distinct cores in the centers which would lead to larger measured $\lambda_{\rm R_{1/2}}$ values.



Figure 2. Comparison of the Magneticum ETGs with ATLAS^{3D}, CALIFA, SLUGGS, and SAMI Pilot observations in the $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ plane. Main panel: Filled circles indicate the Magneticum ETGs, whereas the open blue circles mark the observations as indicated in the legend. The green line defines the threshold between fast and slow rotators. The magenta line shows the theoretical position of edge-on viewed ellipsoidal galaxies with an anisotropy parameter $\delta = 0.7 \times \epsilon_{intr}$ (for further details see Cappellari et al. (2007)). According to the dashed magenta line represents the theoretical position for edge-on projected isotropic galaxies with $\delta = 0$. The number density in the plane is illustrated by the red squares. Upper and right panel: Cumulative number of galaxies (N_{CUM}) normalised by the total number of galaxies (N_{TOT}) of the respective sample for $\lambda_{R_{1/2}}$ and $\epsilon_{1/2}$, respectively.

This observed segmentation suggest a fundamental difference between the slow- and the fast rotating population of ETGs, with only few ETGs in a transition state between them. Thus, it supports the idea that there are distinct formation histories for the fast- and slow rotating populations, and not a slow, continuous transformation from one population to the other.

3.3 Global Anisotropy

Within the theoretical framework provided by Binney (2005), Cappellari et al. (2007) derived a theoretical connection between the anisotropy

$$\delta = 1 - \frac{\Pi_{zz}}{\Pi_{xx}} = 1 - \frac{\sum_{i=1}^{N} M_i \ \sigma_{z,i}^2}{\sum_{i=1}^{N} M_i \ \sigma_{x,i}^2},$$
(11)

 V/σ and the intrinsic ellipticity $\epsilon_{\rm intr}$ from first principles, i.e. the tensor virial theorem, under the assumption of a flattened, axisymmetric galaxy in the edge-on projection. Furthermore, Emsellem et al. (2011) found a tight correlation between $\lambda_{\rm R_{1/2}}$ and V/σ which allows a translation of the theoretical connection into the $\lambda_{\rm R_{1/2}}$ - $\epsilon_{1/2}$ plane:

$$\lambda_{\mathrm{R}_{1/2}} \approx \frac{k \ (V/\sigma)}{\sqrt{1 + k^2 \ (V/\sigma)^2}} \tag{12}$$

$$(V/\sigma) \approx \sqrt{\frac{(1-\delta) \ \Omega(\epsilon_{\rm intr}) - 1}{0.15 \ (1-\delta) \ \Omega(\epsilon_{\rm intr}) + 1}}$$
(13)

where k is estimated to be ≈ 1.1 . The solid magenta line in Fig. 2 and Fig. 3 represents the result for this theoretical model under the assumption of a linear connection between



Figure 3. Main Panel: $\lambda_{R_{1/2}} - \epsilon_{1/2}$ plane for *Magneticum* galaxies with $M_* > 2 \cdot 10^{10} M_{\odot}$ in the edge-on projection. Colours separate the *Magneticum* ETGs (red) and the rejected galaxies (grey) according to Sec. 2.2. The *Magneticum* ETGs are furthermore plit up according to their large scale environment into cluster satellites (filled red triangles), group satellites (filled red squares), field satellites (filled red diamonds), and centrals (filled red circles). The green line defines the threshold between fast and slow rotators given in Eq. 10. The magenta line shows the theoretical position of edge-on viewed ellipsoidal galaxies with an anisotropy parameter $\delta = 0.7 \times \epsilon_{intr}$ (for further details see Cappellari et al. (2007)). Accordingly the dashed magenta line represents the theoretical position for edge-on projected isotropic galaxies with $\delta = 0$. The number density in the plane is illustrated by the red squares. Side panels show the cumulative number of galaxies (N_{CUM}) normalised by the total number of galaxies (N_{TOT}) of the respective sample split up into different environments.

 δ and ϵ_{intr} with a factor 0.7, while the dashed magenta line shows the theoretical result for an isotropic ($\delta = 0$) galaxy. Therefore, these two magenta curves define a generic envelope for the population of "real" fast rotators. This is in substantial agreement with the findings of Cappellari et al. (2007); Cappellari (2016) who showed that for regular rotating ETGs δ lies in the interval $0 \leq \delta \leq 0.7 \cdot \epsilon_{intr}$.

In addition to the anisotropy from this model, that can be assigned to each ETG (δ_{model}), we also calculate the "real" anisotropy from the particle distribution for each ETG (δ_{calc}). We find that δ_{model} and δ_{calc} are not identical for nearly all galaxies, but that δ_{model} usually is lower than δ_{calc} , but by a more or less constant factor (see Fig. A1 in App. A). We show the $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ -plane, with symbols coloured according to their anisotropy δ_{calc} , in Fig. 5, with dotted lines in different colours marking the galaxies δ_{model} according to the model. This does not only show the shift between the two anisotropies, but it also clearly demonstrates that there is no correlation between the difference in the two anisotropies and the position of the galaxy within the $\lambda_{\text{R}_{1/2}}$ - $\epsilon_{1/2}$ -plane.

We also cannot find a change in the anisotropy distribution of the ETGs with redshift, as demonstrated for the fast rotator population in Fig. 4. Only at z = 2, there are slightly higher anisotropies found, albeit that is also the smallest sample of ETGs and this might thus only be a statistical effect.



Figure 4. Anisotropy distribution of fast rotators at different redshifts as indicated in the left upper corner.



Figure 5. $\lambda_{\text{R}_{1/2}}$ - $\epsilon_{1/2}$ plane colour coded according to δ_{calc} . The dashed lines show the result for the theoretical model assuming different δ_{model} as given in the legend.

3.4 Redshift evolution

While we do not see strong evolution trends with redshift in the anisotropy of our ETGs, we find a clear evolution of the $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ plane with redshift: Fig. 6 shows the $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ plane for redshifts from z = 2 to z = 0.

The significant increase in the total number of ETGs towards higher redshifts confirms that most galaxies are still gas-rich at z = 2, and quenching mechanism have a substantial impact blow this redshift. At z = 2, most of the galaxies are in the vicinity or above the magenta relation, while only a very small fraction is classified as slow rotators. Already at this stage of the evolution, only a small fraction of ETGs exceed the isotropic relation. Therefore, the fast rotating population is already in place at z = 2, while the slow rotating population only begins to form. All of the subsequent redshifts show a well defined fast rotating population, well constrained by the magenta curves.

As the redshift decreases, the slow rotator population becomes statistically more significant. Therefore, we conclude that the mechanism which leads to the formation of the slow rotators, starts to take effect below z = 2. This is consistent with the results from Naab et al. (2014) who found that particularly (multiple minor) merger events after z = 2 drive the formation of slow rotating ETGs.

The detailed evolution of $\lambda_{R_{1/2}}$ with redshift can be seen in Fig. 7, which shows the histograms of $\lambda_{R_{1/2}}$ for the four considered redshifts. It clearly confirms that the slow rotating population only begins to build up at z = 2, and gets more prominent towards z = 0: At z = 2, the distribution exhibits a clear peak in the range $0.5 < \lambda_{R_{1/2}} < 0.7$, with a decreasing tail towards lower values. Subsequently, at z = 1 a peak at $\lambda_{R_{1/2}} \approx 0.1$ close to the resolution limit emerges. In the further evolution, more galaxies drop to the slow-rotator regime, enhancing this peak at the resolution limit value.

In the high- $\lambda_{R_{1/2}}$ regime, we observe a continuous shift of the peak towards lower $\lambda_{R_{1/2}}$, implying a general spindown of the complete fast rotating population with decreasing redshift. Due to a simultaneous decrease of $\epsilon_{1/2}$, as seen in Fig. 11, the complete fast rotating population shifts approximately within the region constrained by the magenta curves. Therefore, the most dominant effect in the evolution of the upper population decreases both $\lambda_{R_{1/2}}$ and $\epsilon_{1/2}$ gradually, while the anisotropies do not change, indicating that this mechanism leads to an enhancement of the velocity dispersion while simultaneously shaping the ETGs to be more spherical. The most likely mechanism to be responsible are multiple merger events, again supporting the previously discussed growth mechanisms for ETGs.

Interestingly, the transition region between the fast and slow rotator branch, found in Fig. 3, is present at all redshift. This suggests that ETGs, which change their kinematical flavour from "real" fast rotation to slow rotation, pass through this transition region rapidly. In order to investigate this in more detail we followed the evolution of $\lambda_{\rm R_{1/2}}$ for all slow rotators individually. We find that more than 50% of all slow rotators are formed in a quite short and distinguishable transition from fast to slow rotators. A closer inspections shows, that at least 30% of all slow rotators at z = 0 unambiguously underwent this transition on a timescale of less than 0.5Gyr, and exclusively are associated with a significant merger in the range between 1:5 and 1:1, as shown Fig. 8.

On a final side note: Due to several quenching mechanisms and galactic merger events, galaxies can change their morphologies between the considered redshifts from discy to spheroidal. Furthermore, it is theoretically possible for galaxies to re-accrete fresh cold gas and hence become discy again. To ensure that the evolution trends found with redshift are not driven by the underlying classification criteria used a the different redshifts, we show the $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ evolution with redshift for all galaxies in the *Magneticum* simu-



Figure 6. Redshift evolution of the *Magneticum* ETGs in the $\lambda_{R_{1/2}} \epsilon_{1/2}$ from z = 2 (upper left panel) to z = 0 (right lower panel). Lines and symbols are the same as in Fig.3. The colour of the symbols is according to the stellar mass as indicated in the left upper panel.

lation in App. C. As can be seen from Fig. C1, all reported redshift trends are physical and do not depend on the selection criteria, and are similarly present also for disk-like galaxies.

4 CONNECTING MORPHOLOGY AND KINEMATICS

4.1 The $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ and the M_* - j_* Plane

We want to connect the results deduced in the previous section to the fundamental classification introduced by Fall (1983) and reviewed in Romanowsky & Fall (2012). When investigating a plane spanned by the stellar specific angular momentum $\log_{10}(j_*)$ and the total stellar mass $\log_{10}(M_*)$, they found a continuous linear sequence of morphological types. Furthermore, they found that LTGs and ETGs follow a parallel sequence with a slope of approximately 2/3, implying a power law relation in the non-log plane. This result was confirmed for a larger sample of galaxies within one effective radius by Cortese et al. (2016). Furthermore these findings are in agreement with recent predictions from cosmological simulations (Pedrosa & Tissera 2015; Teklu et al. 2015; Genel et al. 2015). Especially Teklu et al. (2015) performed a rigorous investigation of the M_* - j_* plane, considering the same simulated box as this study, introducing the so-called "b-value" defined as

$$b = \log_{10} \left(\frac{j_*}{\text{kpc km/s}} \right) - \frac{2}{3} \log_{10} \left(\frac{M_*}{M_{\odot}} \right)$$
(14)

to effectively parametrise the position of a galaxy in the plane, with j_\ast given by

$$j_* = \frac{\left|\sum_{j=1}^N m_i \ \vec{r}_i \times \vec{v}_i\right|}{\sum_{j=1}^N m_i}.$$
 (15)

This *b*-value is by definition the *y*-intercept of a linear function with slope 2/3 in the $\log_{10}(M_*)$ - $\log_{10}(j_*)$ plane. As shown by Romanowsky & Fall (2012) and Teklu et al. (2015), at z = 0 objects with $b \approx -4$ are disc-like galaxies, followed by a smooth transition to lenticular and elliptical galaxies with decreasing *b*-value. Therefore, this parameter represents a fundamental connection between three parameters



Figure 7. Statistical distribution of $\lambda_{R_{1/2}}$ presented as histograms of the relative frequency. The colour distinguishes four redshift as given in the legend.



Figure 8. Upper Panel: Temporal evolution of the stellar mass M_* for individual slow rotators which show a rapid decline in $\lambda_{R_{1/2}}$ associated with a merger event in the range between 1:5 and 1:1 representing 30% of all slow rotators. M_* is normalised to the average stellar mass after the drop in $\lambda_{R_{1/2}}$ ($M_{*,drop}$). The time axis is fixed such that the zero point represents the time of the drop. Lower Panel: Temporal evolution of $\lambda_{R_{1/2}}$ demonstrating the rapid decline.

describing a significant part of galaxy formation: the total stellar angular momentum j_* , the total stellar mass M_* , and the present-day morphology. The striking implication of the slope of 2/3 is that, for a given specific angular momentum, which is per definition normalised by the total stellar mass, all different morphologies are possible. We use the *b*-value as a tracer for the morphology within the *Magneticum* ETGs.

Fig. 9 shows the edge-on $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ plane coloured according to *b*-value. For the calculation of M_* and j_* we used all stellar particles within a sphere of radius $3R_{1/2}$ centred on



Figure 9. Edge-on $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ plane for the *Magneticum* sample with colours indicating the *b*-value as indicated in the colourbar. The lines are identical to the ones in Fig. 3.

the galaxy's centre of mass. The mean value for b for the fast rotators is $\overline{b} = -4.7$, while for the slow rotators $\overline{b} = -5.3$, clearly showing that the lower population is dominated by classical spheroidal galaxies. The transition galaxies have a mean *b*-value of $\overline{b} = -4.9$, consistently between the two other populations. This confirms the evolution towards less disc-like galaxies with decreasing $\lambda_{\text{R}_{1/2}}$, as already found in observations and for the progenitors of isolated merger simulation (Bois et al. 2011; Krajnović et al. 2013; Falcón-Barroso et al. 2015).

For the upper population there is a clear trend for the b-value to increase with raising $\lambda_{R_{1/2}}$ and $\epsilon_{1/2}$. The colour gradient seems to evolve continuously along a linear relation indicating a transition from more elliptical-like galaxies towards disc-like morphologies with increasing $\lambda_{R_{1/2}}$ and $\epsilon_{1/2}$. This behaviour suggest also a continuous sequence of accretion histories along the gradient in b, assuming galaxies to usually have high j_* and a disk-like morphology at high redshift. In general, mergers diminish the angular momentum in a moderate fashion while enhancing the stellar mass (Lagos et al. 2017), leading to a simultaneous decrease of b and $\lambda_{\mathbf{R}_{1/2}}$, creating the vertical evolution visible in Fig. 9 (see also Choi & Yi (2017)). Such an effect of merging was already suggested by Romanowsky & Fall (2012) to move within the M_* - j_* Plane. Furthermore, mergers lead to higher dispersions, resulting in a puffed up galaxy with smaller ellipticity responsible for the horizontal evolution. Therefore, within this scenario, the evolution in b is driven by the amount of merging a galaxy has undergone during its recent assembly. Of course, the path of a galaxy in the parameter space spanned by $\lambda_{R_{1/2}}$, b and $\epsilon_{1/2}$ strongly depends on the configurations of the mergers involved in the formation.

A similar behaviour, however much weaker, is visible for slow rotators: galaxies with the smallest and therefore most

elliptical-like \boldsymbol{b} are located in the vicinity of the resolution limit.

To investigate the connection between $\lambda_{R_{1/2}}$ and b in more detail, Fig. 10 displays their direct correlation colour (coded by the half-mass radius). It elucidates the conclusion that $\lambda_{R_{1/2}}$ and b are directly correlated. The distribution is well confined by an upper and lower envelope. For $\lambda_{R_{1/2}} \gtrsim 0.3$, the correlation follows a almost linear relationship with a certain scatter. Below $\lambda_{R_{1/2}} \lesssim 0.3$ the dependency gets flatter until resolution effects get significant. Interestingly, this is approximately the threshold between the upper population and the transition population.

Investigating the distribution of $R_{1/2}$ we find that galaxies exhibiting the same $R_{1/2}$ lie on separated relations parallel shifted to the residual $R_{1/2}$ with a continuous transition from lower $R_{1/2}$ at lower b to larger $R_{1/2}$ at higher b. Therefore for a fixed b galaxies exhibiting a higher $\lambda_{R_{1/2}}$ have smaller $R_{1/2}$. This correlation is especially prominent for fast rotating ETGs with $\lambda_{R_{1/2}} > 0.2$, while in the slow rotating regime the correlation is not present. Therefore, the mechanism driving this correlation is only effective for fast rotating ETGs indicating a different formation pathway for fast and slow rotators.

To further explore the functional form of the relation we use a simplified toy model. The model is based on the following assumptions:

• The stellar density follows a Hernquist profile (Hernquist 1990) given by

$$\rho(r) = \frac{\rho_0}{\frac{r}{r_s} \left(1 - \frac{r}{r_s}\right)^3}.$$
(16)

• The velocity and position vector of the stellar particles are perpendicular to each other $\vec{v} \perp \vec{r}$. This conditions constrains the particles to be on circular orbits, however, the planes of rotation can be tilted with respect to each other.

• The rotational velocity v_{rot} and the velocity dispersion σ are spatially constant and are connected by:

$$V_{vir} = \sqrt{v_{rot}^2 + \sigma^2}.$$
 (17)

This equation states that the total energy in the system obtained from the virial theorem is distributed between ordered and random motion. This relation, however, only holds for systems in virial equilibrium.

• We neglect the projection effects on v_{rot} , σ and $\rho(r)$ occurring in the calculation of $\lambda_{R_{1/2}}$. This is probably the most error-prone assumption.

• We use the mass-size relation determined for the *Magneticum* ETGs in Sec. 2.3.

Applying these assumptions, $\lambda_{R_{1/2}}$ simplifies considerably

$$\lambda_{\mathbf{R}_{1/2}} = \frac{\int_{\partial A} \rho_p(r) r_p | v_{rot,p} | dA}{\int_{\partial A} \rho_p(r) r_p \sqrt{v_{rot,p}^2 + \sigma_p^2} dA} \approx \frac{|v_{rot}|}{\sqrt{v_{rot}^2 + \sigma^2}}, \quad (18)$$

where the subscript p indicates the projected quantities. To calculate b, it is necessary to integrate the stellar angular momentum and the density over the considered volume

$$b = \log_{10} \left(\int_{\partial V} \rho(r) |\vec{r} \times \vec{v}| dV \right) - \frac{2}{3} \log_{10} \left(\int_{\partial V} \rho(r) dV \right).$$
(19)

The only free parameters we have to feed into the model are



Figure 10. Relation between $\lambda_{R_{1/2}}$ and *b*. The colour corresponds to the half-mass radius as given in the colourbar. The black curve represents the theoretical prediction of the model outlined in Sec. 4.1 for a galaxy with $M_* = 5 \cdot 10^{10} M_{\odot}$. The grey shaded area indicates the 1σ error adopted from the scatter in the mass-size relation. The dashed black line marks the resolution limit for $\lambda_{R_{1/2}}$.

 M_* and v_{rot} . The result for $M_* = 5 \cdot 10^{10} M_{\odot}$ with varying v_{rot} in the range $[0, V_{vir}]$, and therefore from a completely dispersion dominated system to a purely rotational supported system, is shown as a black curve in Fig. 10. The grey shaded area marks the 1σ scatter adopted from the mass-size relation. The impact of this error suggests that the trend for the scatter with radius is a translation of the scatter in the mass-size relation into this correlation. It is remarkable that this extremely simplified model can already reproduce the generic functional form visible in Fig. 10. Down to the resolution limit the black curve resembles the lower and upper envelope for the total distribution seen from Magneticum as well as the intrinsic shape for a given $R_{1/2}$.

To investigate whether the correlation found in Fig. 10 is already established at higher redshifts, Fig. 11 displays its evolution with redshift.

The correlation between $\lambda_{R_{1/2}}$ and b is already present at z = 2, however the tail towards lower b is not established yet. At z = 1, the tail region starts to be populated, and becomes more prominent at the subsequent redshifts. Generally, we find a clear trend that the low b-values are established parallel to the appearance of the slow-rotator population in ETGs, in line with the results presented in Fig. 6 & 7. In the fast rotator regime the spin-down in $\lambda_{R_{1/2}}$ is visible, while also a shift towards higher b is present. The shift in b with redshift for disc galaxies was also found by Teklu et al. (2016) in the Magneticum simulation, showing the agreement with the theoretical model presented in Obreschkow et al. (2015).

In the lower right panel of Fig. 11 we include observa-

14 Schulze et al.

tions by the SAMI survey (filled diamonds, Cortese et al. (2016)) and a cross-match of ATLAS^{3D} data ($\lambda_{R_{1/2}}$) and data obtained from Romanowsky & Fall (2012) (R12, *b*-values) shown as open diamonds. In order to obtain the *b*-values for the SAMI data we extracted stellar masses and the inclination corrected stellar specific angular momentum for the elliptical and S0 galaxies from Fig. 2 in Cortese et al. (2016), while $\lambda_{R_{1/2}}$ is extracted from Fig. 7. The SAMI data is in excellent agreement with the *Magneticum* ETGs as well as with our theoretical toy model. The slight disagreement found in both data sets is most probably caused by inclination effects, since only the stellar angular momentum for the SAMI data is inclination corrected.

At z = 2 the ETGs are very compact, as minor mergers as the main driver for an increase of $R_{1/2}$ get more prominent at lower redshifts. This is reflected by the occurrence of more extended ETGs in the further evolution down to z = 0. The growth of $R_{1/2}$ with decreasing redshift is in agreement with the observational findings by Huertas-Company et al. (2013). From the evolution of the colour gradient and the width of the distribution we can furthermore conclude that the scatter increases significantly between 2 < z < 1, followed by a modest increase with decreasing redshift. We checked the evolution of the mass-size relation with redshift and found the same trend in the scatter (see also Remus et al. (2017) for the redshift evolution of the mass-size relation for ETGs in *Magneticum*). This supports the idea that the scatter in the $\lambda_{\mathbf{R}_{1/2}}$ -b correlation is a reflection of the scatter in the mass-size relation.

4.2 Connecting Morphological and Kinematical Properties

Another tracer of a galaxies's morphology is the Sérsic-Index n, since it describes the curvature of the Sérsic-profile fitted to the radial light distribution. The details of the fitting procedure is outlined Sec. 2.5. We investigate the correlation between the Sérsic-Index and the two parameters $\lambda_{R_{1/2}}$ and b.

Fig. 12 shows the direct correlation between $\lambda_{R_{1/2}}$ and the Sérsic-Index, while the colour encodes *b*. We find no clear correlation between *n* and $\lambda_{R_{1/2}}$, in agreement with observational results found for the ATLAS^{3D} sample presented in Krajnović et al. (2013) for a one component Sérsic fit. In a more recent study Cortese et al. (2016), within the SAMI project, speculate that the scatter in this relation also found in their own sample especially in the slow rotating regime is mainly due to inclination effects. Since we are able to exclude inclination effects by using the edge-on projection, we can disprove this speculation and conclude that there is no significant correlation between $\lambda_{R_{1/2}}$ and the Sérsic-Index. Hence, the Sérsic-Index as a morphological parameter of a galaxy is not a sufficient tracer for the kinematical state of a galaxy⁴.

⁴ However, a direct comparison of Sérsic-Indices obtained from simulations and observations has to be interpreted with caution, as they probe different radial regimes of the galaxies, since simulations do not resolve the innermost part of the galaxies while observations have difficulties in detecting the faint, low-luminosity outskirts of the galaxies. We also do not find a correlation between b (see upper panel of Fig. 13) and the Sérsic-Index. A weak, shallow, linearly decreasing trend is visible for galaxies with b > -5.0while in the low-b range absolutely no correlation is present. The lower panel of Fig.13 explains the origin of the described behaviour by colouring fast (blue) and slow (red) rotating galaxies. A clear trend is visible for fast rotating galaxies with high n to possess a lower b in the more elliptical branch in line with what is found in Cortese et al. (2016). In contrast, the slow rotators populate the diffuse low-b region, causing the uncorrelated part of the distribution. This is again demonstrating the fundamental difference between fast and slow rotators, populating different regions of the b-n-plane.

The black symbols in the lower panel of Fig. 13 depict observational results from the SAMI survey (filled diamonds, Cortese et al. (2016)) and R12 (open diamonds). In the SAMI data we again only include elliptical and S0 galaxies. The *Magneticum* ETGs are in good agreement with the SAMI observations, while a significant fraction of the R12 data shows higher *b*-values. The disagreement with the R12 observations is most probably due to projection effects, since the angular momentum is not corrected for inclination.

5 KINEMATICAL GROUPS

Motivated by the fundamental findings of Krajnović et al. (2011) we investigate the specific kinematical features of the line-of-sight velocity maps. The authors in that observational study define six kinematical groups based on features in the velocity maps (see Sec. 1 for the definition of the kinematical groups) and investigate their connection to internal properties of ETGs.

5.1 Classification and Group Frequency

The classification method applied in this study rests upon a visual inspection of each individual velocity map in the edgeon projection. The class of 2σ galaxies is not considered. To reduce possible bias, each velocity map was classified independently by three persons. Discrepancies were solved by either assigning the class with two votes to the galaxy or, in the case of a complete disagreement, a collective reconsideration. The galaxies were classified into the following groups:

• *Regular Rotator (RR)*: The velocity map shows a well defined, ordered rotation around the minor axis, with no kinematical features.

• Non Rotator (NR): The velocity map exhibits no distinct kinematical feature and low-level velocities.

• Distinct Core (DC): The velocity map features a central rotating component surrounded by a low-level or nonrotating component.

• *Kinematically Distinct Core (KDC)*: The velocity map shows a central rotating component surrounded by a region with inclined rotation with respect to the central component. This explicitly includes counter-rotating cores.

• *Prolate Rotator (PR)*: The velocity map shows ordered rotation around the major axis of the galaxy.

• Unclass (U): The galaxy cannot be assigned to any of the previous groups.



Figure 11. The redshift evolution of the correlation between $\lambda_{R_{1/2}}$ and *b*. Each panel corresponds to one redshift as indicated in the left upper corner. The colour of the symbols marks the half-mass radius of each galaxy in comoving coordinates as given in the colourbar. In the lower right panel the black diamonds show observations from the SAMI survey (filled diamonds, Cortese et al. (2016)) and a combination of ATLAS^{3D} data ($\lambda_{R_{1/2}}$) and data obtained from Romanowsky & Fall (2012) (R12, *b*-values) shown as open diamonds. The black dashed line in each panel represents the resolution limit for $\lambda_{R_{1/2}}$, while the thick dashed line shows the relations obtained from our toy model.

For each class, Fig. 14 displays an example of a voronoibinned line-of-sight velocity map, except for the U class. The statistical frequency of each class is summarised in a pie chart in Fig. 15. In agreement with observations, the RR are the most common group in the *Magneticum* sample (69.8%) followed by the NR (13.0%). 3.3% of the *Magneticum* ETGs are assigned to the class of DC corresponding to 26 objects. The reason for the low number of five KDCs (1%) is most probably a resolution issue, since KDCs in ATLAS^{3D} are located in the very centre of the galaxy, which is not resolved in the simulation.

The group of PR contains 17 galaxies, which corresponds to 2.2% of the total sample. Preliminary results from the M3G-project using the MUSE instrument suggest that at stellar masses above $10^{11.5} M_{\odot}$ those galaxies are more

frequent. The PRs in our sample follow a similar massdistribution, being more prominent at higher masses, with a significant increase aroung $10^{11.5} M_{\odot}$ (see middle panel of B1). This is in good agreement with the results from ATLAS^{3D} and CALIFA sample which show a rise in the prolate rotator fraction in the same mass regime, reported in Tsatsi et al. (2017). Interestingly, the number of PRs is rather constant over the complete mass range covered by the *Magneticum* ETGs (see upper panel of B1). This implies, that the increase in the fraction of PRs with higher mass is due to the decrease of the number of other objects. This is in agreement with results found in the Illustris simulation (Ebrová & Lokas 2017). Thus, we conclude, that *Magneticum* represents a powerful tool to investigate their formation and compare to upcoming results from M3G, MAS-



Figure 12. Correlation between Sérsic-index n and $\lambda_{\rm R_{1/2}}.$ The colour indicates the b-value.



Figure 13. Correlation between Sérsic-index n and b-value. Upper panel: Colours according to $R_{1/2}$. Lower panel: Colours code fast rotators (blue filled circles) and slow rotators (red filled circles). Black filled diamonds show observations from the SAMI survey extracted from Cortese et al. (2016), while the open diamonds represent observations from R12.

SIVE, CALIFA and MaNGA, which will be addressed in a future paper.

A direct comparison to the results by Krajnović et al. (2011) is difficult due to the differences in the underlying classification mechanism. Especially the group of PR is not included in the ATLAS^{3D} classification. Furthermore, many of the kinematical features found in the ATLAS^{3D} reside within $0.5R_{1/2}$, which, for various of the *Magneticum* ETGs, is in the region where spatial resolution issues become rel-



Figure 14. Example velocity maps for the 5 kinematical groups defined in Sec. 5 from top to bottom: Regular Rotator, Non Rotator Distinct Core, Kinematically Distinct Core, and Prolate Rotator. Each map is voronoi binned to ensure a proper number of particles per bin. The scaling of the velocity is shown in the colour bar. The shown areas encompasses one half-mass radius, and the galaxies are rotated such that the long axis is along the x-direction. The solid black lines represent contours of constant density.



Figure 15. The statistical distribution of the different kinematical groups within the *Magneticum* ETGs: red are Regular Rotators, blue are Non Rotators, turquoise are Distinct Cores, yellow are Kinematically Distinct Cores, and lilac are Prolate Rotators. Unclass are shown in black.

evant and it is not possible to detect features which are smaller than the spatial resolution of the velocity map. In order to explore the impact of mass resolution on the velocity maps we applied a bootstrapping algorithm to galaxies with the smallest amount of stellar particles. It revealed that the kinematical features in our sample are not sensitive to the particle binning.

5.2 Kinematical Groups and Global Galaxy Properties

The distribution of the five kinematical groups in the $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ plane is shown in the central panel of Fig. 16 for the edge-on projection, while the sideways panels display cumulative distributions of $\lambda_{\mathrm{R}_{1/2}}$ and $\epsilon_{1/2}$ with a different linestyle for each group. As expected the NR they exclusively populate the low- $\lambda_{\mathrm{R}_{1/2}}$ regime, while spanning a wide range of $\epsilon_{1/2}$ with a peak at $\epsilon_{1/2} \approx 0.35$ and are slow rotating. The RRs populate the fast rotating regime, with only a small number of objects in the slow rotating region. The slow rotating RRs show a RR velocity pattern, however their dispersion maps reach higher values than those of the fast rotating RRs, with a comparable rotational velocity, effectively causing the low $\lambda_{R_{1/2}}$ -values. Four of the five KDCs found in the Magneticum ETGs are, in agreement with $\rm ATLAS^{3D}$ observations, classified as slow rotators, while one KDC lays close to the threshold.

For the DCs, the distribution is not as obvious as for the other classes. They span the complete $\lambda_{R_{1/2}}$ range up to ≈ 0.5 , with a larger fraction in the fast rotating regime. The reason for the wide spread within this class is the strong difference in the spatial extend of the rotating component: since the non-rotating surrounding is causing a lower $\lambda_{R_{1/2}}$ value the total $\lambda_{R_{1/2}}$ decreases with smaller radial extend of the rotating component. The DCs with the slowest rotation have small rotating discs in the centre, with an extent of only a third of $R_{1/2}$.

Except for two members, all PRs are located in the slow rotating regime. This is an unexpected behaviour since per definition the stars show a distinct rotation around a common axis. As the formation of those objects is still a matter of debate, we can only speculate about the origin of the low $\lambda_{R_{1/2}}$ -values. From basic physics it is clear that smoothly accreted gas cooling to the centre of a halo always forms a disk rotating around the minor axis. Therefore, the most probable mechanisms to form prolate rotation are galaxy mergers or a flyby interaction with another galaxy. In the context of dwarf spheroidal galaxies, Ebrová & Łokas (2015) showed that mergers of two dwarf disk galaxies can indeed result in prolate rotation, using a collisionless N-Body simulation. They found the most significant prolate rotation in the remnant of a nearly face-on merger of two identical disk dwarfs on a perfectly radial orbit. In a pure N-Body simulation with this settings, the prolate rotation is a consequence of angular momentum conservation around the merger axis. The possibility to form PRs in binary N-Body simulations of more massive non-dwarf galaxies is shown in Tsatsi et al. (2017), where the authors form PRs in a polar merger with the amplitude of prolate rotations depending on the bulgeto-total ration of the progenitors. Of course, the scenario gets significantly more complicated when involving a gaseous component and allowing for non-radial orbits in a fully cosmological environment (Ebrová & Lokas 2017). Therefore, a further analysis regarding the properties and formation pathway of those objects has to be conducted to disentangle their origin.

To further explore the properties of the kinematical groups, Fig. 17 shows the three correlations investigated in the previous sections, split up into the kinematical groups. The upper left panel shows the between $\lambda_{\rm R_{1/2}}$ and the Sérsic-index. None of the groups shows a correlation between those parameters. Except for the KDC, which have n in the range 1 < n < 4, all groups populate a large range in n. Hence, the formation of the various kinematical features do not leave a signature in the stellar density profile out to large radii. We want to allude, that the region where the defining features for the kinematical groups are located is explicitly excluded from the fitting range. Therefore we can not exclude to find group specific features in the stellar density profile when investigating only the central region within one half-mass radius.

In the correlation between n and b (lower left panel of Fig. 17) we see that the uncorrelated lower region is build up by the NR,DC,KDC and PR, while the correlation is build up by the RR. We can furthermore conclude, that RR exhibit more disc-like *b*-values than all the other kinematical groups, while the *b*-value cannot be used to distinguish the other kinematical groups from each other.

This is also visible in the upper right panel where the connection between b and $\lambda_{R_{1/2}}$ is displayed. The only group which populates a distinct region in this plane are the RRs. Interestingly, the special group of PRs covers a large range in b, not reaching the high disc-like values.

Therefore, we conclude that it is not possible to disentangle the kinematical groups using global galaxy parameter. This suggests that the signature of their respective for-



Figure 16. Main Panel: The $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ plane for the *Magneticum* ETGs subdivided into the kinematical groups defined in Sec. 5 at z = 0: red circles are PR, blue triangles are NR, turquoise rectangles are DC, yellow diamonds are KDC, and lilac bow ties are PR. The lines are the same as in Fig. 3. Side panels show the cumulative number of galaxies (N_{CUM}) normalised by the total number of galaxies (N_{TOT}) of the respective sample split up into the various kinematical groups.

mation histories are encoded in local parameters restricted to the central region of the galaxy.

5.3 Kinematical Groups at z = 2

For the 157 galaxies classified as ETGs at z = 2, we apply the same classification according to the kinematical features as for the ETGs at z = 0. From these 157 ETGs, 154 had maps resolved enough for such classifications, while only 3 could not be classified. Fig. 18 shows the $\lambda_{R_{1/2}}$ - $\epsilon_{R_{1/2}}$ plane for the ETGs at z=2, with different symbols and colours marking the same kinematical groups as before. Most of the ETGs at z = 2 are regular rotators (92%), and nearly all fast rotators belong to this kinematical group. Only two fast rotators do not classify as regular rotators but have a clearly visible decoupled core. Both galaxies with decoupled cores live in the transition region, however, it is not possible to conclude from only two objects a "typical" behaviour.

As can clearly be seen, all galaxies (but one) that belong to the slow rotator regime are classified as non-rotators (4%), demonstrating that the split-up between non-rotators and fast-rotators is already established at z = 2. The single slow rotator that is not a non-rotator is the only galaxy at z = 2 that exhibits a prolate rotation. We trace this galaxy from z = 2 to z = 0 and find that the prolate rotation of this object is present at all redshifts, and that its $\lambda_{\text{R}_{1/2}}$ does not change much with time. In our sample at z = 2, there is no galaxy with a kinematically distinct core, however, these galaxies are also rare at z = 0 and thus the lack of such galaxies in our box at z = 2 does not indicate that such galaxies cannot exist at z = 2.

We conclude that the most common kinematical classes are already present at z = 2, and that regular rotators are more frequent at higher redshifts, in agreement with the current picture of non-rotator formation. Furthermore, we already find special kinematic features like decoupled cores and prolate rotation at z = 2, proving that the formation pathways that produce such kinematical features can also occur at redshifts higher than z = 2. As all our ETGs have gas fractions below $f_{gas} = 0.1$, none of these galaxies classify



Figure 17. Various parameter correlations split up into kinematical groups. Left upper panel: Correlation between $\lambda_{R_{1/2}}$ and Sérsic index. Right upper panel: Correlation between $\lambda_{R_{1/2}}$ and b. Left lower panel: Correlation between b and Sérsic index. In the upper row the black dashed line marks the resolution limit for $\lambda_{R_{1/2}}$.

as blue nuggets, but several of them have rather small halfmass radii and as such classify as red nuggets.

6 DISCUSSION AND CONCLUSION

Recent observations revealed a shift in the existing paradigm for early-type galaxies, away from the classical morphological separation into elliptical galaxies and S0 mostly driven by the observed ellipticities, towards a classification based on fundamental kinematical properties. The picture of earlytype galaxies to be kinematically unimpressive has been revolutionised by observing a richness of complex kinematical structures, suggesting a variety of formation histories.

In this study we investigated the kinematical properties of a sample of 784 early-type galaxies at z = 0 extracted from the *Magneticum Pathfinder* simulation, which are a set of hydrodynamical simulations performed with the Tree/SPH code GADGET-3. We selected all galaxies with $M_* > 2 \cdot 10^{10} M_{\odot}$ and a cold gas fraction below $f_{\rm gas} < 0.1$, to ensure maximal comparability to the galaxy selection process adopted for the observations. Furthermore, we follow the evolution of the properties of ETGs from z = 2 to z = 0in a statistical manner applying the same selection criteria.

To characterise the kinematical state of the *Magneticum* ETGs, voronoi-binned line-of-sight velocity maps were constructed and $\lambda_{R_{1/2}}$ is calculated for all galaxies. In addition, we investigate our sample with regard to various parameters characterising ETGs like the *b*-value, anisotropy, ellipticity, and Sérsic-index.

In the first step we compare the distribution of our sample at z = 0 in the $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ plane to the distribution found by ATLAS^{3D}, CALIFA, SLUGGS and SAMI Pilot. With 75% of the sample being classified as fast rotators and 25% as slow rotators, *Magneticum* successfully reproduces the observed strong trend of fast rotators being the dominant kinematical state of early-type galaxies. While the overall distribution is in quantitative agreement with observations, the simulation generates a non-negligible fraction of elongated SR with $\epsilon_{1/2} > 0.4$ that is not found in the observed sample. However, this seems to be a common feature of numerical simulations since this issue was already reported in previous studies (Bois et al. 2010; Naab et al. 2014; Penoyre et al. 2017).



Figure 18. Main Panel: The $\lambda_{\text{R}_{1/2}}$ - $\epsilon_{1/2}$ plane for the *Magneticum* ETGs subdivided into the kinematical groups at z = 2. The lines are the same as in Fig. 3.

While the observations are constrained to the fixed given projection on the sky, we investigate the $\lambda_{\text{R}_{1/2}}$ - $\epsilon_{1/2}$ plane in the more physical edge-on projection. We find that in the edge-on projection two distinct populations are present, clearly separated by an underpopulated region. The upper population is encompassed by an envelope combining the analytic isotropic line and the theoretical line obtained by assuming a linear relation between the anisotropy and the ellipticity. This is in substantial agreement with observations by Cappellari et al. (2007) who found the same limits for a sample of 25 regular rotating ETGs. Since the tensorvirial theorem does not impose those limits, this seems to be a fossil record of the formation history (Cappellari et al. 2007).

Studying the evolution of the $\lambda_{\mathrm{R}_{1/2}}\text{-}\epsilon_{1/2}$ plane with redshift, we find that the upper population is already in place at z = 2, while only a small fraction of ETGs occupy the slow rotating regime. With decreasing redshift, the statistical frequency of the SR increases gradually from 6% to 25%. We therefore conclude that the mechanism which forms slow rotators becomes more frequent at redshifts below z = 2. Accordingly, the frequency of fast rotators decrease from 94% at z = 2 to 75% at z = 0. This strongly indicates that dry (minor) merger events play an important role in the formation of slow rotators as discussed in previous studies (Naab et al. 2014; Moody et al. 2014). Investigating the evolution of $\lambda_{\mathrm{R}_{1/2}}$ for individual slow rotators we find that at least 30% of the slow rotators at z = 0 experience a rapid decrease of $\lambda_{\mathrm{R}_{1/2}}$ on a timescale of 0.5Gyr associated with a significant merger event.

Considering the fast rotators, we find the general trend of a spin down and a shift towards rounder shapes, consistent with the findings of Choi & Yi (2017). The described behaviour is also present when considering all galaxies that satisfy the resolution criteria (i.e. disks and galaxies undergoing a merger event). Clearly demonstrating that the recent formation of the slow rotating population and the spin-down are not due to a galaxy selection bias but in fact driven by physical processes in the low-z universe.

We show that, adopting the *b*-value from Teklu et al. (2015), the fundamental $M_* \cdot j_*$ plane is tightly connected to the $\lambda_{R_{1/2}} \cdot \epsilon_{1/2}$ plane. The continuous change of the *b*-value in the $\lambda_{R_{1/2}} \cdot \epsilon_{1/2}$ plane implies a simultaneous movement in both planes when a galaxy is influenced by external processes like merging, harassment or stripping. We suggest merging as a possible process that is able to cause the observed distributions in both planes.

We find a strong correlation between $\lambda_{R_{1/2}}$ and the *b*-value. This correlation and the associated scatter is already in place at z = 2. Using a toy model to estimate the functional form of the correlation, we show that the scatter in the relation is inherited from the scatter in the mass-size relation.

In order to investigate the connection between morphology and kinematics, we determine the Sérsic-index n as a proxy for the morphology of galaxies. We recover the general trend found in Krajnović et al. (2013) that ETGs with n < 1.5 are preferentially classified as fast rotators, while nis not adequate to disentangle fast and slow rotators.

Furthermore, we classify the Magneticum ETGs into five kinematical groups according to visual features in their velocity maps. Our sample includes all features stated in Krajnović et al. (2011), except for the 2σ galaxies which by definition are not considered in our study. 70% of our ETGs show a regular rotation pattern in excellent agreement with observational results. The class of non-rotating ETGs comprise 13% of the Magneticum ETGs, while 3% exhibit a distinct core and 1% show a kinematically distinct rotating core in the centre.

We find a non-neglectable number (17) of ETGs which show prolate rotation around the morphological major axis. Although this only represents a small fraction of 2% of our total sample their existence confirms that *Magneticum* comprises all the physics to form those observationally discovered objects. Interestingly, except for two galaxies, all PRs are classified as slow rotators. Investigating the mass dependence of the PR fraction we find a relatively flat distribution in the range 10.3 < log(M_*) < 12. In contrast, the fraction of the other groups decreases significantly leading to a rise of the fraction of prolate rotators. The increase in the statistical frequency of prolate rotators is in agreement with recent observational results of CALIFA and ATLAS^{3D}, as reported in Tsatsi et al. (2017) and theoretical results from the Illustris simulation (Ebrová & Lokas 2017).

ACKNOWLEDGMENTS

The Magneticum Pathfinder simulations were partially performed at the Leibniz-Rechenzentrum with CPU time assigned to the Project "pr86re". This work was supported by the DFG Cluster of Excellence "Origin and Structure of the Universe". KD is supported by the DFG Transregio TR33. We are especially grateful for the support by M. Petkova through the Computational Center for Particle and Astrophysics (C2PAP).

REFERENCES

- Arnold J. A., et al., 2014, ApJ, 791, 80
- Bacon R., et al., 2001, MNRAS, 326, 23
- Baldry I. K., et al., 2012, MNRAS, 421, 621
- Barnes J. E., 1988, ApJ, 331, 699
- Barnes J. E., 1992, ApJ, 393, 484
- Beck A. M., et al., 2015, preprint, (arXiv:1502.07358)
- Bender R., Doebereiner S., Moellenhoff C., 1988, A&AS, 74, 385
- Binney J., 2005, MNRAS, 363, 937
- Bois M., et al., 2010, MNRAS, 406, 2405
- Bois M., et al., 2011, MNRAS, 416, 1654
- Bournaud F., Jog C. J., Combes F., 2005, A&A, 437, 69
- Bournaud F., Jog C. J., Combes F., 2007, A&A, 476, 1179
- Cappellari M., 2016, ARAA, 54, 597
- Cappellari M., Copin Y., 2003, MNRAS, 342, 345
- Cappellari M., et al., 2007, MNRAS, 379, 418
- Cappellari M., et al., 2013a, MNRAS, 432, 1709
- Cappellari M., et al., 2013b, MNRAS, 432, 1862
- Choi H., Yi S., 2017, preprint, (arXiv:1702.00517)
- Ciotti L., Bertin G., 1999, A&A, 352, 447
- Cortese L., et al., 2016, MNRAS, 463, 170
- Cox T. J., Dutta S. N., Di Matteo T., Hernquist L., Hop-
- kins P. F., Robertson B., Springel V., 2006, ApJ, 650, 791 D'Eugenio F., Houghton R. C. W., Davies R. L., Dalla
- Bontà E., 2013, MNRAS, 429, 1258
- Davies R. L., et al., 2001, ApJL, 548, L33
- Davis T. A., et al., 2015, MNRAS, 449, 3503
- Dolag K., Vazza F., Brunetti G., Tormen G., 2005, MNRAS, 364, 753
- Dolag K., Borgani S., Murante G., Springel V., 2009, MN-RAS, 399, 497
- Donnert J., Dolag K., Brunetti G., Cassano R., 2013, MN-RAS, 429, 3564
- Ebrová I., Lokas E. L., 2015, ApJ, 813, 10
- Ebrová I., Lokas E. L., 2017, ApJ, 850, 144
- Emsellem E., et al., 2004, MNRAS, 352, 721
- Emsellem E., et al., 2007, MNRAS, 379, 401
- Emsellem E., et al., 2011, MNRAS, 414, 888
- Faber S. M., et al., 1997, AJ, 114, 1771
- Fabjan D., Borgani S., Tornatore L., Saro A., Murante G., Dolag K., 2010, MNRAS, 401, 1670
- Falcón-Barroso J., Lyubenova M., van de Ven G., 2015, in Cappellari M., Courteau S., eds, IAU Symposium Vol. 311, Galaxy Masses as Constraints of Formation Models. pp 78–81 (arXiv:1409.7786), doi:10.1017/S1743921315003439
- Fall S. M., 1983, in Athanassoula E., ed., IAU Symposium Vol. 100, Internal Kinematics and Dynamics of Galaxies. pp 391–398
- Fogarty L. M. R., et al., 2015, MNRAS, 454, 2050
- Forbes D. A., Sinpetru L., Savorgnan G., Romanowsky A. J., Usher C., Brodie J., 2017, MNRAS, 464, 4611
- Genel S., Fall S. M., Hernquist L., Vogelsberger M., Snyder G. F., Rodriguez-Gomez V., Sijacki D., Springel V., 2015, ApJL, 804, L40
- Greene J. E., et al., 2018, ApJ, 852, 36
- Hernquist L., 1990, ApJ, 356, 359
- Hernquist L., 1992, ApJ, 400, 460
- Hirschmann M., Dolag K., Saro A., Bachmann L., Borgani S., Burkert A., 2014, MNRAS, 442, 2304

Ho N., et al., 2012, ApJ, 758, 124

- Hoffman L., Cox T. J., Dutta S., Hernquist L., 2010, ApJ, 723, 818
- Houghton R. C. W., et al., 2013, MNRAS, 436, 19
- Hubble E. P., 1926, ApJ, 64, 321
- Huertas-Company M., et al., 2013, MNRAS, 428, 1715
- Jesseit R., Cappellari M., Naab T., Emsellem E., Burkert A., 2009, MNRAS, 397, 1202
- Kelvin L. S., et al., 2012, MNRAS, 421, 1007
- Komatsu E., et al., 2011, ApJS, 192, 18
- Kormendy J., Bender R., 1996, ApJL, 464, L119
- Krajnović D., et al., 2011, MNRAS, 414, 2923
- Krajnović D., et al., 2013, MNRAS, 432, 1768
- Lagos C. d. P., et al., 2017, preprint, (arXiv:1701.04407)
- Lagos C. d. P., et al., 2018, MNRAS, 473, 4956
- Lange R., et al., 2015, MNRAS, 447, 2603
- Lynden-Bell D., 1967, MNRAS, 136, 101
- McDermid R. M., et al., 2006, NewAR, 49, 521
- Mihos J. C., Hernquist L., 1996, ApJ, 464, 641
- Moody C. E., Romanowsky A. J., Cox T. J., Novak G. S., Primack J. R., 2014, MNRAS, 444, 1475
- Naab T., Burkert A., 2003, ApJ, 597, 893
- Naab T., et al., 2014, MNRAS, 444, 3357
- Obreschkow D., et al., 2015, ApJ, 815, 97
- Oser L., Ostriker J. P., Naab T., Johansson P. H., Burkert A., 2010, ApJ, 725, 2312
- Pedrosa S. E., Tissera P. B., 2015, A&A, 584, A43
- Penoyre Z., Moster B. P., Sijacki D., Genel S., 2017, MN-RAS, 468, 3883
- Querejeta M., et al., 2015, A&A, 579, L2
- Remus R.-S., Dolag K., 2016, in The Interplay between Local and Global Processes in Galaxies,.
- Remus R.-S., Dolag K., Naab T., Burkert A., Hirschmann M., Hoffmann T. L., Johansson P. H., 2017, MNRAS, 464, 3742
- Romanowsky A. J., Fall S. M., 2012, ApJS, 203, 17
- Sandage A., 1961, The Hubble atlas of galaxies
- Sandage A., 1975, Classification and Stellar Content of Galaxies Obtained from Direct Photography. the University of Chicago Press, p. 1
- Sarzi M., et al., 2006, MNRAS, 366, 1151
- Schulze F., Remus R.-S., Dolag K., 2017, Galaxies, 5, 41
- Scott N., Davies R. L., Houghton R. C. W., Cappellari M., Graham A. W., Pimbblet K. A., 2014, MNRAS, 441, 274
- Serra P., et al., 2014, MNRAS, 444, 3388
- Sérsic J. L., 1963, Boletin de la Asociacion Argentina de Astronomia La Plata Argentina, 6, 41
- Springel V., 2005, MNRAS, 364, 1105
- Springel V., Hernquist L., 2003, MNRAS, 339, 289
- Springel V., Yoshida N., White S. D. M., 2001a, NewA, 6, 79
- Springel V., White S. D. M., Tormen G., Kauffmann G., 2001b, MNRAS, 328, 726
- Springel V., Di Matteo T., Hernquist L., 2005, MNRAS, 361, 776
- Steinborn L. K., Dolag K., Hirschmann M., Prieto M. A., Remus R.-S., 2015, MNRAS, 448, 1504
- Teklu A. F., Remus R.-S., Dolag K., Beck A. M., Burkert A., Schmidt A. S., Schulze F., Steinborn L. K., 2015, ApJ, 812, 29
- Teklu A. F., Remus R.-S., Dolag K., 2016, in The Interplay between Local and Global Processes in Galaxies,.



Figure A1. Real anisotropy δ_{calc} calculated from the particle distribution using Eq. 11 against anisotropy obtained from the theoretical model δ_{model} using Eq. 13, for each ETG. The colours and symbols show the different kinematical groups, as described in the legend and Sec. 5.1.

- Thomas J., et al., 2009, MNRAS, 393, 641
- Toomre A., 1977, in Tinsley B. M., Larson D. Campbell R. B. G., eds, Evolution of Galaxies and Stellar Populations. p. 401
- Toomre A., Toomre J., 1972, ApJ, 178, 623
- Tornatore L., Borgani S., Dolag K., Matteucci F., 2007, MNRAS, 382, 1050
- Tsatsi A., Lyubenova M., van de Ven G., Chang J., Aguerri J. A. L., Falcón-Barroso J., Macciò A. V., 2017, A&A, 606, A62
- Veale M., Ma C.-P., Greene J. E., Thomas J., Blakeslee J. P., McConnell N., Walsh J. L., Ito J., 2017, MNRAS, 471, 1428
- Wu X., Gerhard O., Naab T., Oser L., Martinez-Valpuesta I., Hilz M., Churazov E., Lyskova N., 2014, MNRAS, 438, 2701
- de Vaucouleurs G., 1959, Handbuch der Physik, 53, 311
- de Vaucouleurs G., 1963, ApJS, 8, 31
- de Vaucouleurs G., de Vaucouleurs A., Corwin J. R., 1976, in Second reference catalogue of bright galaxies, Vol. 1976, p. Austin: University of Texas Press..
- de Vaucouleurs G., de Vaucouleurs A., Corwin Jr. H. G., Buta R. J., Paturel G., Fouqué P., 1991, Third Reference Catalogue of Bright Galaxies. Volume I: Explanations and references. Volume II: Data for galaxies between 0^h and 12^h . Volume III: Data for galaxies between 12^h and 24^h . de Zeeuw P. T., et al., 2002, MNRAS, 329, 513

APPENDIX A: APPENDIX A: THEORETICAL VERSUS REAL ANISOTROPY IN ETGS

In Sec. 3.3, we discussed the differences between the global anisotropy of an ETG calculated directly from the stellar particles, and the global anisotropy obtained from theoretical modelling using Eq. 13. Fig. A1 shows the correlation between both anisotropies found for each ETG in the *Magneticum* sample. While there is a clear correlation between these anisotropies, albeit with a large scatter, we find the correlation to be offset from the 1:1 correlation, with the theoretical values obtained from the model usually lower than the anisotropies calculated from the particles within the galaxies.

This offset and the scatter are neither coupled with the stellar mass of the ETGs, nor with their sizes, $\lambda_{R_{1/2}}$, the *b*-value, or any other global galaxy parameter tested in this study. The symbols in Fig. A1 demonstrate exemplary that there is also no correlation with the kinematical group features of the ETGs, and only the aforementioned effect, that slow-rotators tend to have slightly higher anisotropies than fast rotators, can be seen again in that the non-rotators generally occupy the area of higher anisotropies in both anisotropy-measurement methods than the regular rotators. The ETGs with prolate rotation occupy the same anisotropy-range as the non-rotators, showing no difference in the anisotropy behaviour. Only the ETGs with decoupled cores show less scatter in the correlation between the anisotropies from the different methods, and tend to have smaller anisotropies than the non-rotators and the prolate rotators in both anisotropy methods. While the former could be most likely due to low-number statistics, the latter indicates a systematic difference in the stellar orbital configurations for the decoupled core galaxies, most likely caused by the regular rotation patterns.

We also tested the anisotropy correlation for the galaxies where a kinematical classification was not possible, but we did not find any difference in their behaviour with respect to the correlation between the real and the calculated anisotropies. Therefore, a detailed investigation of the origin of the offset is needed, however this is beyond the scope of this work, as it is most likely due to the modelling process.

APPENDIX B: APPENDIX B: $\lambda_{R_{1/2}}-M_*$ RELATION

Observations revealed that very massive ETGs usually tend to be round and often slowly rotating ETGs, while the high $\lambda_{R_{1/2}}$ range is preferentially populated by less massive galaxies (e.g. Emsellem et al. 2011). One possible explanation for this behaviour is the formation of such massive systems through multiple minor merger (e.g. Naab et al. 2014; Moody et al. 2014). However, more recently, Ebrová & Lokas (2017) showed that many of the highest massive systems are prolate rotating, albeit the formation channel of these systems is still under debate, as discussed in Sec. 5.2.

The main panel (lowest) of Fig. B1 shows $\lambda_{R_{1/2}}$ against the stellar mass M_* for all ETGs (upper panel), with colours and symbols indicating the different kinematical groups introduced in Sec. 5.1. As can clearly be seen, the regular rotating galaxies dominate the lower mass regime, while non-rotators are present but not common. At the high mass end, the regular rotators become less dominant, while the fraction of galaxies with decoupled cores or prolate rotation increases (as can also be seen from the middle panel of Fig. B1, where the relative fractions of the different kine-



Figure B1. Lower (main) panel: $\lambda_{R_{1/2}}$ versus M_* for all ETGs in the *Magneticum* simulation. Regular Rotators (RR) are shown as red filled circles, Non-Rotators (NR) as blue filled triangles, Decoupled Core (DC) galaxies as cyan filled squares, Kinematically Distinct Core (KDC) galaxies as yellow filled diamonds, and Prolate Rotators (PR) as violet filled bowties. Upper panel: the number of galaxies of a certain kinematical group at different stellar mass bins normalised by the total number of galaxies within each group. The different kinematical groups are colourcoded according to the symbol colour in the lower panel. Middle panel: Relative percentage of each kinematical group at the different stellar mass bins, colour-coded as in the lower panel.

matical groups within each mass bin are shown). Especially the prolate rotators account for a significant fraction of the ETGs at the high mass end, in good agreement with recent observations by Tsatsi et al. (2017). The fraction of non-rotators, interestingly, is at about 10% regardless of the stellar mass, clearly showing that non-rotators occur on all mass bins. Nevertheless, at the highest mass bin above $M_* = 10^{11.75} M_{\odot}$, where the number of galaxies in the simulation is low, all ETGs are regular rotators with high values of $\lambda_{\rm R_{1/2}}$, in contrast to observations.

The upper panel of Fig. B1 shows the number of ETGs within a given kinematical group in each mass bin, normalised to the total number of all ETGs of that group. Both, regular rotators and non-rotators, are most frequent at the low-mass end, and become less frequent with increasing mass. On the contrary, distinct cores and kinematically distinct cores are features that become much more frequent at the higher mass end, albeit this could be a result of the resolution limitation of the simulation, as the sampling of particles at the low-mass end enhances the noise in the kinematical maps and therefore could suppress the signal of such features in the maps.

However, most interestingly, the relative number of prolate rotators is constant over all mass bins. Thus, while prolate rotators become more frequent at the high mass bin due to the decreasing number of galaxies with other kinematic features, their effective occurrence is independent of the mass, which is clearly in agreement with the observational fact that such prolate rotating galaxies are also seen at the dwarf-galaxy mass end (Ho et al. 2012; Ebrová & Lokas 2015). Therefore, a more detailed investigation of the formation mechanisms that lead to prolate rotation features, is needed in the future.

APPENDIX C: APPENDIX C: REDSHIFT EVOLUTION OF THE $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ DISTRIBUTION FOR ALL GALAXIES IN THE *MAGNETICUM* SIMULATION

As the classification criteria for galaxies into early- and latetype galaxies strongly vary between different surveys and especially redshifts, it is important to demonstrate that the redshifts trends found for galaxies in the $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ -plane in this paper are not biased by the classification. Therefore, Fig. C1 shows the same as Fig. 6, but for all galaxies in the *Magneticum* simulation above the resolution limit. Those galaxies that were classified as ETGs are shown as circles, while the galaxies that were rejected (see Sec. 2.2 for the classification criteria) are shown as diamonds, independent whether they are disk-like galaxies, merging systems, or rejected due to other reasons.

As can be seen, there is a clear tendency for all galaxies regardless of their morphology, to have higher $\lambda_{R_{1/2}}$ (and thus be more rotation dominated) and larger ellipticities at higher redshifts, clearly indicating that there exists a general spin-down for disk galaxies towards lower redshifts as well. This indicates that also disk galaxies might suffer from multiple minor merger event, without destroying the disklike structures (e.g., Karademir et al., in prep.). At z = 2, the slow-rotator regime remains nearly empty, and is populated subsequently at lower redshifts with the advent of more massive galaxies and an increasing fraction of ETGs. At all redshifts, the higher- $\lambda_{R_{1/2}}$ -regions are occupied by rejected galaxies (as expected since these galaxies include the disk-like galaxies).

This paper has been typeset from a TeX/ LATeX file prepared by the author.



Figure C1. $\lambda_{R_{1/2}}$ - $\epsilon_{1/2}$ -plane for all galaxies satisfying the resolution criteria, with ETGs shown as circles and the rejected galaxies shown as diamonds. Different colours and symbol-sizes indicate the mass of the galaxy. Upper left panel: z = 2; Upper right panel: z = 1; Lower left panel: z = 0.5; Lower right panel: z = 0. Lines are the same as in Fig. 3.