

The tidal trail of NGC 205?

A. W. McConnachie,^{1*} M. J. Irwin,¹ G. F. Lewis,² R. A. Ibata,³ S. C. Chapman,⁴
A. M. N. Ferguson⁵ and N. R. Tanvir⁶

¹*Institute of Astronomy, Madingley Road, Cambridge, CB3 0HA*

²*Institute of Astronomy, School of Physics, A29, University of Sydney, NSW 2006, Australia*

³*Observatoire de Strasbourg, 11, rue de l'Université, F-67000, Strasbourg, France*

⁴*California Institute of Technology, Pasadena, CA 91125, USA*

⁵*Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str. 1, Postfach 1317, D-85741 Garching, Germany*

⁶*Physical Sciences, University of Hertfordshire, Hatfield AL10 9AB*

Accepted 2004 May 27. Received 2004 May 19; in original form 2004 March 10

ABSTRACT

Using data taken as part of the Isaac Newton Telescope Wide Field Camera (INT WFC) survey of M31, we have identified an arc-like overdensity of blue, presumably metal-poor, red giant branch stars in the north-west quadrant of M31. This feature is $\sim 1^\circ$ (15 kpc) in extent and has a surface brightness of $\Sigma_{V'} \simeq 28.5 \pm 0.5$ mag arcsec⁻². The arc appears to emanate from the dwarf elliptical galaxy NGC 205, and the colour of its red giant branch is significantly different from the M31 disc population but closely resembles that of NGC 205. Further, using data taken with the Deep Imaging Multi-Object Spectrograph (DEIMOS) on Keck II, we identify the radial velocity signature of this arc. Its velocity dispersion is measured to be $\simeq 10$ km s⁻¹, similar to that of the central regions of NGC 205 and typical of stellar streams. Based upon the spatial coincidence of these objects, the surface brightness, the velocity dispersions and the similarity in colour of the red giant branches, we postulate that the arc is part of a stellar stream, the progenitor of which is NGC 205.

Key words: galaxies: dwarf – galaxies: general – galaxies: interactions – Local Group.

1 INTRODUCTION

Over the course of the last decade there has been a lot of renewed interest in the Local Group. This has in part been due to several discoveries of nearby galaxies by various groups (Ibata, Gilmore & Irwin 1994; Whiting, Irwin & Hau 1997; Armandroff, Davies & Jacoby 1998; Armandroff, Jacoby & Davies 1999; Karachentsev & Karachentseva 1999; Whiting, Hau & Irwin 1999) that have been made possible because of significant advances in instrumentation. At the same time as these observational discoveries, hierarchical formation scenarios of structure formation, and in particular cold dark matter (CDM), have reached a sufficiently detailed level for the Local Group to begin acting as a laboratory to test and constrain these theoretical models. Here, larger galaxies such as M31 or the Milky Way are postulated to form via the merging of smaller bodies (White & Rees 1978; Searle & Zinn 1978).

The Sagittarius dwarf galaxy, discovered by Ibata et al. (1994), was subsequently shown to have a substantial stellar stream associated with it (e.g. Mateo, Olszewski & Morrison 1998; Majewski et al. 1999; Dohm-Palmer et al. 2001; Ibata et al. 2001a), a direct result of its interaction with the gravitational potential of the Milky

Way. More recent work has shown that up to 75 per cent of halo M-giants actually belong to the Sagittarius dwarf (Majewski et al. 2003), demonstrating the significant role that this accretion event has had on the make-up of our Galaxy. Tidal streams naturally fit in to hierarchical models of structure formation, and evidence for tidal disruption has been found for many of the Galactic dSph satellites (Irwin & Hatzidimitriou 1995) as well as for many galaxies located at higher redshift (e.g. Chapman et al. 2003).

The Andromeda galaxy, M31, provides the only example of a giant spiral galaxy, other than the Milky Way, in which individual stars can be easily resolved using current generations of ground-based technology. As such, we have conducted a large photometric survey of M31 using the Isaac Newton Telescope Wide Field Camera (INT WFC) and the CFH12K camera on the Canada–France–Hawaii Telescope (Ibata et al. 2001b; Ferguson et al. 2002; McConnachie et al. 2003; Irwin et al., in preparation; hereafter Papers I, II, III and IV, respectively), complete over total area of 40 deg², ranging out to 80 kpc from the centre of this galaxy (see the figures in Paper IV). Study of the structure of Andromeda is in many ways easier than for our own Galaxy, as our location external to M31 allows us to obtain a global view without the projection problems that plague studies of the structure of the Milky Way.

Initial results of our survey were spectacular, revealing a considerable amount of unexpected substructure. Most significant of

*E-mail: alan@ast.cam.ac.uk

all these was the discovery of a giant tidal stream, extending out to some 40 kpc in projection from the centre of M31 (Paper I). A subsequent, deeper survey using the CFH12K camera allowed the three-dimensional position of the stream to be calculated, and showed that this feature is over 140 kpc in extent (Paper III). A follow-up kinematic survey using the Deep Imaging Multi-Object Spectrograph (DEIMOS) on the KECK II telescope has provided radial velocities for over 800 stars in 13 different fields located at various positions in the halo and disc of M31, concentrating on areas of known substructure. Initial results of this survey are detailed in Ibata et al. (2004).

In this Letter, we report on the identification of what appears to be another substantial tidal stream in M31, distinct in both colour and kinematics from the M31 stellar stream that has already been reported in Papers I and III. The projected position of this feature, the colour of its red giant branch (RGB) and its velocity dispersion all suggest that it is tidal debris from NGC 205, the bright dwarf elliptical companion of M31. This satellite has long been known to be tidally perturbed by M31 (e.g. Hodge 1973; Kent 1987; Bender, Paquet & Nieto 1991; Choi, Guhathakurta & Johnston 2002; Demers, Battinelli & Letarte 2003) and the discovery of an associated stream will allow for a better understanding of its evolution. Additionally, if this interpretation is correct, then it is yet more evidence of the ubiquitous nature of streams in galactic haloes which, taken with the other substructure present in M31, reveals the complex formation history of this galaxy.

2 PHOTOMETRY

Details of the search strategy, observations and data reduction of our INT WFC survey of M31 can be found in Paper II, and the completed

maps showing the global two-dimensional spatial distribution of various stellar populations in M31 can be found in Paper IV. These stellar populations were defined by isolating different loci in the colour–magnitude diagrams (CMDs) created from our Johnson V (V') and Gunn i (i') photometry (Paper II). In Paper IV, where we show the distribution of ‘blue’ RGB stars, a prominence of stars is visible centred at approximately $(-0.6, 1.1)$. It is this feature that is analysed here in more depth.

The left-hand panel of Fig. 1 shows a map of blue RGB stars similar to that shown in Paper IV, defined using a sample limited by $20.5 < i' < 22.5$ mag, $24.85 - 2.85 (V' - i') < i' < 26.85 - 2.85 (V' - i')$. An arc-like overdensity is visible in this figure, with a projected length of $\sim 1^\circ$ (15 kpc), but significantly curved. It extends to the north of NGC 205 but then bends back to the east into the disc of M31. At these galactocentric radii, we are unable to follow it further due to the dominant M31 disc population. The morphology of this feature is strongly suggestive of a stellar stream. However, the quantity of substructure in our survey makes it difficult to categorize these features based on morphology alone; for example, it could conceivably also be disrupted M31 outer disc. We estimate the surface brightness of this stellar arc in the V' band to be $\Sigma_{V'} \sim 28.5 \pm 0.5$ mag arcsec $^{-2}$. The Sagittarius stream has a surface brightness of ~ 30 mag arcsec $^{-2}$, although there is substantial variation with azimuthal angle (Mateo et al. 1998), and the M31 stream reported in Paper I has an estimated surface brightness of $\sim 30 \pm 0.5$ mag arcsec $^{-2}$. The arc is therefore typically somewhat brighter than these other remnants, although still well within the expected range for stellar streams.

Fig. 2 shows a CMD for the arc (left-hand panel), NGC 205 (middle panel) and a typical M31 disc field (right-hand panel). Overlaid are four well-studied globular cluster fiducial sequences of




Figure 1. Left-hand panel: the spatial distribution of ‘bluer’ RGB stars, defined by the colour cut described in the text. This cut is designed to highlight the arc-like density enhancement centred at approximately $(-0.6, 1.1)$. The dwarf elliptical galaxy NGC 205 is located at $(-0.6, 0.4)$. The metal-poor dwarf spheroidal galaxy I stands out to the south of the plot, in addition to other complex substructure around the outer south-west disc. Right-hand panel: cartoon showing the location of our outer disc DEIMOS fields with respect to M32, NGC 205 and the stellar arc. In both panels, the 2° -radius ellipse marks the outer boundary of the optical disc of M31, whereas the outer ellipse has a semimajor axis $\simeq 55$ kpc and flattening 0.6. This corresponds to the original limit of the INT survey.




Figure 2. Hess diagrams showing the CMDs for the stellar arc (left-hand panel), NGC 205 (middle panel) and a typical M31 outer disc field (right-hand panel). Overlaid on these are four well-studied globular cluster sequences of different metallicities. From left to right, these are NGC 6397 ($[\text{Fe}/\text{H}] = -1.9$), NGC 1851 ($[\text{Fe}/\text{H}] = -1.3$), 47 Tuc ($[\text{Fe}/\text{H}] = -0.7$) and NGC 6553 ($[\text{Fe}/\text{H}] = -0.2$). Both the arc and NGC 205 have an enhanced bluer RGB component compared with the M31 field, located at a similar galactocentric radius. The enhanced bluer RGB in the left-hand panel appears to be well represented by a metallicity of $[\text{Fe}/\text{H}] \sim -0.9$, in good agreement with that derived for NGC 205 (Mould et al. 1984; Mateo 1998).

different metallicities. From left to right, these are NGC 6397 ($[\text{Fe}/\text{H}] = -1.9$), NGC 1851 ($[\text{Fe}/\text{H}] = -1.3$), 47 Tuc ($[\text{Fe}/\text{H}] = -0.7$) and NGC 6553 ($[\text{Fe}/\text{H}] = -0.2$) (Da Costa & Armandroff 1990; Sagar et al. 1999). Both NGC 205 and the arc show an enhanced bluer RGB component relative to the M31 field located at a similar galactocentric radius. Differential reddening is unlikely to account for such a feature. If the stellar arc is somehow related to the outer disc of M31, it is not clear how the colours of the RGBs could be so markedly different. Comparison of the arc CMD with the globular cluster tracks shows that it is well described by a mean metallicity of $[\text{Fe}/\text{H}] \sim -0.9$. Mould, Kristian & Da Costa (1984) measured a metallicity for NGC 205, as implied by the colour of its RGB, of $[\text{Fe}/\text{H}] \sim -0.9 \pm 0.2$ (see also Mateo 1998). Our INT photometry therefore suggests a link between this feature and NGC 205.

3 KINEMATICS

We have used the DEIMOS instrument on Keck II to measure radial velocities of $\simeq 80$ stars per field in a total of 13 fields in various locations across the extent of our photometric survey, selected to lie upon various interesting photometric features. The locations of the fields in the outer disc are shown in the right-hand panel of Fig. 1. Details of the observations and reduction process, and early results from this survey, can be found in Ibata et al. (2004). The typical uncertainties in the radial velocities that we measure are $5\text{--}10 \text{ km s}^{-1}$. One of our fields (W91) is centred at an RA and Dec of $0^{\text{h}}41^{\text{m}}5^{\text{s}}.63$, $42^{\circ}34'25''.1$. This samples the arc at the point at which it turns to go back into the disc, as indicated in Fig. 1. A histogram of the stars in this field with heliocentric radial velocities in the range $-500 \leq v_{\odot} \leq 80 \text{ km s}^{-1}$, with a Tonry–Davis coefficient (Tonry & Davis 1979) greater than 7.5 and an average continuum flux in the Ca II triplet region > 50 counts per pixel (0.31 \AA) is shown in the right-hand panel of Fig. 3. The left-hand panel shows a comparison field (W80) located at similar galactocentric radius on the other side of the minor axis (Fig. 1). For reference, the systemic heliocentric

radial velocity of M31 of $v_{\odot}(\text{M31}) \simeq -300 \text{ km s}^{-1}$ (Mateo 1998) is indicated by the dashed line.

In both panels of Fig. 3, we can identify various stellar components. The stars with velocities $\geq -100 \text{ km s}^{-1}$ are likely foreground stars in the halo and disc of the Milky Way. The stars with $v_{\odot} < -300 \text{ km s}^{-1}$ are attributable to the halo or bulge of M31, and presumably have an equivalent number with $v_{\odot} > -300 \text{ km s}^{-1}$. However, we attribute the peak at -340 km s^{-1} in W80 and the peak at -215 km s^{-1} in W91 to the M31 disc. As both these fields are located approximately symmetrically on either side of the minor axis of M31, we would then expect the disc components in each to lie approximately symmetrically on either side of the M31 systemic velocity, as we observe. The solid curves in each panel of Fig. 3 are Gaussian fits to the disc component, each with a velocity dispersion of $\sim 20 \text{ km s}^{-1}$ (uncorrected for measuring errors of typically $5\text{--}10 \text{ km s}^{-1}$). This value is consistent with our knowledge of the M31 disc. In W91 there is an additional velocity component centred on -160 km s^{-1} with a corrected velocity dispersion of $\simeq 10 \text{ km s}^{-1}$ (dashed curve in Fig. 3). As this feature is unique to this field, we attribute it to the photometrically identified stellar arc.

NGC 205 has a measured radial velocity of -240 km s^{-1} . The KECK radial velocity field that we have of the arc is located nearly 1° away from NGC 205, and shows a component centred on -160 km s^{-1} with a corrected velocity dispersion of $\simeq 10 \text{ km s}^{-1}$. We would expect a substantial velocity difference between NGC 205 and the arc in this field, whether the two are related or not, as they are separated by $\gtrsim 15 \text{ kpc}$, depending upon the inclination of the arc to the line of sight. Therefore, without detailed modelling or stellar velocities in the region between our field and NGC 205, it is impossible to draw any firm conclusions regarding their association based upon only the magnitude of their radial velocities. On the other hand, the velocity dispersion of the arc is typical of stellar streams and is comparable to that which has been measured for the central regions of NGC 205 ($\sim 14 \text{ km s}^{-1}$; Carter & Sadler 1990; Peterson & Caldwell 1993), which supports a link between the two. We note that the velocity dispersion of NGC 205 is known to show




Figure 3. The heliocentric radial velocity distribution in W80 (left-hand panel) and W91 (right-hand panel). The location of these fields is marked on Fig. 1. The vertical dashed line in both panels represents the systemic velocity of M31. The solid curves are a Gaussian fit to the M31 disc population, with a dispersion of $\sim 20 \text{ km s}^{-1}$. W91 has an additional component present, centred on -160 km s^{-1} with a corrected velocity dispersion of $\simeq 10 \text{ km s}^{-1}$, which we attribute to the stellar arc (dashed curve).

a substantial increase with radius, with a velocity dispersion outside of the nucleus of $40\text{--}50 \text{ km s}^{-1}$ (Bender et al. 1991). This is indicative of tidal disruption and so therefore the presence of a tidal stream would not be surprising.

4 DISCUSSION

The quantity of substructure in M31 makes it difficult to decide upon the origin of any single feature (Fig. 1; see also Papers II and IV). Several interpretations as to the origin of this stellar arc are possible, and here we consider each in turn.

One possibility is that this feature is a continuation of the stream reported in Papers I and III. This seems unlikely, however, as the two structures have significantly different colours; the stellar arc is bluer, suggesting a mean $[\text{Fe}/\text{H}] \sim -0.9$, whereas the previously identified stream is redder, with a mean $[\text{Fe}/\text{H}] \sim -0.5$ (Paper III). Furthermore, the systemic kinematic signatures are quite distinct (Ibata et al. 2004), and so we conclude that the stellar arc is unrelated to the previously identified stellar stream.

A second possibility is that the arc is actually disrupted M31 disc. Although this interpretation is certainly plausible considering the amount of disruption evident in M31, we consider it unlikely for two reasons. First, as Fig. 2 demonstrates, this feature has a substantially different RGB colour from the outer disc, with a significantly enhanced bluer component. We would expect the CMD of any disrupted disc feature to closely resemble the CMD of the undisturbed disc, although this is not the case here. While not disproving a link between the two, it is not clear how this difference can be easily reconciled. Additionally, the velocity dispersion of the arc ($\simeq 10 \text{ km s}^{-1}$) is extremely small for a disc component; we measure a M31 disc radial velocity dispersion of $\sim 20 \text{ km s}^{-1}$ from Fig. 3 and we note that the velocity ellipsoid in the Milky Way disc at the Solar neighbourhood is $\sim 39:23:20 \text{ km s}^{-1}$ (Dehnen & Binney 1998). Again, this suggests that this feature is unrelated to the M31 disc.

The third possibility is that this is a stellar stream. Its morphological appearance, surface brightness and small velocity dispersion are all consistent with this interpretation. Additionally,

the strong spatial coincidence of the ‘base’ of the arc with NGC 205, the close similarity of the RGB colours of these objects and the similarity of the arc velocity dispersion to that for the central region of NGC 205 all allude to this object as the progenitor. As this interpretation is fully consistent with all available photometric and kinematic information, it is our favoured hypothesis. To confirm this, radial velocities in several fields positioned along the arc are required. If correct, then a velocity gradient connecting NGC 205 with the arc in our field will be present. This information will also be vital, should the connection be verified, for dynamical modelling of this intriguing system.

Taking this as our working hypothesis, by counting the excess stars in the on-stream regions compared with off-stream regions, we estimate a stellar luminosity density of $\sim 1 L_{\odot} \text{ arcsec}^{-2}$. If we assume a value for the mass–light ratio in NGC 205 of ~ 9 (Held, Mould & de Zeeuw 1990; Carter & Sadler 1990; Bender et al. 1991; Peterson & Caldwell 1993), then the mass density in this stream is of order $9 M_{\odot} \text{ arcsec}^{-2}$. We estimate the area of the arc to be $\sim 0.2 \text{ deg}^2$, implying a total mass in the arc of $\sim 1.8 \times 10^7 M_{\odot}$, not accounting for projection effects. The mass of NGC 205 is of order $7.4 \times 10^8 M_{\odot}$ (Mateo 1998), and so the visible part of the stream contains ~ 2.5 per cent of the mass of NGC 205.

There has been other evidence presented in the literature that suggests NGC 205 is undergoing tidal disruption. Hodge (1973) was the first to undertake detailed photometry of this object and showed that its outer isophotes are twisted, presumably as a result of tidal interactions with M31. Subsequent photometry was later conducted by Kent (1987), and he found reasonable agreement with Hodge (1973). Most recently, Choi et al. (2002) have found strong evidence for tidal distortion in the isophotes of NGC 205. Fig. 9 of Paper II is an isopleth map of M31 from an APM scan of a 75-min-exposure Palomar Schmidt IIIaJ plate taken by Sydney van den Bergh in 1970, and it shows this effect very clearly. The increase in its velocity dispersion as a function of radius, reported by Bender et al. (1991), also suggests that there has been some event which has disturbed the stars in the outer parts of NGC 205 and caused them to have an unusually large velocity dispersion. Recently, Demers et al. (2003) has shown that the number

of carbon stars in this galaxy is low for its luminosity, and that there are very few carbon stars located at greater than 10 arcmin from the centre, which suggests that NGC 205 has been tidally stripped by M31. Our potential discovery of a tidal stellar stream is therefore unsurprising and is readily testable with further kinematic data. If the stream hypothesis is correct, then the discovery of this stream will greatly aid in the understanding of the evolution of NGC 205 and provide yet further evidence of the prevalence of stellar streams in galactic haloes and of the *ongoing* role of accretion events in galactic evolution.

ACKNOWLEDGMENTS

This paper is based in part on observations made with the Isaac Newton Telescope on the Island of La Palma by the Isaac Newton Group in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. Some of the data presented herein were obtained at the W.M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W.M. Keck Foundation.

AWM would like to thank the University of Sydney Physics Department for their hospitality during a recent collaborative visit. The research of AMNF has been supported by a Marie Curie Fellowship of the European Community under contract number HPMF-CT-2002-01758. GFL acknowledges support from the Australian Research Council under Discovery Project DPO343508.

REFERENCES

- Armandroff T. E., Davies J. E., Jacoby G. H., 1998, *AJ*, 116, 2287
 Armandroff T. E., Jacoby G. H., Davies J. E., 1999, *AJ*, 118, 1220
 Bender R., Paquet A., Nieto J.-L., 1991, *A&A*, 246, 349
 Carter D., Sadler E. M., 1990, *MNRAS*, 245, 12p
 Chapman S. C., Windhorst R., Odewahn S., Yan H., Conselice C., 2003, *ApJ*, 599, 92
 Choi P. I., Guhathakurta P., Johnston K. V., 2002, *AJ*, 124, 310
 Da Costa G. S., Armandroff T. E., 1990, *AJ*, 100, 162
 Dehnen W., Binney J. J., 1998, *MNRAS*, 298, 387
 Demers S., Battinelli P., Letarte B., 2003, *AJ*, 125, 3037
 Dohm-Palmer R. C. et al., 2001, *ApJ*, 555, L37
 Ferguson A. M. N., Irwin M. J., Ibata R. A., Lewis G. F., Tanvir N. R., 2002, *AJ*, 124, 1452 (Paper II)
 Held E. V., Mould J. R., de Zeeuw P. T., 1990, *AJ*, 100, 415
 Hodge P. W., 1973, *ApJ*, 182, 671
 Ibata R. A., Gilmore G., Irwin M. J., 1994, *Nat*, 370, 194
 Ibata R., Irwin M., Lewis G., Ferguson A. M. N., Tanvir N., 2001a, *Nat*, 412, 49 (Paper I)
 Ibata R., Irwin M., Lewis G. F., Stolte A., 2001b, *ApJ*, 547, L133
 Ibata R., Chapman S., Ferguson A. M. N., Irwin M. J., Lewis G. F., McConnachie A. W., Tanvir N., 2004, *MNRAS*, 351, 117
 Irwin M., Hatzidimitriou D., 1995, *MNRAS*, 277, 1354
 Karachentsev I. D., Karachentseva V. E., 1999, *A&A*, 341, 355
 Kent S. M., 1987, *AJ*, 94, 306
 McConnachie A. W., Irwin M. J., Ibata R. A., Ferguson A. M. N., Lewis G. F., Tanvir N., 2003, *MNRAS*, 343, 1335 (Paper III)
 Majewski S. R., Siegel M. H., Kunkel W. E., Reid I. N., Johnston K. V., Thompson I. B., Landolt A. U., Palma C., 1999, *AJ*, 118, 1709
 Majewski S. R., Skrutskie M. F., Weinberg M. D., Ostheimer J. C., 2003, *ApJ*, 599, 1082
 Mateo M. L., 1998, *ARA&A*, 36, 435
 Mateo M., Olszewski E. W., Morrison H. L., 1998, *ApJ*, 508, L55
 Mould J., Kristian J., Da Costa G. S., 1984, *ApJ*, 278, 575
 Peterson R. C., Caldwell N., 1993, *AJ*, 105, 1411
 Sagar R., Subramaniam A., Richtler T., Grebel E. K., 1999, *A&AS*, 135, 391
 Searle L., Zinn R., 1978, *ApJ*, 225, 357
 Tonry J., Davis M., 1979, *AJ*, 84, 1511
 White S. D. M., Rees M. J., 1978, *MNRAS*, 183, 341
 Whiting A. B., Irwin M. J., Hau G. K. T., 1997, *AJ*, 114, 996
 Whiting A. B., Hau G. K. T., Irwin M., 1999, *AJ*, 118, 2767

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.