

# Global fitting of globular cluster age indicators <sup>★</sup>

F. Meissner<sup>1</sup> and A. Weiss<sup>1</sup>

Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str. 1, 85748 Garching, Federal Republic of Germany  
e-mail: (meissner,weiss)@mpa-garching.mpg.de

## ABSTRACT

*Context.* Methods for age determinations of globular clusters are still in need of improvement.

*Aims.* It is attempted to obtain a more objective method of age determinations based on cluster diagrams, avoiding the introduction of biases due to the preference of one single age indicator.

*Methods.* We compute new stellar evolutionary tracks and derive the dependence of age indicating points along the tracks and isochrone – such as the turn-off or bump location – as function of age and metallicity. The same critical points are identified in the colour-magnitude-diagrams of globular clusters from a homogeneous database. Several age indicators are then fitted simultaneously, and the overall best-fitting isochrone is selected to determine the cluster age. We also determine the goodness-of-fit for different sets of indicators to estimate the confidence level of our results.

*Results.* We find that our isochrones provide no acceptable fit for all age indicators. In particular, the location of the bump and the brightness of the tip of the red giant branch are problematic. On the other hand, the turn-off region is very well reproduced and restricting the method to indicators depending on it results in trustworthy ages. Using an alternative set of isochrones improves the situation, but neither leads to an acceptable global fit.

*Conclusions.* We conclude that evolutionary tracks of low-mass metal-poor stars are far from reproducing all aspects of globular cluster colour-magnitude-diagrams and that the determination of cluster ages still depends on the favourite method or indicator chosen.

**Key words.** stars: evolution – globular clusters: general – globular clusters: ages

---

Send offprint requests to: A. Weiss

\* submitted to Astron. Astrophys.

## 1. Introduction

The determination of globular cluster (GC) ages rests on the fact that colour-magnitude-diagrams (CMD) of single-age single-composition stellar populations exhibit specific time-dependent features. Most importantly, this is the location of the turn-off (TO), which – together with the cluster’s distance – serves as the most straightforward and widely used age indicator. However, there are other parts of the CMD, which change their colour or brightness with age, too. Since the sensitivity to time is different for the various parts of the cluster CMD, it is possible to either use various indicators independently, or to use the differences in colour and brightness between pairs of them; these latter methods have the advantage of being distance independent. Finally, one may attempt to fit the overall shape of the cluster diagram with one isochrone, which is the preferred method of Vandenberg (2000).

The decision, which age indicator is used, is the choice of the investigator and may depend on the observational data (e.g. CMDs not deep enough prevent the use of the TO), the isochrones (with or without the horizontal branch), or simply tradition. In the epoch of rather accurate *Hipparcos* distances, the traditional TO brightness has experienced new interest (Gratton et al. 1997; Carretta et al. 2000). Salaris & Weiss (1997, 2002) prefer the brightness difference between horizontal branch (HB) and TO for absolute, and the colour difference between TO and red giant branch (RGB) for relative ages, while Rosenberg et al. (1999) use the former method for an almost empirical relative age scale. Rarely, several age indicators are used simultaneously to check for consistency. Therefore, the choice of the age determination method is subject to a bias.

The aim of this paper is to investigate whether the simultaneous use of several age indicators, which implies a higher degree of objectivity, is possible and whether it leads to overall acceptable isochrone fits. We restrict ourselves to brightness differences, thereby avoiding the problem of distance accuracy. We do not intend to present new or improved age determinations, but rather concentrate on developing a new and more reliable method. The basic idea is to define a selection of crucial points along the cluster sequences and the corresponding ones along theoretical isochrones. These are the TO, and two points close to it, but cooler; the bump and tip of the RGB, and the color of the lower RGB. HB and RGB tip are almost age independent and serve as reference points; they also provide a means of predicting the distance to the cluster. This can be compared to independent distance measurements to check for consistency and for graphical overlay of isochrone and cluster ridge line.

Various brightness and colour differences are defined according to their use in the literature as age indicators. They are derived directly from the photometric data; theoretical values are determined from our isochrones, which we computed specifically for this project, and their age dependence fitted by simple analytical formulae. The deviation between theoretical and observed value as function of age enters a goodness-of-fit test (a  $\chi^2$ -method). The overall minimum of this function determines the best age, and its value the confidence level. The basic result is that using all indicators results in flat minima with low probability, while the restriction to subsets of indicators (equivalent to putting zero weight to the others) allow quite accurate age determinations. The reason for the failure of the complete set of age indicators lies in the inability of the isochrones to fit not just the overall shape, but the individual features of a cluster CMD. More details about our work can be found in Meissner (2005) on which this paper is based.


The age indicators will be introduced and discussed in Sect. 2 with references to earlier work using them. The observational data are presented in Sect. 3 along with the details of determining ridge line and the various features of interest. Then (Sect. 4) follows a short description of our calculations and set of stellar tracks. Section 5 contains the presentation of results, before the discussion in Sect. 6 closes the paper.

## 2. Age indicators from cluster CMD

### 2.1. Indicator points

*Turn-off region:* The classical age indicator (Renzini & Fusi Pecci 1988) in cluster CMDs is the turn-off (indicated in Fig. 1 by “TO”), whose position moves to the lower right (cooler and fainter) with age. Its colour can be determined easily and accurately, if the reddening is small or well-determined, while the brightness is more difficult to fix due to the long and almost vertical extension of the ridge-line (or, equivalently, the isochrone) which can reach up to 0.1 mag, corresponding to an age uncertainty of about 1 Gyr. In this paper we will always indicate TO quantities by the corresponding index.

To avoid the problem of determining  $V_{TO}$  accurately, two alternative indicator points have been introduced, which are 0.05 mag redder than the TO, but lie on less steep portions of the isochrone: the point  $TO^b$  on the post-TO part and  $TO^f$  on the main-sequence. The first alternative was introduced by Chaboyer et al. (1996); it has the same age sensitivity as the TO itself, but its determination is claimed to be a factor of 2 better. The second one was proposed by Buonanno et al. (1998) and is claimed to be additionally less affected by



**Fig. 1.** A sample isochrone ( $\log Z/Z_{\odot} = -3$ ;  $t = 12$  Gyr) in the observable plane and points along it used for the age indicators.

uncertainties such as the convection theory. While this conjecture is plausible on the basis that pre-TO stars are less evolved, it was never investigated in detail.

*Red giant branch:* The brightness of the red giant branch tip depends weakly on composition, but hardly at all on age. Therefore it is used as a standard candle to determine distances to galaxies with resolved populations (see Salaris & Cassisi 1997, 1998 for a discussion).  $V_{\text{tip}}$  is therefore a well-suited reference point, from which brightness difference to time-dependent points can be defined.

On the lower red-giant branch the colour of a point or region is used for colour-based age indicators (also called “horizontal” indicators in contrast to the brightness-based “vertical” ones). We chose the colour of a point 2.5 mag brighter than the TO, and call its color for example  $(B - V)_{\text{RGB}}$ ; Salaris & Weiss (1997) used the mean colour for a range of about 0.5-1.0 mag starting from this point. The uncertainty of  $V_{\text{TO}}$  is not very crucial since the RGB colour changes hardly over 0.1 mag.


The most age-sensitive feature along the RGB is the so-called bump, where evolutionary tracks display a vertical loop, which observationally leads to an enhanced number of stars in a limited magnitude range (see Salaris et al. 2002 for more details). The bump is now routinely found in high-quality cluster photometry, both in the optical (Riello et al. 2003) and near infrared (Valenti et al. 2004). Its location both in colour and brightness is mass-, and therefore age-dependent. It also depends on composition.

As the bump has to be detected from the luminosity function (LF) of a cluster, we show in Fig. 2 the simulated LFs of isochrones with  $\log Z = -3$  for ages of 6 and 12 Gyr. For the initial mass function (IMF) needed to produce the LF the standard Salpeter law (Salpeter 1955) was assumed.


*Horizontal branch:* The brightness of the zero-age horizontal branch (ZAHB) is only very weakly dependent on age and therefore  $V_{\text{ZAHB}}$  is a widely used for vertical age indicators (Vandenbergh et al. 1990; Salaris & Weiss 1997). Theoretically it is easily determined, but observationally one estimates it from the lower envelope of the horizontal branch. From  $V_{\text{ZAHB}}$  one can thus derive the distance to the cluster; in general, the agreement with main-sequence fitting distances is quite good (Salaris & Weiss 1998). The colour of the HB red end is another possible age-dependent feature, but due to our lack of understanding of HB morphology we chose not to use it.

## 2.2. Age indicators

$\Delta V_{\text{TO}}^{\text{ZAHB}}$ : The difference between TO and ZAHB,  $\Delta V_{\text{TO}}^{\text{ZAHB}} = V_{\text{TO}} - V_{\text{ZAHB}}$ , is the most widely used vertical age indicator (see De Angeli et al. 2005 for the most recent application). If  $V_{\text{ZAHB}}$  is taken at the same colour as the TO, it is largely independent of uncertainties in the colour transformations or reddening, and depends more on the relative accuracy of bolometric corrections, as was discussed by Salaris & Weiss (1998). In contrast to that paper and to Salaris & Weiss (2002) we use  $\Delta V_{\text{TO}}^{\text{ZAHB}}$  also for those cluster, which do not have stars on the HB at the TO colour. This requires a less accurate approach: we chose to select the ZAHB level at the closest colour available in the observed CMD. As long as this point is on the horizontal part of the ZAHB,



**Fig. 2.** Luminosity function isochrones for  $\log Z/Z_{\odot} = -3$  and ages of 6 (solid line) and 12 Gyr (dashed). Along the LF main-sequence (MS), sub-giant branch (SGB), red giant branch (RGB), tip of the RGB and the position of the RGB bump are indicated.




**Fig. 3.**  $\Delta V_{TO}^{ZAHB}$  (upper lines) and  $V_{ZAHB}$  (lower lines) for  $Z = 2 \cdot 10^{-4}$  (solid) and  $10^{-2}$  (dashed) as function of time from our theoretical isochrones.

this appears to be rather safe, but for an extremely blue morphology it will be less so. Fig. 3 shows  $\Delta V_{TO}^{ZAHB}$  as function of age as well as  $V_{ZAHB}$  for two metallicities, as obtained from the theoretical isochrones. It is obvious that  $V_{ZAHB}$  is nearly constant over an age range of 8-15 Gyr.

$\Delta V_{TO}^{ZAHB}$  and  $\Delta V_{TO}^{ZAHB}$ : These are the two alternative age indicators equivalent to  $\Delta V_{TO}^{ZAHB}$ , but without the uncertainty connected with the determination of  $V_{TO}$ . All other comments made in the previous paragraph hold. The age and metallicity dependence is very similar to that of  $\Delta V_{TO}^{ZAHB}$ .

$\Delta V_{TO}^{Bump}$ : In case of a HB with a morphology rendering the previous age indicators inaccurate, one can resort to  $\Delta V_{TO}^{Bump} = V_{TO} - V_{Bump}$  instead. Its age sensitivity is slightly lower than that of  $\Delta V_{TO}^{ZAHB}$  for low ages, but almost comparable at the interesting age range around 10 Gyr. The problems with this indicator lie in the need for photometry with a sufficient number of stars to identify the bump and theoretically, as we will see, that it appears to be too bright systematically by a few tenths of a magnitude as is also found in some observational papers (e.g. Zoccali & Piotto 2000).

$\Delta V_{TO}^{tip}$ :  $\Delta V_{TO}^{tip} = V_{TO} - V_{tip}$  is again useful in cases of an insufficiently populated or observed HB. It is particularly sensitive to age for lower metallicities, changing by more than 0.5 mag between 8 and 14 Gyr.



**Fig. 4.**  $\Delta(B - V)_{\text{TO}}^{\text{RGB}}$  (solid lines),  $(B - V)_{\text{TO}}$  (dashed) and  $(B - V)_{\text{RGB}}$  (dash-dotted) from our isochrones. All quantities are given for two metallicities,  $Z = 2 \cdot 10^{-4}$  (thick lines) and  $10^{-2}$  (thin lines).

It was used by Vandenberg & Durrell (1990) for relative age determinations. The obvious difficulty lies in a precision estimate of  $V_{\text{tip}}$ , since the number of stars at the tip is very low due to the increased evolutionary speed in this phase.

$\Delta V_{\text{ZAHB}}^{\text{Bump}}$ : If the photometry is not deep enough to reach the TO, it is in principle possible to use  $\Delta V_{\text{ZAHB}}^{\text{Bump}} = V_{\text{Bump}} - V_{\text{ZAHB}}$  (Ferraro et al. 1999). The age sensitivity is very weak, amounting to  $\approx 0.5$  mag over 10 Gyr only, such that  $\Delta V_{\text{ZAHB}}^{\text{Bump}}$  will not provide any additional age restriction in comparison with the other indicators (Cassisi & Salaris 1997). The colour at which  $V_{\text{ZAHB}}$  is to be taken, is free to choose, but must be consistent between observed data and isochrone.

$\Delta(B - V)_{\text{TO}}^{\text{RGB}}$ : The preferred horizontal (here in  $(B - V)$ ) age indicator is  $\Delta(B - V)_{\text{TO}}^{\text{RGB}} = (B - V)_{\text{RGB}} - (B - V)_{\text{TO}}$  (Vandenberg et al. 1990; Salaris & Weiss 1997; Buonanno et al. 1998). As mentioned in Sect. 2.1, we take  $(B - V)_{\text{RGB}}$  at a brightness 2.5 mag brighter than the TO. This point represents a compromise between minimizing photometric errors and stellar density on the RGB (Rosenberg et al. 1999). The general problems with colours rests with the uncertainties in both the calculations (convection theory) and the transformations between theoretical  $T_{\text{eff}}$  and colour. However, by using a differential quantity, systematic uncertainties related to the models' metallicity can probably be avoided. A further reduction of the influence of these uncertainties can be achieved by using  $\Delta(B - V)_{\text{TO}}^{\text{RGB}}$  for relative ages as in Salaris & Weiss (1998).

$\Delta(B - V)_{\text{TO}}^{\text{Bump}}$ : Alternatively, the colour difference between bump and TO can be used, which is very similar to  $\Delta(B - V)_{\text{TO}}^{\text{RGB}}$ , but is taken at the location of a physical feature.


Hatzidimitriou (1991) presented a further “age calibrator”, using the colour difference between RGB and the red part of the HB. However, in view of the variety of HBs we encounter in a sample of GC and the unknown influence of the second parameter, we chose not to include this in our investigation.

### 3. Cluster data

#### 3.1. Cluster sample

This work is concerned with developing a more complete approach to globular cluster dating. The observational data should be comprehensive in terms of metallicity range and consistent to avoid differential effects in the quality of the method, which could be due to the differences in the database. For this reason, the photometric cluster catalogues by Rosenberg et al. (2000b) or Piotto et al. (2002a) are excellent choices. The first catalogue was used by Salaris & Weiss (2002), and we decided to use the latter one, which contains 74 galactic GCs in the HST F439W and F555W filters. The HST observations concentrated in particular on the central parts of the clusters. We obtained the data from the Padua web page <sup>1</sup>. The catalogue conveniently offers reddening-corrected Johnson colours ( $B$ ,  $V$ ), which were obtained by Piotto et al. (2002a) from an iter-

<sup>1</sup> <http://dipastro.pd.astro.it/globulars>



**Fig. 5.** Mean ridge line for the cluster M 5 as determined by us (crosses) and by Sandquist et al. (1996) from their own data (diamond symbols).

ative data calibration. While the authors do not claim the reddening to be always correct, in particular in the case of highly reddened clusters, our differential age indicators are mostly rather insensitive to reddening.

We deselected 28 clusters due to the low number of stars, or a significant scatter in the CMDs, or a lack of any discernible HB. The final number of clusters used was therefore 46 (Table 6).

For the metallicities ( $[\text{Fe}/\text{H}]$ ) we used the values on the scale of Zinn & West (1984,  $[\text{Fe}/\text{H}]_{\text{ZW}}$ ) given in the cluster catalogue, and transformed them to the Carretta & Gratton (1997,  $[\text{Fe}/\text{H}]_{\text{CG}}$ ) scale with the relation given by Carretta et al. (2001), unless they were available from Rutledge et al. (1997).

### 3.2. Data extraction


#### 3.2.1. Mean ridge line

The first step in preparing the photometric data for our method is to determine the mean ridge line. After exclusion of obvious HB stars by eye, we follow the standard procedure described, e.g. in Cho et al. (2005): After determining the mean colour of stars in brightness bins of 0.25 mag, stars with colour more than  $2.5\sigma$  different from the mean are removed. Then, the determination of mean and variance are repeated until convergence is reached, usually in less than 5 iterations. Typically, around 5% of the initial non-HB objects are removed. We also tested using the median instead of the mean, but the difference in the final ridge line was not recognizable.

Figure 5 shows, as an example, our ridge line and that of Sandquist et al. (1996) for the cluster M 5. It also shows that along the RGB the ridge line data points are not forming a smooth line due to statistical variations at low star numbers per brightness interval. Therefore, we performed a hyperbolic fit to the data following Rosenberg et al. (1999) and Saviane et al. (2000).

#### 3.2.2. ZAHB brightness

The ZAHB brightness was determined using the method presented by Salaris & Weiss (1997): The HB is divided into colour bins with width sufficiently large to contain at least 10 stars. Each colour bin is then subdivided into brightness bins of 0.05 mag, and the number of stars in each of them is used for a histogram of the colour bin. The criterion for the ZAHB is that 90% or more of all stars are above the ZAHB and that the number of stars drops by at least a factor of two from the bin above to that below the ZAHB level. The error is estimated from the brightness width. Not all colour bins allow the determination of the ZAHB at this colour and under these conditions, and thus the ZAHB line is obtained by interpolation between those colours where the method is successful. To check the reliability of our method, we compared with Ferraro et al. (1999), who determined, among other parameters, also  $V_{\text{ZAHB}}$  for 61 GCs. Figure 6 shows the result of the comparison, which confirms our method completely.



**Fig. 6.** Comparison of our derived values for  $V_{\text{ZAHB}}$  and those obtained by Ferraro et al. (1999) for clusters in common. The solid line is a linear fit through the data points, the dashed one the 1:1 line for comparison.

$V_{\text{ZAHB}}$  can be compared directly with the theoretical predictions, and therefore yields immediately a distance to the cluster. We have also compared these distances to those by Ferraro et al. (1999) and found that our distances are on average 0.1 mag larger. This is the immediate consequence of the Straniero et al. (1997) stellar models used by Ferraro et al. (1999). We emphasize again that the distances are not necessary for any age indicator we are using, but are just for the purpose of comparing isochrone with data in the CMD.

### 3.2.3. The turn-off region

For the cluster TO one should select the bluest point along the MS. However, as discussed by Salaris et al. (1997) for the case of M 15, the uncertainties in the mean color of brightness bins together with the near-vertical CMD introduces an uncertainty of 0.1 to 0.2 mag. We followed the method by Chaboyer et al. (1996) who fitted the mean ridge line around the TO by a fifth-order polynomial  $(B - V) = f(V)$ , and then took the bluest point along this curve as the TO. This point does not necessarily have the same colour as the bluest point of the ridge line. The error  $\sigma(B - V)_{\text{TO}}$  introduced is estimated (Press et al. 1992) by the expression

$$\sigma(B - V)_{\text{TO}} = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \bar{x}_i)^2}. \quad (1)$$


Here  $N$  is the number of ridge points used for the fit, and  $x_i$  and  $\bar{x}_i$  are the measured and the fitted colours in each bin. It is implied that this error is also the typical error of the TO colour,  $\sigma(B - V)_{\text{TO}}$ , assuming that the polynomial fits the TO region well. The error in  $V_{\text{TO}}$  can then be estimated<sup>2</sup> by

$$\sigma(V_{\text{TO}}) = \frac{\sigma_{\text{fit}}}{\sqrt{N}}, \quad (2)$$

where  $\sigma_{\text{fit}}$  is the standard deviation in brightness of the  $N$  ridge points within a  $3\sigma(B - V)_{\text{TO}}$  interval in  $(B - V)$  around the TO. The formal errors are typically 0.1 mag in  $V_{\text{TO}}$ , and 0.001 mag in  $(B - V)_{\text{TO}}$ .

The related quantities  $V_{\text{TO}}^b$  and  $V_{\text{TO}}^f$  are both determined from the same polynomial fit and TO colour with a similar estimate of the errors (see Fig. 7 for an illustrative example). Again, we compared our results with others from the literature; in this case we used Rosenberg et al. (1999), who determined  $V_{\text{TO}}$  from ground-based  $VI$ -photometry, for 16 clusters in common. Only in two cases, NGC 1261 and NGC 6093 (M 80), there is a difference of 0.3 mag, in all other cases, the agreement is of order 0.1 mag or better; the rms deviation is 0.125 (0.073 without NGC 1261 and M 80). Noting that  $\sigma(V_{\text{TO}})$  is 0.06–0.09 for all these clusters, the agreement is excellent.

<sup>2</sup> basic statistical properties can be found in many textbooks on the subject; throughout this paper we used the one by Brandt (1999)



**Fig. 7.** Fit to the mean ridge line of NGC 362 in the TO region and the location of TO, TO<sup>b</sup> and TO<sup>f</sup> with their respective errors.

Finally, we compared our  $\Delta V_{\text{TO}}^{\text{ZAHB}}$  values with those by De Angeli et al. (2005), obtained from the same HST photometry. Most values agree within the error bars (the typical error in De Angeli et al. (2005) is around 0.08 mag), but there are some prominent outliers; many of them (like NGC 4147, NGC 6273, and NGC 6544) have very blue HBs, for which our method is more uncertain). For the 35 clusters, for which we have  $\Delta V_{\text{TO}}^{\text{ZAHB}}$  from both sources, the rms is 0.15 mag. De Angeli et al. (2005) obtained  $\Delta V_{\text{TO}}^{\text{ZAHB}}$  also for the ground-based data by Rosenberg et al. (1999); the comparison there resulted in an rms deviation of only 0.09 mag. This better agreement is most likely due to the use of the same method of “template HBs” for determining the HB levels.

### 3.2.4. The red giant branch bump

To identify the RGB bump, the luminosity function (omitting HB and AGB stars) is needed. Piotto et al. (2002a) investigated the completeness of the luminosity function (LF) by artificial star experiments, concluding that the lower MS is complete only to a level of 0.7. In principle, LFs can be used for age determinations, as they allow the identification of the subgiant-branch and bump luminosity and the relative number of stars in different luminosity bins, which is age-sensitive, too. Also, the shape of the LF at the subgiant-branch is age-dependent, and this was used by Zoccali & Piotto (2000) to determine the ages of 18 clusters. However, incompleteness severely affects the usefulness of the LF and therefore we did not pursue this route any further.


To locate the bump, we used brightness bins of 0.05 mag, and followed Ferraro et al. (1999) and Riello et al. (2003) by requiring the coincidence of a stellar overdensity in a brightness bin and of a change of slope in the integrated LF. Figure 8 shows the example of NGC 362. For ten clusters in the sample, the bump could not be identified due to a too large scattering of star numbers along the RGB (Tab. 7). The colour of the bump was determined by using the average  $(B - V)$  of the bump and the two adjacent bins. The error was taken to be of the same size as the standard deviation of colours within these three bins. The theoretical counterpart was obtained by taking the upper bump brightness in the isochrone, which corresponds to the peak in the theoretical LF. Tables 6 and 7 list brightness and color of our age indicator points for all clusters taken into consideration.

## 4. Stellar models and isochrones

### 4.1. Stellar evolution code

For this project we computed new stellar models and isochrones with the Garching stellar evolution code described lately by Weiss & Schlattl (2000). The code incorporates the OPAL equation of state (Rogers et al. 1996) and the OPAL opacity tables (Iglesias & Rogers 1996) supplemented by the molecular opacities of Alexander & Ferguson (1994). In the calculations presented here, the equation of state (EOS) of Irwin (see Cassisi et al. 2003) has been used, which is based on the OPAL EOS. Diffusion of hydrogen and helium





**Fig. 8.** Observed luminosity function (LF) for NGC 362. The lower lines are the differential LF with brightness bins of sizes 0.05 mag (solid) and 0.2 mag (dotted), the upper solid line the integrated LF. The bump is identified by the vertical line, where a slope change in the integrated function coincides with a local overdensity of stars.

was taken into account using the diffusion coefficient treatment of Thoul et al. (1994), while that of heavier elements was ignored as it only adds a minor effect on the evolution (see Weiss & Schlattl 2000). Convection is treated according to standard mixing length theory with the Schwarzschild criterion for stability. The mixing length parameter was obtained from a calibration of the solar model (with diffusion), which yielded a value of 1.71. For the solar mixture, the values by Grevesse & Noels (1993) were employed. No mass loss is included.

Nuclear reaction rates for H-burning are mostly from Adelberger et al. (1998); the S-factor for  $^{14}\text{N}(p, \gamma)^{15}\text{O}$  was taken from the recent result by the LUNA collaboration (Formicola et al. 2004), and that for  $^3\text{He}(^3\text{He}, 2p)^4\text{He}$  from Junker et al. (1998). Helium burning reaction rates are by Caughlan et al. (1985) and Caughlan & Fowler (1988).

#### 4.2. Models calculated

We calculated models in the mass range of  $M/M_{\odot} = 0.3, \dots (0.1) \dots 1.3$ . For the initial helium content  $Y_0$  we used

$$Y_0 = Y_p + \frac{\Delta Y}{\Delta Z} Z$$

with  $Y_p = 0.245$  in agreement with the current concordance cosmological parameters and  $\Delta Y/\Delta Z = 2.5$ . Enhancement of  $\alpha$ -elements is taken into account. Our choice of the enhancement factors was guided by Tab. 1 in Salaris & Weiss (1998), and is similar, but slightly smaller than theirs. Most  $\alpha$ -elements are enhanced by 0.30–0.35 dex, and oxygen by 0.40 dex. The particular choice is not significant at this low metallicity (Salaris & Weiss 1998). At fixed metallicity  $Z$   $\alpha$ -enhancement implies a reduced value for  $[\text{Fe}/\text{H}]$  (see Weiss & Schlattl 2000 for a table) as compared to solar-scaled mixtures. Total metallicities are given in Tab. 1.

Our age indicators require theoretical ZAHB models, which we obtained by the method presented in Serenelli & Weiss (2005). Briefly, for each metallicity one stellar evolution model is followed completely from ZAMS to HB, including the full core helium flash, and without mass loss. From the ZAHB configuration of this model, other ZAHB models of lower mass are obtained by reducing the envelope mass. Serenelli & Weiss (2005) have shown that this method to high accuracy reproduces models obtained from full calculations. As the ZAHB brightness depends hardly on age (a change in age of 6 Gyr results in a change in  $\log L/L_{\odot}$  of only 0.02), it suffices to calculate one ZAHB per metallicity case for the age range of interest.

#### 4.3. Isochrones

To obtain isochrones from the stellar evolution tracks, we follow the method of equivalence points (EP) described, e.g., in Bergbusch & Vandenberg (1992). We defined in total 16 EPs, 5 on the MS, 7 more from the

**Table 1.** Chemical composition of stellar models with initial helium abundance  $Y_0$ , total metallicity  $Z$ , iron abundance with respect to hydrogen  $[\text{Fe}/\text{H}]$  and total metallicity with respect to hydrogen  $[\text{M}/\text{H}]$ 

$Z$	$Y_0$	$X_0$	$[\text{Fe}/\text{H}]$	$[\text{M}/\text{H}]$
$2 \cdot 10^{-4}$	0.2455	0.7543	-2.26	-1.97
$3 \cdot 10^{-4}$	0.2458	0.7539	-2.08	-1.79
$6 \cdot 10^{-4}$	0.2465	0.7529	-1.78	-1.49
$10^{-3}$	0.2475	0.7515	-1.56	-1.27
$2 \cdot 10^{-3}$	0.2500	0.7480	-1.25	-0.96
$3 \cdot 10^{-3}$	0.2525	0.7445	-1.07	-0.78
$6 \cdot 10^{-3}$	0.2600	0.7340	-0.77	-0.48
$10^{-2}$	0.2700	0.7200	-0.54	-0.25

TO to the bottom of the RGB, and 4 more along the RGB, including 2 at the start and end of the bump. Between the EPs, additional points, their number ranging from 10 on the MS to 100 on the RGB, were distributed at equal stellar age increments (for details, see Meissner 2005). This procedure yields a “normalized track” of given mass and metallicity. To get an isochrone of age  $t$  from these normalized tracks,  $T_{\text{eff}}(t)$  and  $\log L(t)/L_{\odot}$  values were interpolated for each mass value available. This results in a number of normalized points where the age of the lower mass track is higher, and that of the higher mass track is lower than  $t$ . Additional isochrone points are thus obtained by interpolating  $T_{\text{eff}}$  and  $\log L/L_{\odot}$  in  $t$  between corresponding normalized points. We constructed isochrones of age  $t = 4.0, \dots (0.5) \dots 18.0$  Gyr for each metallicity.

Finally, isochrones for the cluster metallicity are needed (measured by its  $[\text{Fe}/\text{H}]$ -value). Linear interpolation in metallicity was done between corresponding points on the available isochrones. We tested our method by interpolating to and comparing with available metallicities, and found an accuracy of order 0.001 mag in color, resp. 0.01 in brightness.

The theoretical isochrones were transformed into the observational  $M_V$  vs.  $(B - V)_0$  plane using the transformations by Cassisi et al. (2004), which, however, do not include  $\alpha$ -enhancements in the theoretical atmosphere models. We always chose the transformation with  $[\text{Fe}/\text{H}]_{\text{atm}} = [\text{M}/\text{H}]_{\text{iso}}$ , although this will render the transformed isochrones slightly too red on the RGB (Cassisi et al. 2004).

Alternatively, we have tested the Yale-transformations (Green 1988), which results in bluer RGBs than the Cassisi-transformation. The influence on the determined ages was tested by generating a 10 Gyr isochrone for  $Z = 0.001$  with the transformations by Cassisi et al. (2004), and then fitting this with isochrones using the Yale-transformations. Depending on the age indicator the transformations change the inferred age between 0.3 Gyr (for TO-indicators) and 1.1 Gyr (indicators involving the bump). However, for  $\Delta(B - V)_{\text{TO}}^{\text{RGB}}$  alone the change is +5 Gyr, when using the Yale transformations.

The theoretical luminosity function is obtained from the isochrone easily, as the mass along it is known, after folding it with an appropriate IMF. We choose an IMF exponent of  $s = 1.17$  (appropriate for globular cluster cores, see Zoccali & Piotto 2000), with  $s = 3.35$  being the Salpeter-value. The influence of  $s$  is almost exclusively on the MS; since we need the LF only for the bump location, our results are almost completely independent of  $s$ .

#### 4.4. Theoretical values for age indicators


The determination of the critical points along isochrones has been discussed before, or is straightforward. It is convenient to derive simple relations for their brightness and colour as function of age for all isochrone metallicities. An example ( $V_{\text{Bump}}$ ) is shown in Fig. 9. The analytical relations are obtained in all cases by fitting a third order polynomial through the data (Tables 8 and 9). No fitting of the metallicity dependence was done, however, for sake of accuracy.

## 5. Results

### 5.1. Method

The general method is to calculate for  $n$  indicators and for each isochrone age the value of  $M(t)$  as defined by

$$M(t) = \sum_{i=1}^n w_i \left( \frac{\Delta y_i - \widetilde{\Delta y}_i(t)}{\sigma(\Delta y_i)} \right)^2, \quad (3)$$




**Fig. 9.**  $V_{\text{Bump}}$  as function of age for all metallicities of Table 1 (crosses; increasing from bottom to top) and the third order polynomial fit (lines; see Table 8)

where  $\Delta y_i$  is the observed value of indicator  $i$ ,  $\widetilde{\Delta y}_i(t)$  the corresponding theoretical value for a particular isochrone age  $t$  and  $\sigma(\Delta y_i)$  the observational error (Brandt 1999). The  $w_i$  are additional weights for the different indicators, which will hereafter be set to 0 or 1 only, but could take any value. The value of  $\widetilde{\Delta y}_i(t)$  is computed from the polynomial fit and free of numerical errors. Systematic errors in the isochrones are not taken into account. Eq. 3 is resembling a  $\chi^2$ -test, but due to the unknown error statistics we cannot claim that it really is one; rather, it is a goodness-of-fit measure. Following Press et al. (1992) we introduce the quantity  $Q$  as a measure of the probability that the apparent discrepancies between data (age indicator values) and model (isochrone) are by chance. We generally chose  $Q = 0.001$  as the critical value, below which we reject models. This value is used as an acceptable measure even for cases with presumably not normally distributed errors. Another requirement is that  $M$  should be less than the number of degrees of freedom, which in our case are the 8 age indicators (equivalent to the normalized  $\chi^2 < 1$ ).

Our best estimate for the age  $t$  is then that of the isochrone with minimal  $M(t)$ , i.e. the age of the isochrone which appears to fit best all indicators simultaneously with equal weights assigned to them, and  $Q > 0.001$  implying that the remaining differences could be by chance due to the measurement errors. We add that we are testing only the isochrones available, which implies an age resolution of 0.5 Gyr. A refined grid of isochrones or use of the polynomial fits of Tables 8 and 9 might possibly yield lower  $M$ -values than those given below. We have tested several cases with various combinations of age indicators (equivalent to setting the weights in Eq. 3 to 1); they are given in Table 2.

**Table 2.** Combinations of age indicators. Given is the weight  $w_i$  in Eq. 3 for each indicator.

Indicator	Indicator set								
	A	B	C	D	E	F	BC	EF	S
$\Delta V_{\text{TO}}^{\text{ZAHB}}$	1	1	1	1	1	1	1	1	1
$\Delta V_{\text{TOb}}^{\text{ZAHB}}$	1	1	1	1	1	1	1	1	1
$\Delta V_{\text{TOF}}^{\text{ZAHB}}$	1	1	1	1	1	1	1	1	1
$\Delta V_{\text{TO}}^{\text{Bump}}$	0	1	0	0	0	0	1	0	1
$\Delta V_{\text{ZAHB}}^{\text{Bump}}$	0	0	1	0	0	0	1	0	1
$\Delta V_{\text{TO}}^{\text{tip}}$	0	0	0	1	0	0	0	0	1
$\Delta(B - V)_{\text{TO}}^{\text{RGB}}$	0	0	0	0	1	0	0	1	1
$\Delta(B - V)_{\text{TO}}^{\text{Bump}}$	0	0	0	0	0	1	0	1	1



**Fig. 10.**  $M(t)$  for various age indicators for the cluster NGC 4833; they are:  $\Delta V_{\text{TO}}^{\text{ZAHB}}$  (solid),  $\Delta V_{\text{TO}}^{\text{ZAHB}}$  (dotted),  $\Delta V_{\text{TO}}^{\text{ZAHB}}$  (dashed),  $\Delta V_{\text{TO}}^{\text{Bump}}$  (dash dot),  $\Delta V_{\text{ZAHB}}^{\text{Bump}}$  (dash dot dot dot),  $\Delta(B - V)_{\text{TO}}^{\text{RGB}}$  (long dash) and  $\Delta(B - V)_{\text{TO}}^{\text{Bump}}$  (diamond symbols)

### 5.2. A sample result: NGC 4833

NGC 4833 is a cluster with  $[\text{Fe}/\text{H}]_{\text{CG}} = -1.71$ ; its CMD contains about 6700 stars and shows very well-defined structures. The HB is rather blue, but with a sufficient number of stars at  $(B - V)_{\text{TO}}$  to determine the ZAHB.

Figure 10 shows  $M(t)$  for various indicators. It is immediately clear that they do not agree at all. The three TO-related vertical indicators (case A) yield an age of 13.5 Gyr with  $M(13.5) = 2.6$  and  $Q = 0.457$ , while  $\Delta V_{\text{TO}}^{\text{Bump}}$  (dash-dotted line) has a sharp minimum at 8 Gyr, and  $\Delta(B - V)_{\text{TO}}^{\text{RGB}}$  (long-dashed) at 13.5 Gyr. Notice that  $\Delta(B - V)_{\text{TO}}^{\text{Bump}}$  (diamond symbols) has an extremely flat curve at low  $M$ , which is due to the large error in this indicator. It provides therefore hardly any restriction on the global fit. Using all indicators (case S) results in a minimum at 11.5 Gyr, which however, is pure coincidence, as  $M(11.5) = 191$  and  $Q < 0.001$ . Clearly,  $V_{\text{Bump}}$  is too low (bump too bright; it gets fainter with increasing age), which is also the reason why  $\Delta V_{\text{ZAHB}}^{\text{Bump}}$  results in ages above the maximum age of 18 Gyr for which we have isochrones. Since  $V_{\text{Bump}}$  from the isochrones is too small,  $\Delta V_{\text{ZAHB}}^{\text{Bump}}$  has a too negative value for the actual age. To compensate, age has to be increased, and since the indicator is only weakly sensitive, the increase has to be substantial. The effect is in the opposite direction of that of  $\Delta V_{\text{TO}}^{\text{Bump}}$ , since the latter quantity has an opposite dependence on  $V_{\text{Bump}}$ .

We add that  $\Delta V_{\text{TO}}^{\text{tip}}$  yields an unrealistically low age of 4.5 Gyr, which is most likely due to an underestimate of the tip brightness in the data.


### 5.3. NGC 6934, a moderately metal-rich cluster

Our second example is that of NGC 6934 ( $[\text{Fe}/\text{H}] = -1.30$ ); the CMD contains close to 9500 stars and the HB is well populated. As in the previous case we see (Table 3) that the first three indicators agree very well and thus the “case A” determination yields an age of 9.0 Gyr with  $Q = 0.857$ . Due to the higher number of stars,  $V_{\text{tip}}$  can be determined more accurately and thus  $\Delta V_{\text{TO}}^{\text{tip}}$  yields an age in agreement (set D). In contrast, inclusion of the bump-brightness yields discrepant ages and cases B and BC of Tab. 2 have a much lower confidence level (Fig. 11). The same is true when including the colour indicator  $\Delta(B - V)_{\text{TO}}^{\text{RGB}}$  (case E), which requires a higher age. This implies that the RGB is too blue, relative to the TO, although our colour transformations have a systematic shift to the red (Sect. 4.3). Cases C and F agree quite well with the turn-off indicators. Overall (case S) the age is 9.5 Gyr, but the confidence in this result is very low, and  $M(9.5) = 150$ .

We have also investigated all other clusters obtaining consistent results: the bump in our isochrones appears to be too luminous thereby leading to discrepant results when using  $\Delta V_{\text{TO}}^{\text{Bump}}$  or  $\Delta V_{\text{ZAHB}}^{\text{Bump}}$ . The TO indicators give very consistent results with very high confidence level, and the quality of  $\Delta V_{\text{TO}}^{\text{tip}}$  depends on the number of stars available for the location of the RGB tip, as should be expected.

**Table 3.** Indicator value, age  $t_{min}$ , minimum value  $M(t_{min})$  and fit quality  $Q$  for all age indicators applied to NGC 6934


Indicator	Value	$t_{min}$	$M(t_{min})$	$Q$
$\Delta V_{TO}^{ZAHB}$	$3.24 \pm 0.08$	9.0	0.0840	0.772
$\Delta V_{TOb}^{ZAHB}$	$2.69 \pm 0.05$	9.5	0.0677	0.795
$\Delta V_{TOF}^{ZAHB}$	$4.13 \pm 0.06$	9.0	0.0044	0.947
$\Delta V_{TO}^{Bump}$	$3.62 \pm 0.06$	7.0	0.2170	0.641
$\Delta V_{ZAHB}^{Bump}$	$-0.38 \pm 0.05$	13.0	0.0030	0.956
$\Delta V_{TO}^{tip}$	$6.5 \pm 0.1$	8.0	0.0016	0.968
$\Delta(B - V)_{TO}^{RGB}$	$0.293 \pm 0.005$	12.0	0.0055	0.941
$\Delta(B - V)_{TO}^{Bump}$	$0.39 \pm 0.02$	17.0	0.0001	0.994

**Fig. 11.** Combinations of age indicators for NGC 6934: Case A (solid), Case B (dotted), Case C (dashed), Case D (dash dot), Case E (dash dot dot dot), and Case S (diamonds). Case F is not shown as it coincides with A.

#### 5.4. The special case of M 15

We finally mention the peculiar case of M 15. Applying our method straightforwardly to the data by Piotto et al. (2002b) we obtain around 9 Gyr only (see Table 6), with very low  $Q$ -values. The ZAHB is determined here at the TO-colour. Overlaying the resulting isochrone of 9 (case A) and 11 Gyr (case S) with the determined cluster ridge line aligned at the ZAHB (Fig. 12; upper panel), the corresponding distance modulus amounts to 15.32 mag. The figure shows that the isochrone is a very bad match to the CMD; it is too blue for all of the MS and too red on the RGB (one nicely realizes how the “best” isochrone is a compromise match to all indicators). The inset, however, indicates that there might be a problem with the identification of the HB, as the majority of stars is to the blue and at a higher brightness. Piotto (private communication) pointed out that the data were obtained in HST *snapshot* mode, which does not allow to determine the mean brightness of RR Lyr variables, in which M 15 is very rich. Thus a number of RR Lyr stars will be caught at minimum brightness. This was one reason for Recio-Blanco et al. (2005) to introduce the concept of template clusters. Identifying the blue part as the true HB, we obtain the results of the lower panel: our method yields 10.2–11.7 Gyr for case A, and 11.5 Gyr for case S; the ZAHB level corresponds to a distance modulus of 15.10 mag, and the 12 Gyr isochrone matches the CMD quite well, except for being too blue on the lower main-sequence.

As the photometry appears to be difficult to interpret, we used as an alternative that by Durrell & Harris (1993). The ridge line we determine from this source agrees very well with that derived by Durrell and Harris themselves. Our age indicators now yield ages between 10.5 and 13.5 Gyr. The inferred distance modulus is 15.37. From these data the bump could not be located such that the corresponding indicators cannot be used. The remaining indicators together yield 12.0 Gyr in excellent agreement with the  $11.7 \pm 0.8$  Gyr by Salaris & Weiss (2002). Note that the CMD of Durrell & Harris (1993) does not show the splitting into a bluer, brighter and a redder, fainter part, as that by Piotto et al. (2002b) does (visible also on the original




**Fig. 12.** Fit of the M 15 photometric data by Piotto et al. (2002b). In the upper panel our standard procedure was applied, fitting the HB at the TO colour. Isochrones of 9 and 11 Gyr are overlaid. For the lower panel the HB was fitted at the blue end, where the majority of stars is found, as can be seen in the inset of the upper panel. Here, the isochrone ages are 10 and 12 Gyr.


web-site [dipastro.astro.unipd.it/globulars/databases](http://dipastro.astro.unipd.it/globulars/databases)). It is also less pronounced in the ground-based  $(V - I)$  data by Rosenberg et al. (2000a). Whether the data by Piotto et al. (2002b) are erroneous, or our method of determining the ridge line fails in this case, or whether the core of M 15 indeed shows an anomalous CMD remains to be investigated. This case demonstrates that our method might also be capable of finding inconsistencies in the data.

### 5.5. Quality of age indicators

Figure 13 shows the quality of the case A indicators fits for all clusters investigated. We notice no correlation with metallicity, and that the majority of cases have  $Q > 0.5$ . In fact, all but three clusters (NGC 6093, 6642,



**Fig. 13.** Fit quality of case A (TO-ZAHB-related) age indicators for all 46 clusters as function of  $[\text{Fe}/\text{H}]$ . Diamonds indicate the clusters for which  $Q < 0.001$ .



**Fig. 14.** Fit quality of case B (case A plus  $\Delta V_{\text{TO}}^{\text{Bump}}$ ) age indicators for all 46 clusters as function of  $[\text{Fe}/\text{H}]$  (diamond symbols). Results for bump brightnesses corrected artificially by +0.25 mag are shown as triangles

and 6864) have  $Q > 0.001$ , i.e. the determined age is unlikely to be just a chance result. This implies that our isochrones reproduce the critical TO-region very well.

The situation is very different for case B (Fig. 14; diamond symbols), when we add  $\Delta V_{\text{TO}}^{\text{Bump}}$  as an additional age indicator: no single cluster has  $Q > 0.1$  and 29 clusters are below our rejection value of 0.001. As the bump appears to be too bright by about 0.2–0.3 mag, we artificially made it fainter by applying an ad-hoc correction of +0.25 mag, which is also indicated by Zoccali & Piotto (2000). However, the result is basically the same (triangle symbols in Fig. 14), such that an even larger correction would be needed. Indeed, with 0.35 mag, the ages obtained from  $\Delta V_{\text{TO}}^{\text{Bump}}$  begin to be consistent with those of case A. Note that adding  $\Delta V_{\text{ZAHB}}^{\text{Bump}}$  to case A (i.e. case C) does not degrade the fit quality significantly due to the rather shallow run of  $M(t)$  for this indicator (Fig. 10).

The inclusion of the colour indicators (cases E or F) results in no change of the fit quality at all (case F, because of the large errors for this indicator; see Sect. 5.2), or in a larger number of clusters to be rejected (case E, inclusion of  $\Delta(B - V)_{\text{TO}}^{\text{RGB}}$ ): 15 out of 46 clusters are below  $Q = 0.001$ . Here, the theoretical uncertainty in the colour transformations is crucial and should be included in the calculation of  $M$ , although in this

**Table 4.** Comparison of bump properties for stellar models of  $1 M_{\odot}$  and  $Z = 0.001$ . The references are: Girardi et al. (2000, Padova), Cariulo et al. (2004, Pisa), Pietrinferni et al. (2004, BASTI), and Kim et al. (2002, Y<sup>2</sup>)

	Padova	Pisa	BASTI	Y <sup>2</sup>	this work
Age (Gyr)	6.514	6.331	6.023	6.769	6.222
$\log L/L_{\odot}$	1.971	2.089	2.106	2.101	2.153
$\log T_{\text{eff}}$	3.698	3.689	3.694	3.689	3.693
metal mix.	solar	solar	$\alpha$ -enh.	$\alpha$ -enh.	$\alpha$ -enh.
diffusion	no	yes	no	yes	yes

case the usefulness of  $\Delta(B - V)_{\text{TO}}^{\text{RGB}}$  would just disappear. What is really needed are reliable and accurate transformations.

Finally,  $\Delta V_{\text{TO}}^{\text{tip}}$  leads to systematically too low ages, as is expected from the problem of finding the true RGB tip in the observed CMD. One probably should rather evaluate at what level the observed tip would be, given the number of stars observed in the uppermost bins of the CMD.

Table 5 contains the results for case A, which is the most reliable one, and case S in Table 5 for all clusters in comparison with the ages found by Salaris & Weiss (2002). If there is an age range given, it is that part of  $M(t)$  with  $Q > 0.32$ , which corresponds to the  $1\sigma$  range or a confidence level of 68%. If only one age is listed, it is that where  $M(t)$  has a minimum, but where this minimum has a lower confidence level. This is particular true for all ages determined using all indicators (case S). Note that very narrow age ranges usually are not due to very sharp minima in  $M(t)$ , but due to a minimum value only slightly above the  $1\sigma$  level.

### 5.6. Results for alternative isochrones


The main conclusion of the preceding sections is that our bump brightnesses are too high. We have therefore looked into alternative sets of isochrones. Since Cassisi & Salaris (1997) and Riello et al. (2003) claimed an improved agreement between observed and theoretical  $V_{\text{Bump}}$  for their isochrones, we took the publically available BASTI<sup>3</sup> isochrones by Pietrinferni et al. (2004) and applied our method to them. A first inspection already confirmed that BASTI-bumps are dimmer than ours by a few tenths of a magnitude. Figure 15 shows several fitting functions  $M(t)$  for the two sample clusters NGC 4833 and NGC 6934 using the BASTI isochrones. Obviously, in these cases, the bump indicators are in much better agreement with those of set A, and in addition, the remaining deviations are in both directions, that is, they are unsystematically. The total fit (case S) is still rather poor, but this is mainly due to the colour indicators, included here, too. Figure 16 illustrates the improvement in  $Q$  (case B indicators) for at least 14 clusters when using the BASTI isochrones and Table 5 contains the corresponding ages when considering both TO and bump indicators.

In Fig. 17 we show the bump location as function of  $[\text{Fe}/\text{H}]$  for our isochrones, that of BASTI, and those by Bergbusch & Vandenberg (2001). Obviously, our bump location is systematically lower (brighter) than that of the other two sources. In Table 4 we compare bump locations and ages of  $1 M_{\odot}$  stars with  $Z = 0.001$  for a number of authors. Obviously, there is quite a scatter in the bump properties. The Padova-bump has the lowest brightness due to the additional effect of overshooting from the lower convective boundary. The bump of our model is the brightest, and the age of BASTI's the youngest. We are currently comparing in detail our RGB-models with those by Pietrinferni et al. (2004) to find out the reasons for the differences. At the present time the scatter seems to be an intrinsic uncertainty in the various model calculations. Further evidence for the rather uncertain bump brightness comes from the comparison of a  $0.9 M_{\odot}$  star ( $Z = 0.0001$ ) by Straniero et al. (1997), recomputed by M. Limongi (private communication) with an updated version of his stellar evolution code: the luminosity of the new model agrees with that of our model within  $\Delta \log L/L_{\odot} = 0.009$ , while the BASTI model is 0.058 dex fainter. However, all models of Straniero et al. (1997), which were used in Valenti et al. (2004) for comparison with infrared bump data, were consistently fainter than ours by  $\approx 0.05$  dex.

If we use only the case A indicators, the fit quality of the BASTI is very similar to that of our isochrones, but the ages are on average lower by about 0.5-1.5 Gyr.

<sup>3</sup> URL: <http://www.te.astro.it/BASTI>






**Fig. 15.** Fitting functions for NGC 4833 (upper panel) and NGC 6934 (lower panel) using the BASTI (Pietrinferni et al. 2004) isochrones. The linetypes correspond to case A (solid), B (dotted), C (dashed), and S (dash-dotted) in both cases

## 6. Discussion


The aim of this work has been to attempt for the first time to simultaneously fit several age indicators of globular cluster colour-magnitude-diagrams. Such a more global approach removes a certain arbitrariness in the method of age determination and results in an overall best-fitting isochrone, which would not necessarily be the one reproducing the TO best. The bottom line of this paper is, however, that we are far from being able to do so. Instead, our method pointed to a number of inconsistencies and problems with our theoretical models. Apart from the case of M 15 (see Sect. 5.3), where the data deserve a second look by the observers, we doubt that a different data source would alter our findings. Neither did we modify the original data (Piotto et al. 2002b), for example, by changing the reddening.

The most severe problem lies in the bump brightness, which, for our models consistently is too large by up to 0.3 mag. This may indicate the need for overshooting, which would result in a fainter bump. At the present time, however, it is evident that different theoretical models, which nominally all use up-to-date physical input, differ widely in their predictions such that one first should look into code-specific reasons. We tested our method with the BASTI (Pietrinferni et al. 2004) isochrones, which show lower bump brightness, and found with respect to bump-related age indicators some improvement, but no final ages of consistently high confidence levels.

The second problem is the determination of the RGB tip brightness, which is due to the low number of stars observed in this phase. This leads to a consistent underestimate for the age derived from  $\Delta V_{\text{TO}}^{\text{tip}}$ , which



**Fig. 16.** Fit quality  $Q$  for case B indicators as function of  $[Fe/H]$  for both our own isochrones (diamonds) and those from Pietrinferni et al. (2004, crosses).



**Fig. 17.** Bump brightness for isochrones of 10 Gyr as function of metallicity  $[Fe/H]$  for three sets of models: Bergbusch & Vandenberg (2001, crosses), Pietrinferni et al. (2004, diamonds), and this paper (triangles).

could be cured by either applying a one-sided error-bar, or by a simulation of observations, or by modifying the age indicator, for example, by determining the level of the 10% brightest stars above the HB. Generally, however, this just leads to the assignment of an appreciable error range which then would reduce the weight of that age indicator. This could also be done by assigning a lower weight  $w_i$  in Eq. (3).

Finally, the colour-related indicators on the RGB are unreliable due to the uncertainties of colour-transformations. They, too, should receive a lower weight, or should be used for differential age determinations only, where the uncertainties are of less influence. The solution lies in better and calibrated transformations.

The most reliable age indicators appear to be those related to  $V_{ZAHB}$  and the three TO-brightness points. They give, for all cases, very consistent ages and the confidence level of the solution is always very high. They are, within the errors, often in agreement with  $\Delta V_{ZAHB}^{Bump}$  and  $\Delta V_{TO}^{tip}$ . Note that the overall best solution, even when omitting the most problematic indicators, does not yield the age of the classical indicator,  $\Delta V_{TO}^{ZAHB}$ , alone. This demonstrates that a global fit of cluster CMDs is indeed valuable.

**Table 5.** Ages (in Gyr) for galactic globular clusters from Case A (column 4), from Case S (column 5) and quoted from Salaris & Weiss (2002, column 6, “SW02”). The age range is that for which  $M(t)$  has  $Q > 0.32$  ( $1\sigma$ -range). If only one age is given, it is  $M_{\min}$ , with  $Q(M_{\min}) < 0.32$ . Columns 7 and 8 contain the result when using the BASTI-isochrones (Pietrinferni et al. 2004) for case A, and for case BC (see Table 2), which includes the bump brightness. Case S is not shown as it is not better than for our own isochrones.

Name		[Fe/H]	$t(A)$	$t_{\min}(S)$	$t(SW02)$	$t(A)_{BASTI}$	$t(BC)_{BASTI}$
NGC 104	47 Tuc	-0.78	11.9-13.6	11.5	$10.7 \pm 1.0$	9.9-11.5	9.5
NGC 362		-1.09	7.2-13.2	8.5	$8.7 \pm 1.5$	6.3-11.2	6.8-10.8
NGC 1261		-1.08	8.6-14.6	9.5	$8.6 \pm 1.1$	7.7-12.2	10.5-10.6
NGC 1851		-1.03	8.8-9.9	9.5	$9.2 \pm 1.1$	7.9- 8.7	7.9- 9.8
NGC 1904	M 79	-1.37	10.6-12.0	11.0	$11.7 \pm 1.3$	10.1-11.8	12.0
NGC 2808		-1.11	9.4-10.0	10.0	$9.3 \pm 1.1$	8.4- 8.6	8.0
NGC 3201		-1.24	9.0	4.0	$11.3 \pm 1.1$	7.3- 8.8	4.0
NGC 4147		-1.50	9.5	6.0	—	8.4- 9.0	4.0
NGC 4590	M 68	-2.00	8.5-11.0	9.5	$11.2 \pm 0.9$	9.2-12.5	9.0
NGC 4833		-1.71	12.5-14.0	11.5	—	12.1-14.2	12.0
NGC 5024	M 53	-1.89	13.0	11.0	—	11.4-13.1	10.0
NGC 5634		-1.61	8.4-10.1	9.0	—	8.8-11.2	7.5
NGC 5694		-1.74	12.5	13.0	—	11.2-13.0	12.0
NGC 5824		-1.67	9.2-10.4	11.0	—	8.8- 9.5	9.0
NGC 5904	M 5	-1.12	9.9-10.9	11.0	$10.9 \pm 1.1$	8.7- 9.7	8.5- 9.6
NGC 5927		-0.64	11.3-13.6	8.0	—	8.6-10.4	7.0
NGC 5986		-1.35	12.0	11.0	—	11.0	10.5
NGC 6093	M 80	-1.47	10.0	10.5	$12.4 \pm 1.1$	9.0	8.0
NGC 6171	M 107	-0.95	9.0	9.5	$11.7 \pm 0.8$	7.3- 7.9	7.5
NGC 6205	M 13	-1.33	11.0	15.5	$11.9 \pm 1.1$	9.7-10.7	10.5
NGC 6266	M 62	-1.02	11.0-12.1	13.0	—	10.0	9.5
NGC 6273	M 19	-1.53	11.1-18.0	15.0	—	17.0	17.0
NGC 6284		-1.17	11.0	11.0	—	8.9-10.0	10.0
NGC 6287		-1.90	12.5	10.5	—	10.2-11.6	9.0
NGC 6293		-1.73	7.9-8.8	10.0	—	8.3- 8.7	8.5
NGC 6304		-0.68	12.3-14.9	10.0	—	10.3-12.6	9.0
NGC 6356		-0.69	12.0-18.0	12.5	—	10.1-15.6	10.0
NGC 6362		-0.96	10.5	9.0	$11.0 \pm 1.3$	8.5- 9.5	8.5
NGC 6522		-1.21	13.9-16.1	16.0	—	12.2-15.4	13.0
NGC 6544		-1.20	7.1-8.5	8.0	—	6.7- 7.5	7.0
NGC 6584		-1.30	9.0	10.5	$11.3 \pm 1.4$	7.9- 8.5	8.0
NGC 6624		-0.70	12.0	10.5	$10.6 \pm 1.4$	10.0	8.5
NGC 6637	M 69	-0.78	12.0-14.5	11.5	$10.6 \pm 1.4$	10.0	9.5
NGC 6638		-0.73	12.0	11.5	—	10.0-12.1	10.0
NGC 6642		-1.08	10.0	10.0	—	8.5	8.5
NGC 6652		-0.81	11.5-11.8	8.5	$11.4 \pm 1.0$	9.1-10.8	7.5
NGC 6681	M 70	-1.35	10.9-13.3	9.5	$11.5 \pm 1.4$	10.6-13.0	9.5
NGC 6712		-0.94	10.2-14.7	9.5	$10.4 \pm 1.4$	8.7-12.1	9.5
NGC 6723		-0.96	10.0	10.5	$11.6 \pm 1.3$	8.3- 9.5	8.0
NGC 6838	M 71	-0.73	12.0	11.5	$10.2 \pm 1.4$	10.0	10.0
NGC 6864	M 75	-1.10	9.0	9.5	—	8.0	7.5
NGC 6934		-1.30	8.7-10.2	9.5	$9.6 \pm 1.5$	7.8- 8.7	9.0
NGC 6981	M 72	-1.21	9.5	9.5	—	8.4- 8.5	7.9- 8.5
NGC 7078	M 15	-2.02	8.6-9.2	11.0	$11.7 \pm 0.8$	8.0	9.0
NGC 7089	M 2	-1.31	11.8-13.7	12.5	—	10.9-11.6	11.0
NGC 7099	M 30	-1.92	13.7-15.4	13.0	$11.9 \pm 1.4$	12.8-15.0	14.0

*Acknowledgements.* We thank H. Schlattl for help with the numerical calculations, M. Salaris for valuable comments on the manuscript, S. Cassisi, and D. Vandenberg for making their models available to us, and M. Limongi for doing a separate new calculation. F. Meissner is grateful to the *Max-Planck-Institut für Astrophysik* for financial support.

*This paper has been typeset using a modified version of the Astron. & Astrophys. L<sup>A</sup>T<sub>E</sub>X macro package, version 6.0.*

## References

- Adelberger, E., Austin, S., Bahcall, J., et al. 1998, *Rev. Mod. Phys.*, 70, 1265
- Alexander, D. & Ferguson, J. 1994, *Astrophys. J.*, 437, 879
- Bergbusch, P. A. & Vandenberg, D. A. 1992, *ApJS*, 81, 163
- Bergbusch, P. A. & Vandenberg, D. A. 2001, *ApJ*, 556, 322
- Brandt, S. 1999, *Datenanalyse*, 4th edn. (Spektrum Akademischer Verlag)
- Buonanno, R., Corsi, C. E., Pulone, L., Fusi Pecci, F., & Bellazzini, M. 1998, *A&A*, 333, 505
- Cariulo, P., Degl'Innocenti, S., & Castellani, V. 2004, *A&A*, 421, 1121
- Carretta, E., Cohen, J. G., Gratton, R. G., & Behr, B. B. 2001, *AJ*, 122, 1469
- Carretta, E. & Gratton, R. 1997, *A&AS*, 121, 95
- Carretta, E., Gratton, R., Clementini, G., & Fusi Pecci, F. 2000, *ApJ*, 215
- Cassisi, S. & Salaris, M. 1997, *MNRAS*, 285, 593
- Cassisi, S., Salaris, M., Castelli, F., & Pietrinferni, A. 2004, *ApJ*, 616, 498
- Cassisi, S., Salaris, M., & Irwin, A. 2003, *ApJ*, 588, 862
- Caughlan, G. & Fowler, W. 1988, *Atomic Data and Nuclear Data Tables*, 40, 283
- Caughlan, G., Fowler, W., Harris, H., & Zimmerman, B. 1985, *Atomic Data and Nuclear Data Tables*, 32, 197
- Chaboyer, B., Demarque, P., Kernan, P. J., Krauss, L. M., & Sarajedini, A. 1996, *MNRAS*, 283, 683
- Chaboyer, B., Demarque, P., & Sarajedini, A. 1996, *ApJ*, 459, 558
- Cho, D., Lee, S., Jeon, Y., & Sim, K. 2005, *AJ*, 129, 1922
- De Angeli, F., Piotto, G., Cassisi, S., et al. 2005, *AJ*, 130, 116
- Durrell, P. R. & Harris, W. E. 1993, *AJ*, 100, 1420
- Ferraro, F. R., Messineo, M., Fusi Pecci, F., et al. 1999, *AJ*, 118, 1738
- Formicola, A., Imbriani, G., Costantini, H., et al. 2004, *Phys. Lett. B*, 591, 61
- Girardi, L., Bressan, A., Bertelli, G., & Chiosi, C. 2000, *A&AS*, 141, 371
- Gratton, R. G., Fusi Pecci, F., Carretta, E., et al. 1997, *ApJ*, 491, 749
- Green, E. M. 1988, in *Calbration of Stellar Ages*, ed. A. D. Philip (Schenectady, N.Y.: L. Davis Press), 81
- Grevesse, N. & Noels, A. 1993, *Physica Scripta*, T47, 133
- Hatzidimitriou, D. 1991, *MNRAS*, 251, 545
- Iglesias, C. & Rogers, F. 1996, *Astrophys. J.*, 464, 943
- Junker, M., D'Alessandro, A., Zavatarelli, S., & et al. 1998, *Physical Review C*, 57, 2700
- Kim, Y., Demarque, P., Yi, S., & Alexander, D. 2002, *ApJS*, 143, 499
- Meissner, F. 2005, Master's thesis, University of Munich
- Pietrinferni, A., Cassisi, S., Salaris, M., & Castelli, F. 2004, *ApJ*, 612, 168
- Piotto, G., King, I. R., Djorgovski, S. G., et al. 2002a, *A&A*, 391, 945
- Piotto, G., King, I. R., Djorgovski, S. G., et al. 2002b, *VizieR Online Data Catalog*, 339, 10945
- Press, W. H., Teukolsky, S. A., Vetterling, W. T., & Flannery, B. P. 1992, *Numerical Recipes in Fortran 77*, 2nd edn., Vol. 1 (Cambridge University Press)
- Recio-Blanco, A., Piotto, G., de Angeli, F., et al. 2005, *A&A*, 432, 851
- Renzini, A. & Fusi Pecci, F. 1988, *ARAA*, 26, 199
- Riello, M., Cassisi, S., Piotto, G., et al. 2003, *A&A*, 410, 553
- Rogers, F., Swenson, F., & Iglesias, C. 1996, *Astrophys. J.*, 456, 902
- Rosenberg, A., Aparicio, A., Saviane, I., & Piotto, G. 2000a, *A&AS*, 145, 451
- Rosenberg, A., Piotto, G., Saviane, I., & Aparicio, A. 2000b, *A&AS*, 144, 5
- Rosenberg, A., Saviane, I., Piotto, G., & Aparicio, A. 1999, *AJ*, 118, 2306
- Rutledge, G. A., Hesser, J. E., & Stetson, P. B. 1997, *PASP*, 109, 907
- Salaris, M. & Cassisi, S. 1997, *MNRAS*, 289, 406
- Salaris, M. & Cassisi, S. 1998, *MNRAS*, 298, 166
- Salaris, M., Cassisi, S., & Weiss, A. 2002, *PASP*, 114, 375
- Salaris, M., Degl'Innocenti, S., & Weiss, A. 1997, *ApJ*, 479, 665
- Salaris, M. & Weiss, A. 1997, *A&A*, 327, 107
- Salaris, M. & Weiss, A. 1998, *A&A*, 335, 943
- Salaris, M. & Weiss, A. 2002, *A&A*, 492
- Salpeter, E. 1955, *ApJ*, 121, 161
- Sandquist, E. L., Bolte, M., Stetson, P. B., & Hesser, J. E. 1996, *ApJ*, 470, 910
- Saviane, I., Rosenberg, A., Piotto, G., & Aparicio, A. 2000, *A&A*, 355, 966
- Serenelli, A. & Weiss, A. 2005, *A&A*, 442, 1041
- Straniero, O., Chieffi, A., & Limongi, M. 1997, *ApJ*, 490, 425

- Thoul, A., Bahcall, J., & Loeb, A. 1994, *Astrophys. J.*, 421, 828  
Valenti, E., Ferraro, F. R., & Origlia, L. 2004, *MNRAS*, 354, 815  
VandenBerg, D. A. 2000, *ApJS*, 129, 315  
Vandenberg, D. A., Bolte, M., & Stetson, P. B. 1990, *AJ*, 100, 445  
Vandenberg, D. A. & Durrell, P. R. 1990, *AJ*, 99, 221  
Weiss, A. & Schlattl, H. 2000, *A&A Supplement*, 144, 487  
Zinn, R. & West, M. J. 1984, *ApJS*, 55, 45  
Zoccali, M. & Piotto, G. 2000, *A&A*, 358, 943

# Online Material

**Table 6.** Data of the critical points of all Globular Cluster Color-Magnitude-Diagrams in our sample as determined by us (see Sect. 3); part I: the turn-off region and colours of turn-off, lower RGB and bump.

Name		[Fe/H]	$V_{TO}$	$V_{TO}^b$	$V_{TO}^f$	$(B - V)_{TO}$	$(B - V)_{RGB}$	$(B - V)_{Bump}$
NGC 104	47 Tuc	-0.78	17.43 ± 0.06	17.06 ± 0.01	18.22 ± 0.02	0.502 ± 0.001	0.829 ± 0.002	0.89 ± 0.02
NGC 362		-1.09	18.56 ± 0.07	18.16 ± 0.01	19.51 ± 0.02	0.373 ± 0.002	0.703 ± 0.003	0.78 ± 0.02
NGC 1261		-1.08	20.15 ± 0.08	19.68 ± 0.02	21.15 ± 0.02	0.432 ± 0.004	0.753 ± 0.004	0.85 ± 0.03
NGC 1851		-1.03	19.41 ± 0.07	18.95 ± 0.01	20.32 ± 0.02	0.459 ± 0.002	0.783 ± 0.003	0.87 ± 0.03
NGC 1904	M 79	-1.37	19.58 ± 0.07	19.10 ± 0.01	20.47 ± 0.02	0.441 ± 0.003	0.706 ± 0.003	0.88 ± 0.03
NGC 2808		-1.11	18.86 ± 0.06	18.42 ± 0.01	19.82 ± 0.02	0.396 ± 0.002	0.707 ± 0.002	0.78 ± 0.03
NGC 3201		-1.24	17.41 ± 0.08	17.02 ± 0.04	18.23 ± 0.03	0.493 ± 0.003	0.790 ± 0.010	0.74 ± 0.03
NGC 4147		-1.50	20.32 ± 0.07	19.84 ± 0.02	21.07 ± 0.04	0.401 ± 0.004	0.659 ± 0.008	0.65 ± 0.03
NGC 4590	M 68	-2.00	18.92 ± 0.09	18.39 ± 0.03	19.82 ± 0.03	0.382 ± 0.003	0.644 ± 0.006	0.73 ± 0.02
NGC 4833		-1.71	18.05 ± 0.09	17.57 ± 0.02	18.96 ± 0.02	0.407 ± 0.003	0.672 ± 0.004	0.79 ± 0.03
NGC 5024	M 53	-1.89	20.23 ± 0.08	19.72 ± 0.02	21.10 ± 0.02	0.389 ± 0.002	0.657 ± 0.002	0.75 ± 0.03
NGC 5634		-1.61	20.91 ± 0.09	20.46 ± 0.02	21.64 ± 0.03	0.369 ± 0.003	0.634 ± 0.003	0.71 ± 0.02
NGC 5694		-1.74	21.65 ± 0.07	21.19 ± 0.02	22.52 ± 0.03	0.424 ± 0.003	0.683 ± 0.004	—
NGC 5824		-1.67	21.51 ± 0.06	20.98 ± 0.02	22.42 ± 0.03	0.395 ± 0.002	0.659 ± 0.002	0.76 ± 0.02
NGC 5904	M 5	-1.12	18.35 ± 0.06	17.90 ± 0.02	19.25 ± 0.03	0.431 ± 0.001	0.732 ± 0.003	0.81 ± 0.03
NGC 5927		-0.64	18.73 ± 0.07	18.43 ± 0.01	19.50 ± 0.02	0.575 ± 0.003	0.954 ± 0.007	0.99 ± 0.03
NGC 5986		-1.35	19.25 ± 0.08	18.75 ± 0.01	20.30 ± 0.02	0.447 ± 0.003	0.727 ± 0.003	0.82 ± 0.02
NGC 6093	M 80	-1.47	18.90 ± 0.06	18.53 ± 0.02	19.38 ± 0.02	0.579 ± 0.005	0.774 ± 0.004	0.85 ± 0.02
NGC 6171	M 107	-0.95	18.14 ± 0.07	17.74 ± 0.02	18.92 ± 0.04	0.650 ± 0.004	0.944 ± 0.008	1.02 ± 0.04
NGC 6205	M 13	-1.33	18.33 ± 0.07	17.89 ± 0.01	19.07 ± 0.03	0.471 ± 0.003	0.719 ± 0.003	—
NGC 6266	M 62	-1.02	18.21 ± 0.07	17.77 ± 0.02	19.15 ± 0.02	0.522 ± 0.002	0.805 ± 0.003	0.89 ± 0.03
NGC 6273	M 19	-1.53	18.71 ± 0.07	18.16 ± 0.02	19.70 ± 0.02	0.465 ± 0.003	0.729 ± 0.004	—
NGC 6284		-1.17	19.92 ± 0.07	19.48 ± 0.02	20.71 ± 0.02	0.502 ± 0.003	0.774 ± 0.005	0.88 ± 0.02
NGC 6287		-1.90	18.58 ± 0.08	18.10 ± 0.04	19.33 ± 0.03	0.533 ± 0.006	0.776 ± 0.009	0.78 ± 0.05
NGC 6293		-1.73	18.62 ± 0.07	18.12 ± 0.02	19.54 ± 0.03	0.365 ± 0.002	0.631 ± 0.006	—
NGC 6304		-0.68	18.24 ± 0.06	17.91 ± 0.01	18.98 ± 0.02	0.542 ± 0.004	0.898 ± 0.005	0.93 ± 0.03
NGC 6356		-0.69	20.26 ± 0.09	19.88 ± 0.01	20.96 ± 0.02	0.547 ± 0.003	0.875 ± 0.003	0.94 ± 0.04
NGC 6362		-0.96	18.50 ± 0.06	18.12 ± 0.02	19.30 ± 0.04	0.488 ± 0.002	0.794 ± 0.007	0.88 ± 0.02
NGC 6522		-1.21	18.94 ± 0.09	18.27 ± 0.04	19.63 ± 0.02	0.599 ± 0.005	0.820 ± 0.020	0.90 ± 0.20
NGC 6544		-1.20	16.21 ± 0.11	15.76 ± 0.02	17.22 ± 0.03	0.535 ± 0.009	0.830 ± 0.020	—
NGC 6584		-1.30	19.49 ± 0.07	19.09 ± 0.02	20.34 ± 0.03	0.420 ± 0.002	0.705 ± 0.006	—
NGC 6624		-0.70	18.73 ± 0.06	18.45 ± 0.02	19.47 ± 0.04	0.571 ± 0.003	0.901 ± 0.005	0.94 ± 0.03
NGC 6637	M 69	-0.78	19.02 ± 0.08	18.65 ± 0.01	19.86 ± 0.02	0.541 ± 0.003	0.867 ± 0.004	0.93 ± 0.03
NGC 6638		-0.73	19.06 ± 0.07	18.67 ± 0.02	20.04 ± 0.02	0.532 ± 0.004	0.810 ± 0.010	0.88 ± 0.05
NGC 6642		-1.08	18.64 ± 0.09	18.33 ± 0.03	19.32 ± 0.02	0.524 ± 0.008	0.760 ± 0.040	0.90 ± 0.20
NGC 6652		-0.81	19.17 ± 0.07	18.87 ± 0.02	19.95 ± 0.03	0.535 ± 0.004	0.876 ± 0.005	0.92 ± 0.02
NGC 6681	M 70	-1.35	18.93 ± 0.08	18.46 ± 0.02	19.83 ± 0.02	0.455 ± 0.002	0.736 ± 0.004	0.82 ± 0.02
NGC 6712		-0.94	18.35 ± 0.08	17.93 ± 0.02	19.08 ± 0.03	0.494 ± 0.004	0.818 ± 0.006	0.90 ± 0.05
NGC 6723		-0.96	18.73 ± 0.08	18.37 ± 0.01	19.59 ± 0.03	0.513 ± 0.002	0.813 ± 0.003	0.87 ± 0.02
NGC 6838	M 71	-0.73	17.22 ± 0.08	16.86 ± 0.02	17.87 ± 0.03	0.521 ± 0.007	0.870 ± 0.010	—
NGC 6864	M 75	-1.10	20.40 ± 0.09	20.07 ± 0.02	21.11 ± 0.02	0.479 ± 0.003	0.792 ± 0.003	0.87 ± 0.03
NGC 6934		-1.30	20.07 ± 0.06	19.51 ± 0.02	20.95 ± 0.03	0.424 ± 0.002	0.717 ± 0.005	0.81 ± 0.02
NGC 6981	M 72	-1.21	20.01 ± 0.07	19.65 ± 0.02	20.86 ± 0.03	0.414 ± 0.003	0.704 ± 0.004	0.77 ± 0.02
NGC 7078	M 15	-2.02	18.92 ± 0.07	18.39 ± 0.01	19.85 ± 0.02	0.396 ± 0.002	0.647 ± 0.002	0.76 ± 0.02
NGC 7089	M 2	-1.31	19.30 ± 0.07	18.77 ± 0.01	20.22 ± 0.02	0.387 ± 0.003	0.666 ± 0.004	—
NGC 7099	M 30	-1.92	18.64 ± 0.08	18.07 ± 0.02	19.50 ± 0.04	0.407 ± 0.002	0.672 ± 0.004	—

**Table 7.** As Table 6, but displaying horizontal branch, RGB bump and tip brightness as well as the cluster distances inferred from  $V_{\text{ZAHB}}$  and  $V_{\text{tip}}$ .  $(B - V)_{\text{ZAHB}}$  is the colour of the point on the ZAHB closest to the TO-colour

Name	$V_{\text{ZAHB}}$	$(m - M)_{\text{ZAHB}}$	$V_{\text{tip}}$	$(m - M)_{\text{tip}}$	$V_{\text{Bump}}$	$(B - V)_{\text{ZAHB}}$
NGC 104	13.94 ± 0.06	13.24 ± 0.03	11.39 ± 0.04	13.6 ± 0.1	14.30	0.626
NGC 362	15.34 ± 0.05	14.66 ± 0.03	12.19 ± 0.09	14.8 ± 0.1	15.26	0.383
NGC 1261	16.80 ± 0.20	16.09 ± 0.03	13.80 ± 0.10	16.5 ± 0.1	16.62	0.451
NGC 1851	16.18 ± 0.05	15.48 ± 0.02	13.16 ± 0.05	15.7 ± 0.1	16.05	0.459
NGC 1904	16.16 ± 0.05	15.55 ± 0.04	13.20 ± 0.10	16.0 ± 0.1	15.24	0.249
NGC 2808	15.63 ± 0.05	15.03 ± 0.03	12.67 ± 0.02	15.3 ± 0.1	15.60	0.396
NGC 3201	14.22 ± 0.09	13.66 ± 0.04	13.00 ± 0.10	15.7 ± 0.1	15.72	0.165
NGC 4147	17.06 ± 0.05	16.39 ± 0.03	13.80 ± 0.40	16.7 ± 0.4	18.11	0.401
NGC 4590	15.70 ± 0.10	15.25 ± 0.04	12.40 ± 0.50	15.4 ± 0.4	15.07	0.107
NGC 4833	14.52 ± 0.05	13.95 ± 0.04	12.20 ± 0.10	15.2 ± 0.1	14.18	0.220
NGC 5024	16.70 ± 0.05	16.30 ± 0.01	13.67 ± 0.08	16.7 ± 0.1	—	0.389
NGC 5634	17.70 ± 0.10	17.15 ± 0.04	14.10 ± 0.20	17.1 ± 0.2	17.28	0.100
NGC 5694	18.16 ± 0.05	17.89 ± 0.03	15.25 ± 0.08	18.2 ± 0.1	—	0.424
NGC 5824	18.28 ± 0.05	17.59 ± 0.01	14.90 ± 0.04	17.8 ± 0.1	17.80	0.395
NGC 5904	15.02 ± 0.05	14.35 ± 0.03	12.22 ± 0.08	14.9 ± 0.1	14.90	0.431
NGC 5927	15.28 ± 0.08	15.00 ± 0.03	13.24 ± 0.05	15.2 ± 0.2	15.90	0.856
NGC 5986	15.82 ± 0.05	15.26 ± 0.03	12.59 ± 0.08	15.4 ± 0.1	15.58	0.447
NGC 6093	15.49 ± 0.05	15.30 ± 0.03	12.55 ± 0.07	15.4 ± 0.1	15.40	0.579
NGC 6171	14.95 ± 0.05	14.34 ± 0.03	12.30 ± 0.20	14.8 ± 0.2	14.83	0.650
NGC 6205	14.90 ± 0.07	14.38 ± 0.03	11.80 ± 0.10	14.6 ± 0.1	—	0.125
NGC 6266	14.81 ± 0.05	14.18 ± 0.02	12.03 ± 0.04	14.6 ± 0.1	14.82	0.525
NGC 6273	14.89 ± 0.05	14.74 ± 0.03	12.04 ± 0.05	14.9 ± 0.1	—	0.465
NGC 6284	16.52 ± 0.05	15.99 ± 0.03	13.60 ± 0.10	16.3 ± 0.1	16.30	0.502
NGC 6287	15.02 ± 0.05	15.00 ± 0.02	12.40 ± 0.20	15.4 ± 0.2	15.98	0.533
NGC 6293	15.63 ± 0.05	14.85 ± 0.04	12.20 ± 0.10	15.2 ± 0.1	—	0.100
NGC 6304	14.69 ± 0.08	14.22 ± 0.01	12.81 ± 0.08	14.9 ± 0.2	15.40	0.778
NGC 6356	16.60 ± 0.20	15.99 ± 0.02	14.07 ± 0.05	16.2 ± 0.2	17.19	0.585
NGC 6362	15.14 ± 0.05	14.20 ± 0.02	12.60 ± 0.30	15.0 ± 0.3	15.23	0.488
NGC 6522	15.14 ± 0.05	14.63 ± 0.03	12.09 ± 0.09	14.8 ± 0.1	15.63	0.599
NGC 6544	13.21 ± 0.07	12.33 ± 0.04	11.00 ± 0.20	13.7 ± 0.2	—	0.100
NGC 6584	16.29 ± 0.05	15.61 ± 0.03	13.30 ± 0.20	16.1 ± 0.2	—	0.420
NGC 6624	15.32 ± 0.05	14.74 ± 0.02	12.99 ± 0.08	15.1 ± 0.2	15.83	0.700
NGC 6637	15.52 ± 0.07	14.82 ± 0.03	13.03 ± 0.07	15.3 ± 0.1	15.90	0.726
NGC 6638	15.69 ± 0.05	14.91 ± 0.01	13.09 ± 0.06	15.5 ± 0.1	15.89	0.532
NGC 6642	15.34 ± 0.15	14.72 ± 0.03	12.70 ± 0.10	15.3 ± 0.1	15.39	0.524
NGC 6652	15.78 ± 0.08	15.14 ± 0.04	13.30 ± 0.10	15.5 ± 0.2	16.15	0.750
NGC 6681	15.47 ± 0.07	14.95 ± 0.02	12.60 ± 0.20	15.4 ± 0.2	15.37	0.175
NGC 6712	14.90 ± 0.10	14.41 ± 0.03	13.18 ± 0.09	15.7 ± 0.1	15.11	0.545
NGC 6723	15.47 ± 0.05	14.91 ± 0.03	13.20 ± 0.10	15.7 ± 0.1	15.50	0.513
NGC 6838	13.73 ± 0.07	13.33 ± 0.04	12.30 ± 0.30	14.5 ± 0.3	—	0.697
NGC 6864	17.17 ± 0.05	16.65 ± 0.02	14.35 ± 0.05	17.0 ± 0.1	17.19	0.479
NGC 6934	16.81 ± 0.05	16.24 ± 0.04	13.50 ± 0.10	16.3 ± 0.1	16.44	0.155
NGC 6981	16.81 ± 0.05	16.03 ± 0.02	13.80 ± 0.20	16.5 ± 0.2	16.57	0.414
NGC 7078	15.86 ± 0.05	15.32 ± 0.03	12.42 ± 0.07	15.5 ± 0.8	14.97	0.396
NGC 7089	15.77 ± 0.05	15.05 ± 0.02	13.06 ± 0.08	15.8 ± 0.1	—	0.225
NGC 7099	14.99 ± 0.05	14.69 ± 0.04	12.80 ± 0.10	15.8 ± 0.1	—	0.336



**Table 8.** Polynomial relations for points along our isochrones used as age indicators. The relations are all of the form  $f(t) = a_0 + a_1t + a_2t^2 + a_3t^3$ , where  $t$  is in Gyr. Part I:  $V$ -magnitudes

[Fe/H]	$a_0$	$a_1$	$a_2$	$a_3$
$V_{TO}$				
-2.26	1.081588	0.409828	-0.019137	0.000357
-2.08	1.108813	0.416247	-0.019509	0.000357
-1.78	1.100272	0.458946	-0.023650	0.000468
-1.56	1.155703	0.478126	-0.026089	0.000546
-1.25	1.217895	0.527360	-0.032193	0.000736
-1.07	1.128014	0.589114	-0.038899	0.000941
-0.77	2.053031	0.376031	-0.021472	0.000476
-0.54	2.557615	0.261690	-0.012317	0.000242
$V_{TO}^b$				
-2.26	0.539520	0.468912	-0.027354	0.000614
-2.08	0.603893	0.467406	-0.027372	0.000616
-1.78	0.785738	0.448451	-0.026057	0.000585
-1.56	0.941462	0.428725	-0.024491	0.000544
-1.25	1.324353	0.354857	-0.018317	0.000382
-1.07	1.428730	0.346346	-0.017664	0.000366
-0.77	1.515883	0.353536	-0.018211	0.000382
-0.54	1.657861	0.338788	-0.016849	0.000346
$V_{TO}^t$				
-2.26	1.058614	0.571462	-0.026232	0.000411
-2.08	1.058448	0.603465	-0.030062	0.000525
-1.78	1.016492	0.686102	-0.039767	0.000821
-1.56	0.989632	0.755390	-0.048332	0.001092
-1.25	1.391756	0.737361	-0.051244	0.001246
-1.07	1.928974	0.632944	-0.044782	0.001121
-0.77	3.212045	0.311654	-0.018641	0.000434
-0.54	3.506210	0.238451	-0.012321	0.000258
$V_{ZAHB}$				
-2.26	0.540172	-0.039658	0.002869	-0.000080
-2.08	0.532918	-0.019872	0.001615	-0.000046
-1.78	0.526569	0.002118	0.000071	-0.000007
-1.56	0.532144	0.007506	-0.000312	0.000003
-1.25	0.586536	0.010611	-0.000563	0.000009
-1.07	0.633460	0.010605	-0.000585	0.000010
-0.77	0.718433	0.008860	-0.000476	0.000007
-0.54	0.776018	0.008516	-0.000483	0.000008
$V_{Bump}$				
-2.26	-1.642720	0.178577	-0.008972	0.000172
-2.08	-1.650644	0.214845	-0.012553	0.000283
-1.78	-1.349552	0.186709	-0.009940	0.000213
-1.56	-1.232649	0.204869	-0.011501	0.000252
-1.25	-0.825191	0.163242	-0.007843	0.000159
-1.07	-0.614190	0.161057	-0.008018	0.000167
-0.77	-0.324075	0.157983	-0.007189	0.000135
-0.54	0.149489	0.098844	-0.002696	0.000030
$V_{tip}$				
-2.26	-3.115167	0.011530	-0.000371	0.000003
-2.08	-3.144452	0.019023	-0.000891	0.000016
-1.78	-3.153072	0.028388	-0.001289	0.000021
-1.56	-3.134376	0.039542	-0.001889	0.000033
-1.25	-3.065595	0.057298	-0.002933	0.000057
-1.07	-2.983806	0.065208	-0.003290	0.000065
-0.77	-2.835034	0.099049	-0.004695	0.000090
-0.54	-2.477340	0.118030	-0.005648	0.000109

**Table 9.** As Table 8. Part II: Colours

[Fe/H]	$a_0$	$a_1$	$a_2$	$a_3$
$(B - V)_{TO}$				
-2.26	-0.191148	0.103417	-0.006435	0.000145
-2.08	-0.168337	0.101817	-0.006405	0.000146
-1.78	-0.111845	0.097322	-0.006293	0.000147
-1.56	-0.059863	0.092706	-0.006109	0.000147
-1.25	0.052195	0.076287	-0.004949	0.000120
-1.07	0.133400	0.062945	-0.003886	0.000094
-0.77	0.269861	0.040226	-0.001963	0.000043
-0.54	0.329375	0.036700	-0.001614	0.000034
$(B - V)_{RGB}$				
-2.26	0.753462	-0.019023	0.001069	-0.000021
-2.08	0.771068	-0.019883	0.001132	-0.000023
-1.78	0.809573	-0.023631	0.001472	-0.000031
-1.56	0.841130	-0.025759	0.001680	-0.000037
-1.25	0.899441	-0.030508	0.002189	-0.000052
-1.07	0.974050	-0.041904	0.003286	-0.000084
-0.77	0.977762	-0.021655	0.001561	-0.000037
-0.54	1.016195	-0.011496	0.000721	-0.000015
$(B - V)_{Bump}$				
-2.26	0.799812	-0.001122	0.000000	0.000002
-2.08	0.816658	-0.002806	0.000184	-0.000004
-1.78	0.832050	-0.000746	0.000032	-0.000001
-1.56	0.856822	-0.001531	0.000096	-0.000001
-1.25	0.881010	0.003478	-0.000307	0.000009
-1.07	0.921353	0.000461	0.000038	-0.000002
-0.77	0.979223	0.003045	-0.000231	0.000007
-0.54	1.026973	0.005509	-0.000360	0.000108