

Improved correlation between the variability and peak luminosity of gamma-ray bursts

Li-Xin Li¹★ and Bohdan Paczyński²★

¹Max-Planck-Institut für Astrophysik, 85741 Garching, Germany

²Princeton University Observatory, Princeton, NJ 08544, USA

Accepted 2005 November 8. Received 2005 October 27; in original form 2005 September 26

ABSTRACT

A new procedure for smoothing a gamma-ray burst (GRB) light curve and calculating its variability is presented. Applying the procedure to a sample of 25 long GRBs, we have obtained a very tight correlation between the variability and the peak luminosity. The only significant outlier in the sample is GRB 030329. With this outlier excluded, the data scatter is reduced by a factor of ~ 3 compared to that of Guidorzi et al., measured by the deviation of fit. Possible causes for the outlier are discussed.

Key words: methods: data analysis – gamma-rays: bursts.

1 INTRODUCTION

Despite exciting progress in observations, the nature of gamma-ray bursts (GRBs) remains a big puzzle [see van Paradijs, Kouveliotou & Wijers (2000) and Piran (2004) for recent reviews]. In such a situation, it is extremely important to identify some good correlations between the apparent (easy to measure or calculate) and intrinsic properties of GRBs. Several such correlations have indeed been found. For example, an anticorrelation between the peak luminosity and spectral lag of GRBs has been found by Norris, Marani & Bonnell (2000), and a correlation between the peak luminosity and the variability of GRB light curves has been found by Fenimore & Ramirez-Ruiz (2000, hereafter FR00) and Reichart et al. (2001, hereafter R01). A correlation between the total isotropic energy and the peak energy of the spectrum (Amati et al. 2002), or the collimation-corrected total energy and the peak energy of the spectrum (Ghirlanda, Ghisellini & Lazzati 2004), has also been discovered.

Recently, Guidorzi et al. (2005, hereafter G05) tested the correlation between the variability and peak luminosity of GRBs, using an expanded sample of 32 GRBs with measured redshifts. The definitions of the variability and the peak luminosity are the same, but the size of the GRB sample of G05 is about three times bigger than that of R01. The existence of a correlation was confirmed, but the scatter in the correlation is significantly larger than that found by R01 (see, however, Reichart & Nysewander 2005). Although the issue is in debate (Guidorzi 2005; Reichart 2005; Reichart & Nysewander 2005), it is clear that with the definition of variability given by R01, the correlation between the variability and the peak luminosity is not tight.

In this paper, we present a new definition for the variability of GRB light curves. We then apply it to a sample of 25 long-duration

GRBs with measured redshifts, whose data are publicly available. We show that, with the new definition of the variability, the correlation between the variability and the peak luminosity of GRBs is significantly improved: the data scatter is considerably reduced.

2 NEW DEFINITION OF THE VARIABILITY

To measure the variability of a light curve, first we must define a *reference light curve*. The reference light curve should be sufficiently smoother than the original raw light curve. Since an ultimate model for GRBs does not exist yet, there is no first principle to guide us in choosing a reference light curve. What people usually do is to smooth the raw light curve with a linear ‘boxcar’ filter (or moving window), which smoothes the light curve with linear average (FR00; R01).

Here we use a *Savitzky–Golay filter* (Press et al. 2002) to smooth a light curve. The Savitzky–Golay filter is a more general and more powerful approach for smoothing noisy data than the linear boxcar filter. The basic idea of Savitzky–Golay filtering is to approximate the underlying function (i.e. the reference light curve) within the moving window by a polynomial of higher order. An advantage of the Savitzky–Golay filter to the linear filter is that the former preserves high moments while the latter does not.

A Savitzky–Golay filter is specified by three numbers: the order of the polynomial (m), the number of points used to the left of a data point (n_L) and the number of points used to the right of a data point (n_R). To apply the Savitzky–Golay filter, the data must be binned with constant spacing. For more details see Press et al. (2002).

We use a third-order Savitzky–Golay filter. That is, we set $m = 3$. The width of the filter, that is, $n_P \equiv n_L + n_R + 1$, is determined with the approach of R01: n_P is set to be equal to the number of data points corresponding to a time-scale T_f – the time spanned by the brightest 100 f per cent of the total counts above the background (for details see R01). It turns out that $f = 0.5$ most suits our

★E-mail: lxl@mpa-garching.mpg.de (L-XL); bp@astro.princeton.edu (BP)

purpose (rather than the $f = 0.45$ used by R01 and G05; see Section 5). So, throughout this paper we use n_p determined by $T_{0.5}$.

We then define $n_L = \text{int}[(n_p - 1)/2]$, and $n_R = n_p - n_L - 1$. If n_p is odd, we have $n_R = n_L$, that is, the filter is symmetric about the point to smooth. If n_p is even, we have $n_R = n_L + 1$, the filter is asymmetric. In this case, we smooth the light curve twice: first we use the n_L and n_R defined above, then we switch n_L and n_R . The results are then averaged.

Suppose we have obtained a reference light curve by applying the Savitzky–Golay filter to the raw light curve. Let us denote the count of the raw data in the i th time bin by C_i , the count given by the reference light curve by Y_i . The total squared deviation of the raw light curve from the reference light curve is then $\sum_{i=1}^{N_{\text{bin}}} (C_i - Y_i)^2$, where N_{bin} is the total number of bins to be summed. We obtain the intrinsic squared deviation by subtracting the Poisson noise N_{Poisson}

$$\Delta C^2 = W \sum_{i=1}^{N_{\text{bin}}} (C_i - Y_i)^2 - N_{\text{Poisson}}, \quad (1)$$

where $W \equiv n_p/(n_p - m - 1)$ is a statistical weight accounting for the fact that among the n_p data points only $n_p - m - 1$ are statistically independent. The inclusion of W allows us to apply the variability definition to any light curve with $n_p > m + 1$.

The summation in equation (1) is from time t_1 (corresponding to $i = 1$) to time t_2 (corresponding to $i = N_{\text{bin}}$), enclosing a major part of the light curve. Following FR00 and R01, we define t_1 to be the start of T_{90} , t_2 to be the end of T_{90} , where T_{90} is the time during which the cumulative counts of the GRB increase from 5 to 95 per cent above background (Kouveliotou et al. 1993).

The Poisson noise is calculated by

$$N_{\text{Poisson}} = \sum_{i=1}^{N_{\text{bin}}} [C_i + (\xi - 1)C_{\text{bg},i}], \quad (2)$$

where $C_{\text{bg},i}$ is the background, the factor ξ is the ratio of the background fluctuation to the Poisson noise of the background (given by the reduced χ^2 of the background fit). For GRBs detected by *Swift* we found that ξ is often significantly larger than unity, indicating that the background fluctuation is quite non-Poissonian.

The *variability* of the light curve is defined by the normalized squared deviation. We find that the following definition leads to the tightest correlation between the variability and the peak luminosity

$$V = \frac{\Delta C^2}{(N_{\text{bin}} - 1) C_{\text{max}}^2}, \quad (3)$$

where $\Delta C^2/(N_{\text{bin}} - 1)$ is the average of the squared deviation, C_{max} is the net peak count (i.e. the background is subtracted).

Our variability is defined in the observer’s frame, so that the information of GRB redshift is not needed. This not only makes the computation of variability simple, but also eliminates the uncertainty arising from the assumption about the dependence of light-curve variability on photon energy. The effect of GRB redshift was considered by R01 and FR00 who defined their variabilities in the GRB frame, but the dependence of variability on redshift turned out to be very weak.

As we applied our smoothing procedure to the GRBs in the sample described in Section 3, we found that the tightest correlation between the variability and the peak luminosity is obtained if we iteratively apply the Savitzky–Golay filter N_{iter} times, where N_{iter} is the integer closest to T_{90}/T_f (i.e. N_{iter} is roughly the number of moving windows contained in T_{90}).

In summary, our definition of variability differs from that of FR00 and R01 in the following aspects: (i) Our variability is defined in

the observer’s frame, while the variabilities of FR00 and R01 are defined in the frame of the GRB. (ii) We normalize the average of the squared deviation by the squared peak count (the same as FR00), while R01 normalize the total squared deviation by the sum of squared counts. (iii) FR00 and R01 use a linear boxcar filter, while we use a non-linear Savitzky–Golay filter.

3 DESCRIPTION OF THE SAMPLE

Our GRB sample contains 19 GRBs from the sample in G05, and six more GRBs detected by French Gamma Telescope (FREGATE)/*HETE-2* and Burst Alert Telescope (BAT)/*Swift*. So, the total number of GRBs in our sample is 25. They are listed in Table 1, with measured redshift, calculated isotropic-equivalent peak luminosity, calculated variability, and the number of iterations in applying the Savitzky–Golay filter. We have chosen to use the GRBs with data available publicly, which include GRBs detected by Burst and Transient Source Experiment (BATSE)/*Compton Gamma Ray Observatory (CGRO)*, *HETE-2*, and BAT/*Swift*.¹ To obtain a reliable calculation of variability, we have only selected GRBs with more than 30 per cent of total counts above the 3σ of background. As a result, those GRBs with too low signal-to-noise ratios (S/Ns) are not included in our sample.

The 19 GRBs from G05 are: 970508, 971214, 980425, 980703, 990123, 990506, 990510, 991216, 000131, 010921, 020124, 020813, 030328, 030329, 041006, 050401, 050505, 050525 and 050603. Their peak luminosities are taken from the same paper. The rest 13 GRBs in G05 are not included in our sample for various reasons: either their data are not publicly available (notably the seven GRBs detected by *BeppoSAX* but not by BATSE or *HETE-2*), or their data are incomplete or have too low S/Ns. Although 050315 and 050319 in G05 were detected by BAT/*Swift*, their data were not available to us when this paper was written, because the archive of *Swift* only contained data taken after 2005 April 1.

The six newly added GRBs are 030115a, 030528, 050408 (from *HETE-2*), 050802, 050803 and 050820a (from *Swift*). Their peak luminosities are calculated in exactly the same way as in R01 and G05. For GRBs detected by *HETE-2*, we used the spectral fit provided by Sakamoto et al. (2005) and the *HETE* website. For GRBs detected by *Swift*, peak spectra are extracted from their event files, then fitted with a power law which is sufficient for all cases (in the energy range 15–350 keV).

Although Golenetskii et al. (2005), Pal’shin et al. (2005) and Cummings et al. (2005) have reported the detection of a second, larger episode of emission from GRB 050820a (by both *Konus/Wind* and BAT/*Swift*), only the data for the first episode from *Swift* are available to us. Thus, for GRB 050820a, the variability and peak luminosity listed in Table 1 refer to its first episode (of the duration of ~ 30 s).

The time bin of each GRB light curve is as follows. For GRBs detected by BATSE, except 000131, the time bin is 64 ms. GRB 000131 does not have a 64-ms light curve, so we use its 1.024-s light curve. Although a bin of 1.024 s is somewhat large, the duration of 000131 is about 100 s which is much larger than its time bin. For GRBs detected by *HETE-2*, the time bin is 164 ms. While for GRBs

¹ Although the light-curve data of some GRBs detected by *Konus/WIND* are available from <http://gcn.gsfc.nasa.gov/gcn/konus-grbs.html>, we have chosen not to use them since the data are highly incomplete in the part of pre- and post-burst.

Table 1. Variability versus peak luminosity for 25 GRBs with known redshift.

GRB	z^a	Mission ^b	N_{iter}^c	V^d	L^e	z references ^f
970508	0.835	B/BS/U/K	6	0.0018 ± 0.0009	9.43 ± 1.89	(1)
971214	3.418	B/BS/U/K/N/R	4	0.0106 ± 0.0018	360 ± 65	(2)
980425	0.0085	B/BS/U/K	6	0.00041 ± 0.00041	0.0007 ± 0.0002	(3)
980703	0.966	B/BS/U/K/R	4	0.0030 ± 0.0005	26.4 ± 5.6	(4)
990123	1.6004	B/BS/U/K	3	0.0100 ± 0.0017	840 ± 121	(5)
990506	1.3	B/BS/U/K/R	8	0.0083 ± 0.0004	583 ± 121	(6)
990510	1.619	B/BS/U/K/N	12	0.0064 ± 0.0003	300 ± 50	(7)
991216	1.02	B/BS/U/N	4	0.0132 ± 0.0004	1398 ± 200	(8)
000131	4.5	B/U/K/N	8	0.0113 ± 0.0007	3600 ± 900	(9)
010921	0.45	H/BS/U/K	3	0.0020 ± 0.0017	8.0 ± 2.0	(10)
020124	3.198	H/U/K	5	0.0097 ± 0.0045	300 ± 60	(11)
020813	1.25	H/U/K/O	5	0.0100 ± 0.0013	340 ± 70	(12)
030115a	2.2	H	12	0.0057 ± 0.0036	57.0 ± 8.0	(13)
030328	1.52	H/U/K	3	0.0043 ± 0.0015	90 ± 18	(14)
030329	0.168	H/U/K/O/RH	4	0.0070 ± 0.0010	6.1 ± 1.2	(15)
030528	0.782	H	4	0.0018 ± 0.0023	1.4 ± 0.5	(16)
041006	0.712	H/K/RH	6	0.0037 ± 0.0008	66 ± 10	(17)
050401	2.90	BSw	7	0.0120 ± 0.0022	740 ± 100	(18)
050408	1.2357	H	9	0.0072 ± 0.0033	148 ± 67	(19)
050505	4.27	BSw	8	0.0090 ± 0.0051	250 ± 50	(20)
050525	0.606	BSw	3	0.0071 ± 0.0012	80 ± 10	(21)
050603	2.821	BSw	6	0.0100 ± 0.0006	1200 ± 200	(22)
050802	1.71	BSw	3	0.0042 ± 0.0038	104 ± 23	(23)
050803	0.422	BSw	6	0.0026 ± 0.0009	2.4 ± 0.6	(24)
050820a	2.612	BSw	5	0.0058 ± 0.0038	63 ± 13	(25)

^aMeasured redshift. ^bMission: B (BATSE/CGRO), H (HETE-2), BS (BeppoSAX), K (Konus/WIND), BSw (BAT/Swift), U (Ulysses), S (ROSS-C), N (NEAR), R (RossiXTE), O (Mars Odyssey), RH (RHESSI): the data used are taken from the first mission mentioned.

^cNumber of iterations in applying the Savitzky–Golay filter (see Section 2). ^dCalculated variability (equation 3) and error. ^eIsotropic-equivalent peak luminosity in 10^{50} erg s^{-1} in the rest-frame 100–1000 keV band, for peak fluxes measured on a 1-s time-scale ($H_0 = 65$ km s^{-1} Mpc $^{-1}$, $\Omega_m = 0.3$, and $\Omega_\Lambda = 0.7$). ^fReferences for the redshift measurements: (1) Metzger et al. (1997), (2) Kulkarni et al. (1998), (3) Tinney et al. (1998), (4) Djorgovski et al. (1998), (5) Kulkarni et al. (1999), (6) Bloom et al. (2003), (7) Beuermann et al. (1999), (8) Vreeswijk et al. (1999), (9) Andersen et al. (2000), (10) Djorgovski et al. (2001), (11) Hjorth et al. (2003a), (12) Price et al. (2002), (13) <http://space.mit.edu/HETE/Bursts/Data/> and Smith et al. (2005), (14) Martini, Garnavich & Stanek (2003), (15) Greiner et al. (2003), (16) Rau, Salvato & Greiner (2005), (17) Fugazza et al. (2004), (18) Fynbo et al. (2005a), (19) Berger, Gladders & Oemler (2005a) and Prochaska et al. (2005a), (20) Berger et al. (2005b), (21) Foley et al. (2005), (22) Berger & Becker (2005), (23) Fynbo et al. (2005b), (24) Bloom et al. (2005), (25) Prochaska et al. (2005b) and Ledoux et al. (2005).

detected by BAT/Swift, we extract 64-ms light curves from their event files available in the public data archive.

As we analyse the GRB light curves we use the total counts in the energy range specified as follows: 25–300 keV for GRBs detected by BATSE, 30–400 keV for those detected by HETE-2 and 25–350 keV for those detected by BAT/Swift. We have tried to make the range of energy for GRBs detected by different instruments to have largest overlap.

Finally, we remark that for the GRBs detected by BATSE, the light-curve data in the pre- and post-burst sections were divided equally into 64-ms time bins from their original 1.024-s time bins. When we calculate the Poisson noise in those sections, we multiply the result by a factor of 0.0625 ($= 0.064/1.024$) to take into account the reduction in the Poisson noise arising from the change in time bin.

4 RESULTS

We have applied the smoothing procedure described in Section 2 to our GRB sample. The obtained variabilities and errors, as well as the number of iterations in applying the Savitzky–Golay filter, are listed in Table 1. The errors of variability are principally caused by photon noises. However, the uncertainties arising from changing N_{iter} (see Section 2) by ± 1 are also taken into account: we find the

changes in the variability for $N_{\text{iter}} \pm 1$ and add in quadrature the maximum to the statistical error.

The peak luminosity versus variability is shown in Fig. 1. Clearly, a very tight correlation between the two quantities exists, with only one prominent outlier: GRB 030329. Given the large error in its variability, it is unclear if GRB 980425 – which is famous for its smallest redshift, lowest peak luminosity and least total energy, and association with SN 1998bw (Galama et al. 1998) – is off trend as in other kind of correlations (e.g. the total energy–peak energy correlation). In fact, the error in variability is larger than or about the same value as the variability itself for 980425 and 030528 (see Table 1).

We have made a least- χ^2 linear fit to $\log L$ – $\log V$, where L is the peak luminosity in erg s^{-1} . To take into account both the errors in L and V , we have made use of the program FITEXY in Press et al. (2002). The asymmetric errors in $\log L$ and $\log V$ are averaged. That is, we take

$$\sigma_{\log q} = \frac{1}{2} [\log(q + \sigma_q) - \log(q - \sigma_q)], \quad (4)$$

where $q = L$ or V . Since the errors in $\log V$ are infinite for 980425 and 030528, these two GRBs do not contribute to the total χ^2 . GRB 030329 was not considered due to its large offset. With 980425, 030329 and 030528 excluded, the total number of GRBs in the final sample to fit is 22. We have obtained the following result (the solid




Figure 1. Peak luminosity versus variability of 25 GRBs. The solid straight line is the least- χ^2 linear fit to the data, with GRBs 980425, 030329 and 030528 excluded (see the text): $\log L = 3.25 \log V + 59.42$ (L in units of erg s^{-1}). The $\chi^2/\text{d.o.f.} = 38.58/20 = 1.93$. The two dashed lines mark the 1σ deviation of the fit. (Filled circles: GRBs detected by BATSE; squares: GRBs detected by *HETE-2*; triangles: GRBs detected by BAT/*Swift*.)

straight line in Fig. 1)

$$\log L = a \log V + b, \quad (5)$$

where

$$a = 3.25 \pm 0.26, \quad b = 59.42 \pm 0.53. \quad (6)$$

The reduced chi-square $\chi_r^2 \equiv \chi^2/\text{d.o.f.} = 1.93$, where the degrees of freedom (d.o.f.) = $22 - 2 = 20$. The smallness of χ_r^2 indicates a very tight correlation between the peak luminosity and variability, for details see Section 5.

The two dashed lines in Fig. 1 mark the 1σ width of the fit, measuring the scatter of data and defined by the deviation of fit (see Section 5). However, this should not be mixed with the error in the predicted $\log L$, which we discuss below.

In Fig. 2, we plot the confidence intervals in the a - b plane. The best-fitting values of a and b , which correspond to the minimum of χ^2 , are indicated by the cross sign in the figure. The three contours enclose the regions containing, respectively, 68.3, 90 and 95.4 per cent of normally distributed data. The highly elongated shape of the contours indicates that a and b are highly correlated.

For any measured value of $\log V$, equation (5) predicts a value of $\log L$. The error in the predicted $\log L$ comes from two sources: the error in $\log V$ and the uncertainties in a and b

$$\sigma_{\log L} = \sqrt{a^2 \sigma_{\log V}^2 + \sigma_{ab}^2}. \quad (7)$$

Generally, the errors are asymmetric in the upward and downward directions. Let us denote $\sigma_{ab,1}$ and $\sigma_{ab,2}$ as upward error and downward error, respectively, due to uncertainties in a and b . That is, the predicted log luminosity is $\log L_{-}^{+\sigma_{ab,1}}$, if $\sigma_{\log V} = 0$. In Fig. 3, we show the numerically calculated values of $\sigma_{ab,1}$ (filled circles) and $\sigma_{ab,2}$ (open circles) versus the predicted log luminosity. From




Figure 2. The confidence intervals in the a - b space. The cross sign denotes the values of a and b at the minimum of χ^2 ($a = 3.25$, $b = 59.42$ and $\chi^2 = 38.58$). The contours correspond to the increase in χ^2 , $\Delta\chi^2 = 2.30$, 4.61 and 6.17, respectively. The region enclosed by each contour contains 68.3, 90 and 95.4 per cent of normally distributed data.

Fig. 2, we find that $\delta b \sim 2.05 \delta a$ when δa and δb are not close to zero, which leads to $\sigma_{ab} \sim (\log V + 2.05) \delta a$. Thus, σ_{ab} is minimized at $\log V \approx -2.05$, that is, at $\log L \approx 52.8$, consistent with the numerical results.

For the convenience of application of the $\log L$ - $\log V$ relation (5) to the prediction of GRB's peak luminosity, we have attempted to approximate the numerical results for σ_{ab} by several analytic formulae. The results are summarized in Appendix A, and represented by the solid lines in Fig. 3.

Now we apply the above results to GRB 030329 to check how far it deviates from the $\log L$ - $\log V$ relation. From Table 1, for GRB 030329 we have $V = 0.0070$ and $\sigma_V = 0.0010$. Submitting $\log V = -2.15$ into equation (5), we obtain a predicted log




Figure 3. The error in the $\log L$ predicted by equation (5), due to the uncertainties in a and b . Filled circles are numerical solutions for the upward error ($\sigma_{ab,1}$). Open circles are numerical solutions for the downward error ($\sigma_{ab,2}$). Solid lines are analytical approximations to the numerical solutions given by equations (A1)–(A6).

luminosity for GRB 030329: $\log L = 52.4$. Since GRB 030329 lies below the straight line in Fig. 1, the relevant error in the predicted log luminosity is determined by $\sigma_{ab,2}$ and $\sigma_{\log V} = \log V - \log(V - \sigma_V) \approx 0.067$. By equation (A4) we have $\sigma_{ab,2} \approx 0.08$. Then, by equation (7), the total error in the predicted luminosity is $\sigma_{\log L} \approx 0.23$. From Table 1, the measured luminosity of GRB 030329 is $\log L \approx 50.8$, with an upward measured error $\sigma_{\log L} \approx 0.08$ that is smaller than the error in the predicted log luminosity. So, GRB 030329 deviates from the correlation (the straight line in Fig. 1) by $(52.4 - 50.8)/0.23 \approx 7\sigma$.

5 COMPARISON TO GUIDORZI ET AL.'S RESULTS

To make a quantitative comparison of our results to that of G05, we need a parameter that measures the scatter of data points around the best-fitting model. The parameter should not be sensitive to the variation in data errors caused by different definitions of variability. Reichart (2001) has provided such a parameter based on the maximum-likelihood method, which is called the ‘sample variance’ or ‘slop’. However, here we choose to use the ‘variance of fit’ defined with the approach of least- χ^2 (Bevington & Robinson 1992), which is much less sophisticated but enough for our current purpose.

Suppose a data set $\{x_i, y_i\}$, with errors $\{\sigma_{x,i}, \sigma_{y,i}\}$, is fitted by a model $y = f(x)$. The fit minimizes the χ^2 defined by

$$\chi^2 \equiv \sum_{i=1}^N \frac{1}{\sigma_i^2} [y_i - f(x_i)]^2, \quad (8)$$

where N is the total number of data points, and

$$\sigma_i^2 = \sigma_{y,i}^2 + \left(\frac{df}{dx}\right)_{x=x_i}^2 \sigma_{x,i}^2. \quad (9)$$

The reduced χ^2 , that is, the χ^2 per d.o.f., is

$$\chi_r^2 = \chi^2/\text{d.o.f.} = \chi^2/(N - p), \quad (10)$$

where p is the number of parameters in the function $f(x)$.

The variance of the fit, measured along the y -direction, is defined by (Bevington & Robinson 1992)

$$s_y^2 \equiv \frac{\langle \sigma_i^2 \rangle}{N - p} \sum_{i=1}^N \frac{1}{\sigma_i^2} [y_i - f(x_i)]^2, \quad (11)$$

where $\langle \sigma_i^2 \rangle$ is the weighted average of the individual data variance

$$\langle \sigma_i^2 \rangle \equiv \left(\frac{1}{N} \sum_{i=1}^N \frac{1}{\sigma_i^2} \right)^{-1}. \quad (12)$$

The *variance of fit* measures the spread of data around the model. We denote the variance of fit by s^2 (then s is the *deviation of fit*), to distinguish it from the *data variance* σ^2 which represents the error in the measurement of each data point.

The appearance of $\langle \sigma_i^2 \rangle$ in the numerator and σ_i^2 in the denominator in equation (11) shows that the variance of fit is not sensitive to the variation in data variance.

By the definition of χ_r^2 , equation (11) can be recast into a simpler form

$$s_y^2 = \langle \sigma_i^2 \rangle \chi_r^2. \quad (13)$$




Figure 4. Definition of the variance of fit. The thick solid curve is the fit $y = f(x)$. The y -component of the variance of fit, s_y^2 , is defined by equation (11). The two dashed curves, called ‘the 1σ width’ of $y = f(x)$, are defined by $y = f(x) \pm s_y$. The x -component of the variance of fit, s_x^2 , is related to s_y^2 by equation (14). The scatter of data points around $y = f(x)$ is measured by $s = s_x s_y / \sqrt{s_x^2 + s_y^2}$, the length of the straight line OC that is perpendicular to the straight line AB. [The straight line AB locally approximates the curve $y = f(x)$ when s_y is sufficiently small or the curvature of the curve is negligible.]

We define the variance of the fit measured along the x -direction by

$$s_x^2 = s_y^2 \left(\frac{df}{dx}\right)^{-2}, \quad (14)$$

where df/dx is the slope of the curve $y = f(x)$. Note, by definitions, s_y is always a constant, but s_x is a function of x unless $f(x)$ is a linear function of x .

We measure the scatter of data points around the curve $y = f(x)$ by the variance of fit along a direction locally perpendicular to the curve, which is (see Fig. 4)

$$s^2 = \frac{s_x^2 s_y^2}{s_x^2 + s_y^2} = \frac{s_y^2}{1 + (df/dx)^2}. \quad (15)$$

For the problem in this paper, we have $x = \log V$, $y = \log L$ and $f(x) = ax + b$. Thus, $df/dx = a$ is a constant, and $s_x = s_y/a$, $\sigma_i^2 = \sigma_{y,i}^2 + a^2 \sigma_{x,i}^2$. The *deviation of fit* is then

$$s = \frac{s_y}{\sqrt{1 + a^2}}. \quad (16)$$

With the above tools, we are ready to compare our results to that of G05 quantitatively. To make the comparison well defined, we analyse the overlapping sample which contains 19 GRBs (see Section 3). In Fig. 5, we plot the variability with our definition versus that with the R01’s definition (the latter has been copied from G05). The two variabilities are highly correlated, but the scatter is large. This large scatter is necessary for obtaining a better correlation between the peak luminosity and the variability of GRBs.

The largest scatter in Fig. 5 occurs at GRB 980425, for which our background fit leads to $T_{90} = 60.29$ s, in contrast to the 34.88 s reported in the BATSE web page. If we adopted $T_{90} =$




Figure 5. The variability with our definition versus the variability with definition of R01 for 19 GRBs. The straight line is defined by the equation ‘our variability = variability of R01 divided by 15’.

34.88 and $\xi = 1$ (ξ is the ratio of the background fluctuation to the Poisson noise; see Section 2), we would obtain $V = 0.00053 \pm 0.00057$ for 980425. This value of V is only slightly larger than that listed in Table 1, so 980425 remains to have the largest scatter in Fig. 5. With our definition of variability, GRB 980425 is much less offset from the variability–luminosity correlation.

Fig. 5 also shows that the values of the variability with our definition are systematically smaller than that with the definition of R01, as indicated by the straight line defined by ‘our variability = R01’s variability divided by 15’. This is mainly caused by the different normalization in the two definitions. In addition, our variability generally has a larger error bar than that of R01, caused by the fact that we normalize our variability by peak count, which suffers larger photon noise than total counts; and we have considered the error arising from changing N_{iter} by ± 1 .

Now, we make a linear fit to $\log L - \log V$ for the overlapping sample, with our definition of variability. For the reasons explained in Section 4, we exclude GRBs 980425 and 030329 during the fit. Thus, the total number of GRBs to fit is $N = 17$. We obtain $\log L = 3.10 \log V + 59.13$, and $\chi^2/\text{d.o.f.} = 2.43$ where d.o.f. = 15. The weighted average of the individual data variance is $\langle \sigma_i^2 \rangle = 0.02715$. Thus, for the fit with our definition of variability, we have $s_y = 0.257$, $s_x = 0.0829$ and $s = 0.079$.

Next, we make a linear fit to $\log L - \log V$ for the overlapping sample, with R01’s definition of variability. Again, GRBs 980425 and 030329 are excluded, to make the GRB members in the sample remain the same. We obtain $\log L = 1.77 \log V + 54.06$, and $\chi^2/\text{d.o.f.} = 19.89$. The weighted average of the individual data variance is $\langle \sigma_i^2 \rangle = 0.009240$. Thus, for the fit with R01’s definition of variability, we have $s_y = 0.429$, $s_x = 0.2417$ and $s = 0.211$.

The scatter of data points around the fitted $\log L - \log V$ is measured by s . Thus, according to the above numbers, for the overlapping sample the scatter in our $\log L - \log V$ relation is smaller than that of G05 by a factor of $0.211/0.079 = 2.7$, although the weighted average of the individual data error ($\equiv \langle \sigma_i^2 \rangle^{1/2}$) with our definition




Figure 6. Comparison of our results (filled circles and thick lines) to that of G05 (open circles and thin lines). Straight lines represent the least- χ^2 fit and the 1σ width. (Our variability has been multiplied by a factor of 15.)

of variability is larger than that with R01’s definition of variability by a factor of $(0.02715/0.009240)^{1/2} = 1.7$.

The above results, for the overlapping sample excluding 980425 and 030329, are summarized in Fig. 6, where filled circles and thick lines represent the results with our data, open circles and thin lines represent the results with G05’s data. The luminosities in the two set of data are the same, so each GRB is represented by a horizontal line in Fig. 6. However, the variabilities in the two set of data differ, so we see the offset along the horizontal direction of each pair of filled circle–open circle. The solid straight lines are least- χ^2 fits to each data set, and the dashed straight lines mark the ‘ 1σ width’ of the fits. We have set the unitary length of the horizontal and vertical axes to be the same, so that for each data set s is just the half-width of the region bounded by the two dashed lines that you see. The figure clearly shows that the scatter of our data set is much narrower than that of G05’s data set.

With the above defined scatter parameter s , we can also check how sensitive our results are to the parameters in our definition of variability. As mentioned in Section 2, we have chosen $T_f = T_{0.5}$, and N_{iter} to be the integer closest to the ratio T_{90}/T_f . With




Figure 7. Variation of the deviation of the linear fit to $\log L$ – $\log V$, with respect to the choice of N_{iter} . The parameter α , which the $T_{0.5}$ in the definition of N_{iter} is multiplied by, is set to be the same for all GRBs in the sample. The variabilities presented in this paper correspond to the case $\alpha = 1$.

this choice, for the GRB sample described in Section 3 (excluding 980425, 030329 and 030528), we obtained $\chi^2/\text{d.o.f.} = 1.93$ and $s = 0.078$ (Fig. 1, where the two dashed lines are obtained by shifting the solid line upward/downward by $s_y = 0.265$). If we follow R01 and G04 to choose $T_f = T_{0.45}$, we would obtain $\chi^2/\text{d.o.f.} = 2.53$ and $s = 0.087$ in the linear fit to $\log L$ – $\log V$. A slightly larger value of s indicates that the correlation is slightly looser.

To test how sensitive the results are to the choice of N_{iter} , let us redefine N_{iter} by the integer closest to $T_{90}/\alpha T_{0.5}$, where α is a parameter that is set to be the same for all GRBs in the sample. When $\alpha = 1$, we return to the definition that we have adopted. We now let α vary from 0.2 to 2, and calculate the corresponding s in the least- χ^2 fit to $\log L$ – $\log V$. The calculated s versus α are presented in Fig. 7, which shows that the results are stable to the choice of N_{iter} . That is, the variance of fit does not change drastically with a small change in α . The figure also shows that $\alpha = 1$ is around the minimum of $s(\alpha)$.

6 DISCUSSION AND CONCLUSIONS

We have presented a new definition for the variability of GRB light curves. The new variability is defined in the observer’s frame, and the light curve is smoothed with the Savitzky–Golay filter. The former makes it easy to apply the procedure to all GRBs with well-measured light curves (i.e. with a significant S/N, and a reasonable size of time bin – neither too large to erase the variability on short time-scales, nor too small to introduce too large Poisson noises), without referring to the redshift information.

We have applied our smoothing procedure to a sample of 25 long-duration GRBs with measured redshifts and publicly available data. A very tight correlation between the variability and the peak luminosity is found, with only one prominent outlier: GRB 030329. Excluding 030329, 980425 and 030528 (the latter two are due to

their too large errors in variability), a linear fit to the log of variability and the log of peak luminosity is obtained (equations 5 and 6), with $\chi^2/\text{d.o.f.} = 1.93$. The smallness of the reduced χ^2 indicates that the data scatter is very small (Figs 1 and 6).

We note that, although the existence of a second, larger episode of emission from GRB 050820a has been claimed (Cummings et al. 2005; Golenetskii et al. 2005; Pal’shin et al. 2005) and we have only made use of the data for its first episode, GRB 050820a fits the relation perfectly well.

Our results are a significant improvement to that of G05 (as well as that of FR00 and R01), who used a different definition of variability and obtained $\chi^2/\text{d.o.f.} = 1167/30 = 38.9$. An analysis on the overlapping sample of 17 GRBs (after GRBs 980425 and 030329 being excluded) shows that the scatter in our data is smaller than that in the G05’s data by a factor of 2.7, measured by the deviation of fit (Section 5).

The improvement to the correlation is caused by using a new smoothing procedure and a different normalization in the definition of our variability, not by neglecting the effect of GRB redshift. R01 have shown that the dependence of variability on the redshift is extremely weak. The fact that a very tight correlation between the peak luminosity and the variability is obtained without including the corrections to the variability from the GRB redshift also leads us to believe that such corrections are not necessary.

A remarkable feature of the correlation that we have found is that it does not rely on the corrections to the luminosity from the collimation of GRB jets. In other words, the peak luminosity used above is simply the isotropic-equivalent peak luminosity. If this correlation is confirmed by future bursts, it will provide a convenient calibration to GRB luminosity and distance.

GRB 030329, which is considered to be a firm case for the connection of GRBs with supernovae (Hjorth et al. 2003b; Stanek et al. 2003), is offset from the correlation by 7σ (Fig. 1). Although the cause is not clear, we have noted that 030329 and 050525 differ from other GRBs in the sample in the following way: their light curves consist of two distinctly separated pulses with each containing at least 30 per cent of total counts. Motivated by this observation and the fact that GRB 050820a fits the relation well although only the first episode of its emission has been used, we have recalculated the variabilities of 030329 and 050525 by dividing the smoothing window by a factor of 2 (i.e. dividing the n_p obtained from $T_{0.5}$ by 2) but keeping N_{iter} unchanged.² We obtained $V = 0.0022 \pm 0.0003$ for 030329, which well fits the solid straight line in Fig. 1. For 050525, we obtained $V = 0.0032 \pm 0.0007$, which fits the straight line equally well as the value listed in Table 1. Whether this treatment is correct must be tested when more GRBs with light curves similar to that of 030329 and 050525 are available.

ACKNOWLEDGMENTS

LXL thanks C. Guidorzi and T. Sakamoto for useful communications. The authors thank the anonymous referee for a very helpful report that has led to significant improvements to this paper. This research has made use of BATSE and BAT/*Swift* data obtained from the High-Energy Astrophysics Science Archive Research Center (HEASARC), provided by NASA Goddard Space Flight Center; and data obtained from the *HETE-2* science team via the website <http://space.mit.edu/HETE/Bursts>.

² Roughly speaking, this is equivalent to smooth each GRB pulse separately.

REFERENCES

- Amati et al., 2002, A&A, 390, 81
 Andersen M. I. et al., 2000, A&A, 364, L54
 Berger E., Becker G., 2005, GCN 3520
 Berger E., Gladders M., Oemler G., 2005a, GCN 3201
 Berger E., Bradley Cenko S., Steidel C., Reddy N., Fox D. B., 2005b, GCN 3368
 Beuermann K. et al., 1999, A&A, 352, L26
 Bevington P. R., Robinson D. K., 1992, Data Reduction and Error Analysis for the Physical Sciences. McGraw-Hill, New York
 Bloom J. S., Berger E., Kulkarni S. R., Djorgovski S. G., Frail D. A., 2003, AJ, 125, 999
 Bloom J. S., Perley D., Foley R., Prochaska J. X., Chen H. W., Starr D., 2005, GCN 3758
 Cummings J. et al., 2005, GCN 3858
 Djorgovski S. G., Kulkarni S. R., Bloom J. S., Goodrich R., Frail D. A., Piro L., Palazzi E., 1998, ApJ, 508, L17
 Djorgovski S. G. et al., 2001, GCN 1108
 Fenimore E. E., Ramirez-Ruiz E., 2000, preprint (astro-ph/0004176) (FR00)
 Foley R. J., Chen H.-W., Bloom J., Prochaska J. X., 2005, GCN 3483
 Fugazza D. et al., 2004, GCN 2782
 Fynbo J. P. U. et al., 2005a, GCN 3176
 Fynbo J. P. U. et al., 2005b, GCN 3749
 Galama T. J. et al., 1998, Nat, 395, 670
 Ghirlanda G., Ghisellini G., Lazzati D., 2004, ApJ, 616, 331
 Golenetskii S. et al., 2005, GCN 3846
 Greiner J. et al., 2003, GCN 2020
 Guidorzi C., 2005, MNRAS, 364, 163
 Guidorzi C., Frontera F., Montanari E., Rossi F., Amati L., Gomboc A., Hurley K., Mundell C. G., 2005, MNRAS, 363, 315 (G05)
 Hjorth J. et al., 2003a, ApJ, 597, 699
 Hjorth J. et al., 2003b, Nat, 423, 847
 Kouveliotou C., Meegan C. A., Fishman G. J., Bhat N. P., Briggs M. S., Koshut T. M., Paciesas W. S., Pendleton G. N., 1993, ApJ, 413, L101
 Kulkarni S. R. et al., 1998, Nat, 393, 35
 Kulkarni S. R. et al., 1999, Nat, 398, 389
 Ledoux et al., 2005, GCN 3860
 Martini P., Garnavich P., Stanek K. Z., 2003, GCN 1980
 Metzger M. R., Djorgovski S. G., Kulkarni S. R., Steidel C. C., Adelberger K. L., Frail D. A., Costa E., Frontera F., 1997, Nat, 387, 878
 Norris J. P., Marani G. F., Bonnell J. T., 2000, ApJ, 534, 248
 Pal'shin V., Frederiks D., 2005, GCN 3852
 Piran T., 2004, Rev. Mod. Phys., 76, 1143
 Press W. H., Teukolsky S. A., Vetterling W. T., Flannery B. P., 2002, Numerical Recipes in C++. Cambridge Univ. Press, Cambridge
 Price P. A., Bloom J. S., Goodrich R. W., Barth A. J., Cohen M. H., Fox D. W., 2002, GCN 1475
 Prochaska J. X., Bloom J. S., Chen H.-W., Foley R. J., Roth K., 2005a, GCN 3204
 Prochaska J. X., Bloom J. S., Wright J. T., Butler R. P., Chen H. W., Vogt S. S., Marcy G. W., 2005b, GCN 3833
 Rau A., Salvato M., Greiner J., 2005, preprint (astro-ph/0508394)
 Reichart D. E., 2001, ApJ, 553, 235
 Reichart D. E., 2005, preprint (astro-ph/0508529)
 Reichart D. E., Nysewander M. C., 2005, ApJ, submitted (astro-ph/0508111)
 Reichart D. E., Lamb D. Q., Fenimore E. E., Ramirez-Ruiz E., Cline T. L., Hurley K., 2001, ApJ, 552, 57 (R01)
 Sakamoto T. et al., 2005, ApJ, 629, 311
 Smith I. A. et al., 2005, A&A, 439, 987
 Stanek et al., 2003, ApJ, 591, L17
 Tinney C., Stathakis R., Cannon R., Galama T., 1998, IAU Circ. 6896
 van Paradijs J., Kouveliotou C., Wijers R. A. M. J., 2000, ARAA, 38, 379
 Vreeswijk P. M. et al., 1999, GCN 496

APPENDIX A: ANALYTIC APPROXIMATION TO THE ERROR IN THE LOG LUMINOSITY

The numerical results for the error in the $\log L$ predicted by equation (5) due to the uncertainties in a and b , σ_{ab} , can be approximated by the following formulae ($\sigma_{ab,1}$ for the upward error, $\sigma_{ab,2}$ for the downward error):

(1) When $\log L < 52$

$$\sigma_{ab,1} \approx 7.2017 - 0.16777x + 0.0006006x^2, \quad (\text{A1})$$

$$\sigma_{ab,2} \approx 8.0929 - 0.18000x + 0.0005134x^2, \quad (\text{A2})$$

where $x \equiv \log L$, L is in units of erg s^{-1} . (2) When $52 \leq \log L < 54$

$$\sigma_{ab,1} \approx 2274.034248 - 124.984716x + 2.286986x^2 - 0.013930421x^3, \quad (\text{A3})$$

$$\sigma_{ab,2} \approx 3028.186992 - 167.524125x + 3.087093x^2 - 0.018948436x^3. \quad (\text{A4})$$

(3) When $\log L \geq 54$

$$\sigma_{ab,1} \approx -6.8502 + 0.12989x, \quad (\text{A5})$$

$$\sigma_{ab,2} \approx -5.7594 + 0.10933x. \quad (\text{A6})$$

The fractional errors in σ_{ab} given by these formulae are always < 5 per cent.

This paper has been typeset from a \TeX/L\TeX file prepared by the author.