

The ATLAS^{3D} project – XX. Mass-size and mass- σ distributions of early-type galaxies: bulge fraction drives kinematics, mass-to-light ratio, molecular gas fraction and stellar initial mass function

Michele Cappellari,^{1*} Richard M. McDermid,² Katherine Alatalo,³ Leo Blitz,³
Maxime Bois,⁴ Frédéric Bournaud,⁵ M. Bureau,¹ Alison F. Crocker,⁶ Roger L. Davies,¹
Timothy A. Davis,^{1,7} P. T. de Zeeuw,^{7,8} Pierre-Alain Duc,⁵ Eric Emsellem,^{7,9}
Sadegh Khochfar,¹⁰ Davor Krajnović,⁷ Harald Kuntschner,⁷ Raffaella Morganti,^{11,12}
Thorsten Naab,¹³ Tom Oosterloo,^{11,12} Marc Sarzi,¹⁴ Nicholas Scott,^{1,15} Paolo Serra,¹¹
Anne-Marie Weijmans¹⁶ and Lisa M. Young¹⁷

¹ *Sub-department of Astrophysics, Department of Physics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford OX1 3RH*

² *Gemini Observatory, Northern Operations Centre, 670 N. A'ohoku Place, Hilo, HI 96720, USA*

³ *Department of Astronomy, Campbell Hall, University of California, Berkeley, CA 94720, USA*

⁴ *Observatoire de Paris, LERMA and CNRS, 61 Av. de l'Observatoire, F-75014 Paris, France*

⁵ *Laboratoire AIM Paris-Saclay, CEA/IRFU/SAp, CNRS, Université Paris Diderot, 91191 Gif-sur-Yvette Cedex, France*

⁶ *Department of Astrophysics, University of Massachusetts, 710 North Pleasant Street, Amherst, MA 01003, USA*

⁷ *European Southern Observatory, Karl-Schwarzschild-Str. 2, 85748 Garching, Germany*

⁸ *Sterrewacht Leiden, Leiden University, Postbus 9513, 2300 RA Leiden, the Netherlands*

⁹ *Université Lyon 1, Observatoire de Lyon, Centre de Recherche Astrophysique de Lyon and Ecole Normale Supérieure de Lyon, 9 avenue Charles André, F-69230 Saint-Genis Laval, France*

¹⁰ *Max-Planck Institut für extraterrestrische Physik, PO Box 1312, D-85478 Garching, Germany*

¹¹ *Netherlands Institute for Radio Astronomy (ASTRON), Postbus 2, 7990 AA Dwingeloo, The Netherlands*

¹² *Kapteyn Astronomical Institute, University of Groningen, Postbus 800, 9700 AV Groningen, The Netherlands*

¹³ *Max-Planck Institut für Astrophysik, Karl-Schwarzschild-Str. 1, 85741 Garching, Germany*

¹⁴ *Centre for Astrophysics Research, University of Hertfordshire, Hatfield, Herts AL1 9AB, UK*

¹⁵ *Centre for Astrophysics & Supercomputing, Swinburne University of Technology, PO Box 218, Hawthorn, VIC 3122, Australia*

¹⁶ *Dunlap Institute for Astronomy & Astrophysics, University of Toronto, 50 St. George Street, Toronto, ON M5S 3H4, Canada*

¹⁷ *Physics Department, New Mexico Institute of Mining and Technology, Socorro, NM 87801, USA*

Accepted 2013 April 15. Received 2013 March 24; in original form 2012 August 17

arXiv:1208.3523v2 [astro-ph.CO] 15 Apr 2013

ABSTRACT

In the companion Paper XV of this series we derive accurate total mass-to-light ratios $(M/L)_{\text{JAM}} \approx (M/L)(r = R_e)$ within a sphere of radius $r = R_e$ centred on the galaxy, as well as stellar $(M/L)_{\text{stars}}$ (with the dark matter removed) for the volume-limited and nearby mass selected (stellar mass $M_* \gtrsim 6 \times 10^9 M_\odot$) ATLAS^{3D} sample of 260 early-type galaxies (ETGs, ellipticals Es and lenticulars S0s). Here we use those parameters to study the two orthogonal projections $(M_{\text{JAM}}, \sigma_e)$ and $(M_{\text{JAM}}, R_e^{\text{maj}})$ of the thin Mass Plane (MP) $(M_{\text{JAM}}, \sigma_e, R_e^{\text{maj}})$ which describes the distribution of the galaxy population, where $M_{\text{JAM}} \equiv L \times (M/L)_{\text{JAM}} \approx M_*$. The distribution of galaxy properties on both projections of the MP is characterized by: (i) the same zone of exclusion (ZOE), which can be transformed from one projection to the other using the scalar virial equation. The ZOE is roughly described by two power-laws, joined by a break at a characteristic mass $M_{\text{JAM}} \approx 3 \times 10^{10} M_\odot$, which corresponds to the minimum R_e and maximum stellar density. This results in a break in the mean $M_{\text{JAM}} - \sigma_e$ relation with trends $M_{\text{JAM}} \propto \sigma_e^{2.3}$ and $M_{\text{JAM}} \propto \sigma_e^{4.7}$ at small and large σ_e respectively; (ii) a characteristic mass $M_{\text{JAM}} \approx 2 \times 10^{11} M_\odot$ which separates a population dominated by flat fast rotator with disks and spiral galaxies at lower masses, from one dominated by quite round slow rotators at larger masses; (iii) below that mass the distribution of ETGs properties on the two projections of the MP tends to be constant along lines of roughly constant σ_e , or equivalently along lines with $R_e^{\text{maj}} \propto M_{\text{JAM}}$ respectively (or even better parallel to the ZOE: $R_e^{\text{maj}} \propto M_{\text{JAM}}^{0.75}$); (iv) it forms a continuous and parallel sequence with the distribution of spiral galaxies; (v) at even lower masses, the distribution of fast rotator ETGs and late spirals naturally extends to that of dwarf ETGs (Sph) and dwarf irregulars (Im) respectively.

We use dynamical models to analyse our kinematic maps. We show that σ_e traces the bulge fraction, which appears the main driver for the observed trends in the dynamical $(M/L)_{\text{JAM}}$ and in indicators of the $(M/L)_{\text{pop}}$ of the stellar population like $H\beta$ and colour, as well as in the molecular gas fraction. A similar variation along contours of σ_e is also observed for the mass normalization of the stellar Initial Mass Function (IMF), which was recently shown to vary systematically within the ETGs galaxy population. Our preferred relation has the form $\log_{10}[(M/L)_{\text{stars}}/(M/L)_{\text{Salp}}] = a + b \times \log_{10}(\sigma_e/130 \text{ km s}^{-1})$ with $a = -0.12 \pm 0.01$ and $b = 0.35 \pm 0.06$. Unless there are major flaws in all stellar population models, this trend implies a transition of the mean IMF from Kroupa to Salpeter in the interval $\log_{10}(\sigma_e/\text{km s}^{-1}) \approx 1.9 - 2.5$ (or $\sigma_e \approx 90 - 290 \text{ km s}^{-1}$), with a smooth variation in between, consistently with what was shown in Cappellari et al. (2012). The observed distribution of galaxy properties on the MP provides a clean and novel view for a number of previously reported trends, which constitute special two-dimensional projections of the more general four-dimensional parameters trends on the MP. We interpret it as due to a combination of two main effects: (i) an increase of the bulge fraction, which increases σ_e and decrease in R_e , and greatly enhances the likelihood for a galaxy to have its star formation quenched, and (ii) dry merging, increasing galaxy mass and R_e by moving galaxies along lines of roughly constant σ_e (or steeper), while leaving the population nearly unchanged.

Key words: galaxies: elliptical and lenticular, cD – galaxies: evolution – galaxies: formation – galaxies: structure – galaxies: kinematics and dynamics

1 INTRODUCTION

Much of our understanding of galaxy formation and evolution comes from the study of dynamical scaling relations relating galaxy luminosity or mass, size and kinematic (e.g. Faber & Jackson 1976; Kormendy 1977; Dressler et al. 1987; Faber et al. 1987; Djorgovski & Davis 1987) or regular trends in the distribution of galaxy properties as a function of their scaling parameters (e.g. Bender, Burstein & Faber 1992; Burstein et al. 1997; Kauffmann et al. 2003b; Galazzi et al. 2006), and from the study of their evolution with redshift (e.g. van Dokkum & Franx 1996; Kelson et al. 1997; van Dokkum et al. 1998; Treu et al. 2005; Franx et al. 2008).

The volume-limited ATLAS^{3D} sample of nearby early-type galaxies (Cappellari et al. 2011a, hereafter Paper I) constitute an ideal benchmark for studying scaling relations and the distribution of galaxy properties, given the availability of a high-quality multi-wavelength dataset. In particular in Cappellari et al. (2013, hereafter Paper XV) we used the state-of-the-art integral-field kinematics, in combination with detailed axisymmetric dynamical models, to derive accurate masses and global dynamical parameter. We found that galaxies lie, with very good accuracy, on a thin Mass Plane (MP) describing galaxies in the parameter space defined by mass, velocity dispersion and projected half-light radius $(M_{\text{JAM}}, \sigma_e, R_e^{\text{maj}})$. Here M_{JAM} is a very good estimate of the galaxy total stellar mass, R_e^{maj} is the major axis of the ‘effective’ isophote containing half of the observed galaxy light and σ_e is the

* E-mail: cappellari@astro.ox.ac.uk

velocity dispersion measured within that isophote. The existence of this MP is mainly due to the virial equilibrium condition

$$M_{\text{JAM}} \propto \sigma_e^2 R_e^{\text{maj}} \quad (1)$$

combined with a smooth variation of galaxy properties with σ_e . For this reason by itself it contains no useful information on galaxy formation. All the useful constraints on galaxy formation models come from the inhomogeneous distribution of galaxies in non-edge-on views of the MP and from the distribution of galaxy properties along the MP.

This paper is devoted to a study of the non edge-on projections of the MP to see what we learn from it on galaxy formation. This is done in the spirit of the classic papers by Bender, Burstein & Faber (1992) and Burstein et al. (1997). However the fact that we have accurate dynamical masses implies that our MP is extremely thin and it follows the scalar virial relation (1) quite accurately. For this reason we can ignore edge-on views of the plane and focus on non edge-on projections only. The thinness of the MP implies that any inclined projection shows essentially the same information, after a change of coordinates. We can use standard and easy-to-understand observables as our main coordinates, instead of trying to observe the plane at a precisely face-on view.

In this paper, in Section 2 we summarize the sample and data, in Section 3 we present our projections of the MP. We illustrate the distribution of a number of quantities on the $(M_{\text{JAM}}, \sigma_e)$ and $(M_{\text{JAM}}, R_e^{\text{maj}})$ projection. We show the variation of the $(M/L)_{\text{JAM}}$, as well as of population indicators of M/L . We show the variation of galaxy concentration, intrinsic shape, morphology and stellar rotation. The variations of the IMF are separately presented in Section 4, together with a review of previous results on the IMF variation. In Section 5 we discuss the implications of our findings for galaxy formation and briefly summarize our paper in Section 6.

2 SAMPLE AND DATA

2.1 Selection and completeness confirmation

The galaxies studied in this work are the 260 early-type galaxies which constitute the volume-limited and nearly mass-selected ATLAS^{3D} sample (Paper I). The objects were morphologically selected as early-type according to the standard criterion (Hubble 1936; de Vaucouleurs 1959; Sandage 1961) of not showing spiral arms or a disk-scale dust lane (when seen edge-on). The early-types are extracted from a parent sample of 871 galaxies of all morphological types brighter than $M_K = -21.5$ mag, using 2MASS photometry (Skrutskie et al. 2006), inside a local ($D < 42$ Mpc) volume of 1.16×10^5 Mpc³ (see full details in Paper I).

In Paper I (section 2.3) we discussed possible incompleteness in the volume-limited parent sample of 871 galaxies, from which the ATLAS^{3D} sample of ETGs was extracted, due to lack of redshift determinations at the faint end limit of our selection. We concluded that the parent sample was 99% complete. We pointed out that the 2MASS redshift survey (2MRS) was going to change the situation by providing redshifts for the all galaxies brighter than the apparent total magnitude $K_T < 11.6$ mag, which makes a galaxy a potential candidate for satisfying our absolute magnitude cut of $M_K < -21.5$ mag, within the adopted $D = 42$ Mpc volume. The 2MRS was recently released with a completeness of 98% for $K_T \leq 11.75$ mag (Huchra et al. 2012). This allowed us to explicitly verify the completeness level estimated in Paper I. For this we

used the redshifts from the 2MRS, in combination with the available redshift-independent distance determinations, to repeat the selection of the ATLAS^{3D} parent sample as described in Paper I. We confirmed that our parent sample has a completeness of 98%, quite close to our original estimation.

In Paper XV we compared our sample to previous samples for which accurate dynamical masses have been determined either via gravitational lensing (e.g. Bolton et al. 2008b; Auger et al. 2010a) or dynamics (e.g. Magorrian et al. 1998; Cappellari et al. 2006; Thomas et al. 2009) and conclude that it provides a major step forward in sample size and accuracy.

2.2 Stellar kinematics and imaging

Various multi-wavelengths datasets are available for the sample galaxies (see a summary in Paper I). In this work we make use of the SAURON (Bacon et al. 2001) integral-field stellar kinematics within about one half-light radius R_e . The kinematics of all galaxies in the ATLAS^{3D} sample was homogeneously extracted as described in Paper I, using the pPXF software¹ (Cappellari & Emsellem 2004) and the full MILES stellar library (Sánchez-Blázquez et al. 2006; Falcón-Barroso et al. 2011) as templates. For the subset of 48 early-types in the SAURON survey (de Zeeuw et al. 2002), the kinematics had already been presented in Emsellem et al. (2004).

The photometry used in this work comes from the Sloan Digital Sky Survey (SDSS, York et al. 2000) data release eight (DR8 Aihara et al. 2011) and was supplemented by our own photometry taken at the 2.5-m Isaac Newton Telescope in the same set of filters and with comparable signal to noise for the rest of the sample galaxies (Scott et al. 2013, hereafter Paper XXI).

2.3 Measuring galaxy global parameters: M/L , R_e and σ_e

The dynamical masses used in this paper were obtained with the dynamical models of the ATLAS^{3D} sample presented in Cappellari et al. (2012) and described in more detail in Paper XV. In brief the modelling approach starts by approximating the observed SDSS and INT r -band surface brightness distribution of the ATLAS^{3D} galaxies using the Multi-Gaussian Expansion (MGE) parametrization (Emsellem, Monnet & Bacon 1994), with the fitting method and software¹ of Cappellari (2002). Full details of the approach and examples of the resulting MGE fits are given in Paper XXI. The MGE models are used as input for the Jeans Anisotropic MGE (JAM) modelling method¹ (Cappellari 2008) which calculates a prediction of the line-of-sight second velocity moments $\langle v_{\text{los}}^2 \rangle$ for a given set of model parameters and fits this to the observed $V_{\text{rms}} \equiv \sqrt{V^2 + \sigma^2}$ using a Bayesian approach (Gelman et al. 2004). The models assume axisymmetry and include a spherical dark halo, which is parametrised according to six different sets of assumptions (See Paper XV for full details). In Cappellari et al. (2012) we showed that the adopted assumptions on the halo have insignificant influence on the measured trend of M/L . For this reason in this paper we only use the two simplest sets of models from that paper (using the same notation):

(A) Self-consistent axisymmetric JAM model, in which the dark matter is assumed to be proportional to the stellar one. We call the total (luminous plus dark) M/L of this model $(M/L)_{\text{JAM}}$;

¹ Available from <http://purl.org/cappellari/idl>

(B) Axisymmetric JAM model with spherical Navarro, Frenk & White (1996) (NFW) dark matter halo. Here the M/L of the stars alone $(M/L)_{\text{stars}}$ is measured directly.

Lablanche et al. (2012, hereafter Paper XII) studied the accuracy in the M/L obtained with the JAM method, using N -body simulations that resemble real galaxies, and concluded that the M/L of unbarred galaxies can be measured with negligible bias $< 1.5\%$, while the M/L of barred galaxies that resemble typical bars found in the ATLAS^{3D} sample can bias the determination by up to 15% in our tests, depending on the position angle of the bars.

The M/L of the stellar population $(M/L)_{\text{Salp}}$ in the SDSS r -band was presented in Cappellari et al. (2012). It was extracted using the penalized pixel-fitting (pPXF) method and software of Cappellari & Emsellem (2004) for full spectrum fitting. In this case the model templates consisted of a regular rectangular grid of 26 ages and 6 metallicities $[M/H]$, assuming a Salpeter (1955) IMF for reference, from the MILES models² of Vazdekis et al. (2012). We used the pPXF keyword REGUL to enforce linear regularization (Press et al. 1992, eq. 18.5.10) in the recovered set of templates weights w_j from the fit. The regularization parameter was chosen for every galaxy to obtain an increase $\Delta\chi^2 = \sqrt{2 \times N_{\text{pix}}}$ in the χ^2 , with respect to a non-regularized fit. In this way the recovered w_j solution constitutes the smoothest one consistent with the observed spectrum (see fig. 20 of Onodera et al. 2012 for an example). This regularized approach helps reducing the noise in the $(M/L)_{\text{Salp}}$ determination, however the results are quite insensitive to the precise value of this parameter and remain essentially unchanged for a range of plausible values. Once the weights for the N model templates have been obtained with pPXF, the properly mass-weighted M/L of the combined population is given by

$$(M/L)_{\text{Salp},r} = \frac{\sum_{j=1}^N w_j M_j^{\text{nogas}}}{\sum_{j=1}^N w_j L_{j,r}}, \quad (2)$$

where M_j^{nogas} is the mass of the j th model in living stars and stellar remnants (black holes and neutron stars), but excluding the gas lost during stellar evolution, and $L_{j,r}$ is the corresponding r -band luminosity of the population. The values are given in Table 1. The details on our spectral fitting of the SAURON data and the analysis of the resulting weight distributions is presented in McDermid et al. (in preparation).

Effective radii are defined as the projected half-light radii of the isophote containing half of the total analytic light of the MGE models from Paper XXI. Both circularized radii R_e and the more robust major axes R_e^{maj} were extracted as described in Paper XV and are taken from table 1 there.

Effective velocity dispersions σ_e were measured by co-adding all the spectra in the SAURON datacube contained within the half-light isophote. The resulting ‘effective’ spectrum was fitted with pPXF using the full MILES stellar library as templates, and assuming a Gaussian line-of-sight velocity distribution. Gas emission around the possible H β and O III emission lines were systematically excluded from the fits and the CLEAN keyword of pPXF was used to reject possible remaining outliers from the spectra. The values for the sample are given in table 1 of Paper XV.

3 PROJECTIONS OF THE MASS PLANE

3.1 Total M/L variations

We have shown in Paper XV that the existence of the FP is due, with good accuracy, to a virial equilibrium condition combined with a smooth variation of galaxy properties, mainly the total mass-to-light ratio M/L , with velocity dispersion. Once this is clarified, the edge-on projection of the MP becomes uninteresting from the point of view of the study of galaxy formation, as it merely states an equilibrium condition satisfied by galaxies and it does not encode any memory of the formation process itself. This is in agreement with previous findings with simulations (Nipoti, Londrillo & Ciotti 2003; Boylan-Kolchin, Ma & Quataert 2006). All information provided by scaling relations on galaxy formation is now encoded in the non edge-on projections of the MP, and first of all in the distribution of M/L on that plane. In Paper XV we also confirmed that M/L correlates remarkably tightly with σ_e (Cappellari et al. 2006). This is especially true (i) for slow rotators, (ii) for galaxies in clusters and (iii) at the high-end of the σ_e range. Here we look at the entire MP and try to clarify the reason for these and other galaxy correlations.

In a classic paper Bender, Burstein & Faber (1992) studied the distribution of hot stellar systems in a three-dimensional space, they called κ space, defined in such a way that one of the axes was empirically defined to lie nearly orthogonal to the plane. This made it easy to look at both the edge-on and face-on versions of the plane. In this paper, thanks to the availability of state-of-the-art integral-field kinematics and the construction of detailed dynamical models, we can use mass as one of the three variables $(M_{\text{JAM}}, \sigma_e, R_e)$. We have shown in Paper XV that in these variables the plane is extremely thin and follows the scalar virial equation $M_{\text{JAM}} \propto \sigma_e^2 R_e^{\text{maj}}$ within our tight errors. This implies that any projection of the plane contains the same amount of information, except for a change of coordinates. Instead of looking at the plane precisely face-on, we constructed special projections which correspond to physically-meaningful and easy-to-interpret quantities.

Our selection of meaningful projections of the MP is shown in Fig. 1. We use as horizontal axis in all plots our main mass variable

$$M_{\text{JAM}} \equiv L \times (M/L)_{\text{JAM}} \approx 2 \times M_{1/2} \approx M_{\text{stars}}, \quad (3)$$

where $(M/L)_{\text{JAM}}$ is the total (luminous plus dark) dynamical M/L obtained using self-consistent JAM models, L is the total galaxy luminosity (both quantities were taken from table 1 of Paper XV) and $M_{1/2}$ is the total mass within a sphere of radius $r_{1/2}$ enclosing half of the total galaxy light, where $r_{1/2} \approx 1.33R_e$ (Hernquist 1990; Ciotti 1991; Wolf et al. 2010; Paper XV). Although the self-consistency assumption, where the total mass is proportional to the stellar mass, is not justified at large radii, it is accurate within the region where we have stellar kinematics (about $1R_e$). It was shown in Williams, Bureau & Cappellari (2010) and in Paper XV that $(M/L)_{\text{JAM}}$ closely reproduces the total $(M/L)(r = R_e)$ inside a sphere (actually an iso-surface) with mean radius the projected half-light radius R_e , derived using models which explicitly include dark matter. Different assumptions on the dark halo produce minor differences in $(M/L)(r = R_e)$. Given that $(M/L)(r = R_e)$ is nearly insensitive to the choice of the halo assumptions, we choose the self-consistent one, being the simplest.

As illustrated in Paper XV (see also Williams, Bureau & Cappellari 2009), most of the galaxies in our sample are consistent with having small fractions of dark matter within a sphere of radius $r = R_e$, with a median value of just 13% of DM within that

² Available from <http://miles.iac.es/>




Figure 1. The MP and its projections. The top two panels show the two main projections of the MP in the $(M_{\text{JAM}}, \sigma_e)$ and $(M_{\text{JAM}}, R_e^{\text{maj}})$ coordinates. Overlaid are lines of constant $\sigma_e = 50, 100, 200, 300, 400, 500 \text{ km s}^{-1}$ (dashed blue), constant $R_e^{\text{maj}} = 0.1, 1, 10, 100 \text{ kpc}$ (dot-dashed red) and constant $\Sigma_e = 10^8, 10^9, 10^{10}, 10^{11} M_\odot \text{ kpc}^{-2}$ (dotted black) transformed in different panels using the scalar virial relation $M_{\text{JAM}} = 5.0 \times \sigma_e^2 R_e / G$. The observed $(M_{\text{JAM}}, \sigma_e, R_e^{\text{maj}})$ points follow the scalar virial relation so closely that the virial coordinates provide a unique mapping on these diagrams and one can reliably infer all characteristics of the galaxies from any individual projection. In each panel the galaxies are coloured according to the (LOESS smoothed) $\log(M/L)_{\text{JAM}}$, as shown in the colour bar at the top. The LOESS method tries to estimate the mean values of the underlying galaxy population, namely the values one should expect to obtain by averaging much larger galaxy samples. In all panels the thick red line shows the same ZOE relation given by equation (4), again transformed in different panels according to the scalar virial relation. The thin white line in the top-left panel is a LOESS smoothed version of the $M_{\text{JAM}} - \sigma_e$ relation, while the thick dark grey line is a double power-law fit (equation 5) with trends $M_{\text{JAM}} \propto \sigma_e^{2.3}$ and $M_{\text{JAM}} \propto \sigma_e^{4.7}$ below or above $\sigma_e \approx 140 \text{ km s}^{-1}$ respectively. While the top two panels contain different observable quantities, the bottom two panels merely apply a coordinate transformation to the quantities in the top two panels, to show the effective phase-space density $f_{\text{eff}} \equiv 1/(\sigma R_e^2)$ and effective mass surface density $\Sigma_e \equiv M_{\text{JAM}}/(2\pi R_e^2)$. Two galaxies stand out for being significantly above the ZOE in the $(M_{\text{JAM}}, \sigma_e)$ and $(M_{\text{JAM}}, \Sigma_e)$ projections. The top one is NGC 5845 and the bottom one is NGC 4342.

radius for our standard models (B). This implies that $M_{1/2}$ is dominated by the stellar mass. For this reason M_{JAM} is a quantity that very closely represents and is directly comparable to the total stellar galaxy mass used in numerous previous studies. Stellar mass seems to relate well with galaxy properties and is often used to study galaxy formation (e.g. Kauffmann et al. 2003a,b; Hyde & Bernardi 2009b). The difference of our mass parameter is that it does not suffer from the uncertainties related to the stellar population models (e.g. Maraston et al. 2006; Gallazzi & Bell 2009; Conroy, Gunn & White 2009; Longhetti & Saracco 2009; Wuyts et al. 2009) moreover it automatically includes the effects of a non-universal IMF (van Dokkum & Conroy 2010; Cappellari et al. 2012). Being a measure of the total enclosed mass within a spherical region, and thus

being directly related the dynamics, M_{JAM} is the ideal parameter to use in scaling relations. Note that M_{JAM} , unlike mass determinations obtained via strong lensing, does not include the possible contribution of dark matter along the cylinder parallel to the line-of-sight (Dutton et al. 2011a), which provides an useful additional constraint on dark matter at large radii, but complicates the interpretation of scaling relations in the galaxy centres.

In the top-left and top-right panels of Fig. 1 we show the projections of the MP along the $(M_{\text{JAM}}, \sigma_e)$ and $(M_{\text{JAM}}, R_e^{\text{maj}})$ axes respectively. The colour in these diagrams represent the dynamical (total) M/L inside a sphere of radius R_e . Three important results are clear from these plots:

- (i) Both projections are equivalent and provide basically the

same picture, apart from a coordinate transformation. This is expected from the tightness of the MP. Moreover, given that the MP nearly follows the scalar virial relation, the expression $M_{\text{JAM}} = 5.0 \times \sigma_e^2 R_e / G$ (Cappellari et al. 2006) can be used to provide an accurate estimate of lines of constant R_e on the $(M_{\text{JAM}}, \sigma_e)$ projection or the lines of constant σ_e on the $(M_{\text{JAM}}, R_e^{\text{maj}})$ projection;

(ii) Galaxies define a clear zone-of-exclusion (ZOE) at small sizes or large densities, as already pointed out by Bender, Burstein & Faber (1992) and Burstein et al. (1997), however we find a clear break in the ZOE at a mass $M_{\text{JAM}} \approx 3 \times 10^{10} M_\odot$ and two nearly power-law regimes above or below this value. The ZOE is approximated by the equation

$$R_e^{\text{maj}} = R_{e,b} \left(\frac{M_{\text{JAM}}}{M_{\text{JAM},b}} \right)^\gamma \left[\frac{1}{2} + \frac{1}{2} \left(\frac{M_{\text{JAM}}}{M_{\text{JAM},b}} \right)^\alpha \right]^{(\beta-\gamma)/\alpha} \quad (4)$$

with $R_{e,b} = 0.7$ kpc, $\alpha = 8$, $\beta = 0.75$, $\gamma = -0.20$. The relation has an asymptotic trend $R_e \propto M_{\text{JAM}}^{0.75}$ above $M_{\text{JAM},b} = 3.0 \times 10^{10} M_\odot$, and a sharp transition into $R_e \propto M_{\text{JAM}}^{-0.20}$ below this break. The values were determined by simultaneously matching the observed boundary in the galaxy distribution in all MP projections (but see Fig. 2 for a quantitative definition). These values are nearly the same to those we already reported in Paper I (eq. 4 there), using K -band luminosity in place of mass and 2MASS R_e instead of MGE ones. The maximum in the galaxy density at $M_{\text{JAM},b}$ that we infer from our sample is also clearly visible in much larger SDSS samples (e.g. fig. 2 of van Dokkum et al. 2008). This shows that the relation is valid for the general galaxy population. The ZOE relation of equation (4) can be converted into a $(M_{\text{JAM}}, \sigma_e)$ one, or into other projections using the scalar virial relation $M_{\text{JAM}} = 5.0 \times \sigma_e^2 R_e / G$. The corresponding trend becomes $\sigma_e \propto M_{\text{JAM}}^{0.6}$ ($M_{\text{JAM}} \propto \sigma_e^{1.67}$) below the break, which flattens to $\sigma_e \propto M_{\text{JAM}}^{0.125}$ ($M_{\text{JAM}} \propto \sigma_e^{8.0}$) above the break. The location of the break we find agrees with the value at which scaling relations of large sample of SDSS galaxies show a subtle deviation from a straight line (Hyde & Bernardi 2009a). Interestingly the characteristic mass at the cusp also coincides with the value reported by Kauffmann et al. (2003b), as the fundamental dividing line between the two distinct families of passive and star-forming galaxies.

(iii) The contours of constant $(M/L)_{\text{JAM}}$ closely follow lines of constant σ_e in the $(M_{\text{JAM}}, \sigma_e)$ projection (above $\sigma_e \gtrsim 120 \text{ km s}^{-1}$) and equivalently $R_e^{\text{maj}} \propto M_{\text{JAM}}$ in the $(M_{\text{JAM}}, R_e^{\text{maj}})$ projection. A comparable agreement is reached using lines parallel to the ZOE $R_e^{\text{maj}} \propto M_{\text{JAM}}^{0.75}$. Neither mass M_{JAM} , nor size R_e nor surface mass density Σ_e provide a comparably good approximation to the $(M/L)_{\text{JAM}}$ contours, although Σ_e provides a better approximation than the other two. This is consistent and explains the finding by Cappellari et al. (2006) that σ_e and not mass or luminosity, is the best tracer of $(M/L)_{\text{JAM}}$. At lower $\sigma_e \lesssim 120 \text{ km s}^{-1}$ the $(M/L)_{\text{JAM}}$ contours start deviating from the lines of constant σ_e and tend to lie closer to lines of constant M_{JAM} . Given that the M/L describes the deviations between the FP and the MP, this twist in the contours demonstrates that the FP is warped with respect to the MP, and explains the sensitivity of the FP parameters to the region included in the fit (D’Onofrio et al. 2008; Gargiulo et al. 2009; Hyde & Bernardi 2009b).

To quantitatively define the meaning of the ZOE, which is shown in various plots of this paper, we estimated the probability density underlying the set of discrete measurements for our galaxies in the M-size plane. For this we used the kernel density estimator method. We adopted an Epanechnikov kernel and determined the optimal smoothing parameter in each dimension following Silver-




Figure 2. Probability density contours for ETGs on the M-size plane. Colours indicate the contours enclosing a given fraction of the total probability, with the outermost (lowest) contour enclosing 99% of the total probability, above our survey mass limit ($M_{\text{JAM}} \gtrsim 6 \times 10^9 M_\odot$). The thick red curve is the same as in Fig. 1 and provides a convenient approximation for the lowest 99% contour. The dashed line indicates the relation $(R_e^{\text{maj}}/\text{kpc}) = 4.2 \times [M_{\text{JAM}}/(10^{11} M_\odot)]^{0.46}$, which approximates the peak ridge-line of the ETGs M-size distribution.

man (1986). The resulting estimate of the ETGs density is shown in Fig. 2. The plot shows that the adopted functional form of the ZOE in equation (4) provides a convenient approximation of the lower contour enclosing 99% of the ETG population, above our survey mass limit $M_{\text{JAM}} \gtrsim 6 \times 10^9 M_\odot$. We argue that, for volume-limited samples, this line constitutes a more meaningful reference to define the lower limit of the sizes of the normal galaxy population than the mean M-size relation and its scatter (e.g. Shen et al. 2003). The reason is that the ZOE takes the galaxies mass function into account. Low-mass galaxies are more numerous than high-mass ones. This implies that, for a given rms scatter σ_{R_e} , low-mass galaxies are more likely than high-mass galaxies to deviate more than σ_{R_e} from the mean M-size relation.

Two galaxies stand out for being significantly above the ZOE in the $(M_{\text{JAM}}, \sigma_e)$ and $(M_{\text{JAM}}, \Sigma_e)$ projections. The top one is NGC 5845 and the bottom one is NGC 4342. The two objects have very high surface brightness and consequently excellent kinematic and photometric data. They are genuine examples of dense objects in the nearby Universe. These two outliers were already presented in Cappellari (2011a) and their compactness was later discussed by Jiang et al. (2012). However they have smaller masses ($M_{\text{JAM}} \approx 3 \times 10^{10} M_\odot$) than their high-redshift counterparts (Cimatti et al. 2008; van Dokkum et al. 2008).

In the top-left panel of Fig. 1 we overplot a version of the $M_{\text{JAM}} - \sigma_e$ relation which was adaptively smoothed using the one-dimensional Locally Weighted Regression robust technique

(dubbed LOESS³) of Cleveland (1979). The relation is equivalent to the Faber & Jackson (1976) but uses dynamical mass instead of luminosity. Like the ZOE, the mean $M_{\text{JAM}} - \sigma_e$ relation appears well approximated by a double power-law of the form:

$$\sigma_e = \sigma_{e,b} \left(\frac{M_{\text{JAM}}}{M_{\text{JAM,b}}} \right)^\gamma \left[\frac{1}{2} + \frac{1}{2} \left(\frac{M_{\text{JAM}}}{M_{\text{JAM,b}}} \right)^{\alpha\gamma} \right]^{(\beta-\gamma)/\alpha}, \quad (5)$$

with best-fitting parameters (excluding NGC 5845 and NGC 4342) $\log(M_{\text{JAM,b}}/M_\odot) = 10.675 \pm 0.090$, $\log(\sigma_{e,b}/\text{km s}^{-1}) = 2.153 \pm 0.031$, $\alpha = 8$, logarithmic slope $\beta = 0.215 \pm 0.027$ at large masses and $\gamma = 0.442 \pm 0.028$ at small masses. When the two power-laws are written as a function of σ_e , the trends become $M_{\text{JAM}} \propto \sigma_e^{1/\beta} = \sigma_e^{4.65 \pm 0.58}$ for $\sigma_e \gg 140 \text{ km s}^{-1}$, and $M_{\text{JAM}} \propto \sigma_e^{1/\gamma} = \sigma_e^{2.26 \pm 0.14}$ for $\sigma_e \ll 140 \text{ km s}^{-1}$.

In all panels of Fig. 1, and in all subsequent similar figures for other parameters, the measured $(M/L)_{\text{JAM}}$ have been adaptively smoothed using the LOESS technique, straightforwardly generalized to two dimension as in Cleveland & Devlin (1988). In all plots we adopt a regularization factor $f = 0.6$, and a linear local approximation. To deal with the different scales of the axes and the elongated shape of the galaxy distributions, before applying LOESS we rotated and re-normalized the coordinates so that the ellipse of inertia of the galaxy distribution reduces to a circle in every projection³. This change of coordinates improves the definition of neighbouring points during the LOESS smoothing. By design the LOESS smoothed maps try to remove observational errors and intrinsic scatter to estimate the mean values of the underlying galaxy population. The maps try to predict the mean values one would obtain via simple histograms of much larger samples of galaxies (see also discussion in Section 3.2). We do not show here the original data, as the scatter in $(M/L)_{\text{JAM}}$, projected along the σ_e coordinate, was already analysed in detail in Paper XV. Given that most of the variation in the smooth surface happens orthogonally to the constant σ_e lines, the scatter from the smooth surface of 27% is nearly unchanged from the value of 29% in the global $(M/L) - \sigma_e$ relation, further confirming that most of the variation happens orthogonally to σ_e .

In the bottom left and right panels of Fig. 1 we show the effective phase-space density, defined following Hernquist, Spergel & Heyl (1993) as $f_{\text{eff}} \equiv 1/(G\sigma_e R_e^2)$ and the mass surface density, defined as $\Sigma_e \equiv M_{\text{JAM}}/(2\pi R_e^2)$. Note that, while we use R_e^{maj} for the mass-size plane, one has to use R_e for f_{eff} and Σ_e . The two bottom panels do not plot new data, but are obtained by rearranging the three variables shown in the top panels, however they illustrate how the above trends relate to other physical quantities. In particular the phase-space density is interesting because it can only decrease during collisionless galaxy mergers, due to Liouville’s theorem (Carlberg 1986; Hernquist, Spergel & Heyl 1993). Interestingly the ZOE we find is nearly flat in f_{eff} below $M_{\text{JAM,b}} = 3 \times 10^{10} M_\odot$ (a value $\gamma = 1/3$ gives $f_{\text{eff}} = \text{const}$ in the virial case) and starts decreasing at larger masses, as one would expect when dry mergers start becoming more important.

3.2 Stellar population indicators of $(M/L)_{\text{pop}}$

We have shown in Paper XV that the stellar matter dominates in the regions we study. If this is indeed the case we would expect stellar

population indicators of the $(M/L)_{\text{pop}}$ of the stellar population to closely follow the behaviour of the dynamical $(M/L)_{\text{JAM}}$. Before addressing this question with our own data, the reader is strongly encouraged to compare Fig. 1 to the right panel of figure 15 of Gallazzi et al. (2006), which shows the luminosity-weighted age versus the stellar mass in the (σ, M_*) projection. Even though our Fig. 1 uses dynamical quantities, which are measured via dynamical models, while Gallazzi et al. (2006) derives population from line strengths, the resemblance of the two plots is striking. This includes rather subtle details like the change of orientation of the contours, which in both cases tend to be horizontal above $\sigma \gtrsim 120 \text{ km s}^{-1}$ and become nearly vertical at lower σ . This comparison already indicates a close link between $(M/L)_{\text{JAM}}$ and stellar population (mainly age), in broad agreement with Cappellari et al. (2006). Importantly, given that our sample is 100 times smaller than the one of Gallazzi et al. (2006), this comparison also provides a validation of the adopted LOESS technique, which appears to recover as expected the underlying distribution of galaxy properties from our much smaller galaxy sample. This comparison confirms that the various LOESS smoothed plots in this paper should be interpreted as estimates of the average trends in the galaxy distribution we would observe if we had much larger samples of galaxies than we have in ATLAS^{3D}.

We now address the relation between dynamical and population M/L using our own dataset and a different approach. Two main simple tracers of $(M/L)_{\text{pop}}$ have been proposed in the past: (i) the $B - R$ colour, which was shown by (Bell & de Jong 2001) to trace $(M/L)_{\text{pop}}$ alone, for a wide range of metallicities, and (ii) the line-strength index $H\beta$, which was shown to satisfy a similar property (fig. 16 of Cappellari et al. 2006) for a wide range of metallicities (see also Worthey 1994). Here we adopt the $H\beta$ determination for our ATLAS^{3D} galaxies, as derived from the same very high S/N ‘effective’ SAURON spectra from which we derived σ_e . The homogeneous extraction of the line-strength parameters for the ATLAS^{3D} sample is given in McDermid et al. (in preparation). The parameters for the SAURON survey (de Zeeuw et al. 2002) subset of 48 galaxies was already given in Kuntschner et al. (2006). As choice of colour we use the SDSS $g - i$ one, which is available for 223 ATLAS^{3D} galaxies in SDSS DR8 and was shown by Gallazzi & Bell (2009) to provide on average the smallest uncertainties and the most stable results, among the SDSS bands.

The results are presented in Fig. 3 and show a quite good agreement between simple population estimators of M/L and the dynamical one. Both $H\beta$ (top panel) and colour (middle panel) are nearly constant along lines of constant σ_e as it was the case for Fig. 1. This confirms the fact that the key driver of the total dynamical $(M/L)_{\text{JAM}}$ is the stellar population. We do notice some systematic differences at the low- σ end, with the contours of constant $(M/L)_{\text{JAM}}$ not quite following the population trends. As the effect is visible both in colour and $H\beta$ we believe it is significant. It may be due to the presence of extended star formation episodes, which are expected to be more prevalent at low mass (Heavens et al. 2004; Thomas et al. 2005) and cause deviations from the simple one-to-one relations between population observables and M/L . The study of the relation between our parameters and the actual physical parameters of the population, like age, metallicity and abundance ratios, is presented in McDermid et al. (in preparation). Analysis of the stellar population parameters for the SAURON survey (de Zeeuw et al. 2002) subset of 48 galaxies was already presented in Kuntschner, Emsellem & et al. (2010).

In fig. 15 of Young et al. (2011, hereafter Paper IV) we showed that galaxies with the largest $H\beta$, and consequently younger mean

³ Our implementations CAP_LOESS_1D and CAP_LOESS_2D of the one-dimensional and two-dimensional LOESS method as used in this paper are available from <http://purl.org/cappellari/idl>




Figure 3. Indicators of the stellar population $(M/L)_{\text{pop}}$ on the MP. *Top Panels:* Same as in Fig. 1, with colours showing the $\log H\beta$ stellar absorption line-strength, from McDermid et al. (in preparation). *Middle Panels:* Same as in Fig. 1, with colours indicating the SDSS galaxy colour $g - i$. *Bottom Panels:* Same as in Fig. 1 with colour indicating the fraction of molecular hydrogen H_2 (from table 1 of Paper IV) with respect to M_{JAM} . All three indicators $H\beta$, $g - i$ and the H_2 fraction show a similar trends as the $(M/L)_{\text{JAM}}$ in the top panels of Fig. 1. This suggests that the stars dominate the region we observe and that the dynamical $(M/L)_{\text{JAM}}$ is driven by the stellar population, in agreement with the quantitative results via dynamical models in Paper XV.

ages, contain higher fractions of molecular hydrogen H_2 as traced by CO emission lines. The trend makes sense given that H_2 reservoir is expected to fuel star formation (see a more detailed discussion in Paper IV). Given the trend we observed for $H\beta$ on the M-size plane, one should expect a similar trend for the H_2 fraction. This is indeed the case as presented in the bottom panel of Fig. 3, which shows the mass $M(H_2)$ as fraction of M_{JAM} . Given that not all galaxies have a CO detection in Paper IV, we assigned to undetected galaxies a uniformly distributed random H_2 mass between zero and the quoted upper limit. A similar plot is obtained by assigning to all undetected galaxies a fixed mass $M(H_2) = 10^6 M_\odot$ below our CO survey sensitivity, or by assigning the upper limit, showing that the trend does not depend critically on our treatment of the undetected galaxies. The plot shows that the molecular gas content, like other population indicators, also tends to follow contours of constant σ_e . However some notable differences exist between this plot and the $(M/L)_{JAM}$ or the $H\beta$ ones. In particular above the mass $M_{JAM} \approx 2 \times 10^{11} M_\odot$, the likelihood of finding significant fractions of molecular gas seems to depend more on mass than σ_e . This mass is a characteristic one, where a number of other observables appear to change as we will show in Section 3.4.

3.3 Inferring bulge fractions from stellar kinematics

In the previous sections we showed that galaxy properties follow smooth trends in the projections of the MP. We have seen that the dynamical M/L near the galaxy centre ($1R_e$) is driven by a variation of the stellar population (including IMF), with the dynamical M/L constituting an accurate tracer of the galaxy population! These trends distribute galaxies in the MP along lines of nearly constant σ_e on the (M_{JAM}, σ_e) plane, or equivalently $R_e^{maj} \propto M_{JAM}$ on the (M_{JAM}, R_e^{maj}) plane. In Paper I we showed that the luminosity-size relation changes gradually with morphological type. Galaxy size decreases as a function of the bulge fraction as indicated by the morphological classification. We explained the size decrease as an increase of the bulge fraction between different morphological types. In Paper II and III, we showed that the vast majority of local ETGs is constituted by disk-like systems, the fast rotators, and in Cappellari et al. (2011b, hereafter Paper VII) we discussed how fast-rotators morphologically resemble spirals with a variety of bulge fractions. We concluded that fast rotators simply constitute the end point of a smooth sequence of disks with increasing bulges, and this led us to propose a fundamental change in our view of galaxy morphology (Paper VII). If this picture is correct, we should be able to find direct evidence for a systematic change in the bulge fraction, in the fast rotators class, while moving from the region of the MP populated by spiral galaxies towards the ZOE, dominated by the oldest and reddest ETGs, with the largest M/L . This is what we show in this section.

A detailed photometric bulge-disk decomposition of the unbarred galaxies subset of the ATLAS^{3D} sample is presented Krajinović et al. (2013, hereafter Paper XVII). Although it provides a well established definition of the bulge luminosity, it is a complex technique which depends on the extraction details and suffers from degeneracies. In particular it is well established that, even assuming galaxies to be axisymmetric, the intrinsic stellar density cannot be uniquely recovered from the images unless galaxies are edge-on (Rybicki 1987; Gerhard & Binney 1996; van den Bosch 1997; Magorrian et al. 1998). For this reason, unless we knew galaxies to be accurately described by Sersic (1968) bulges and exponential disks (Freeman 1970), we should not expect to be able to uniquely decompose bulges and disks of inclined galaxies, as this would im-

ply we can uniquely infer their intrinsic densities. This is likely one of the reasons why we do not find trends in the bulge fraction (parametrized by the disk-over-total luminosity ratio D/T) of fast rotators on the M-size plane. An exception is the important and clear lack of disks above the characteristic $M_{JAM} \gtrsim 2 \times 10^{11} M_\odot$.

In our survey we have SAURON integral-field kinematics (Paper I) for all the galaxies in the sample. Here we exploit that extra information to robustly infer the bulge fractions even at low inclinations. The reason why this is possible is because the kinematics retains information on the *flattening* of the galaxy components even at low inclinations, when no such information can be inferred from the images. In this situation the photometric decomposition has to rely entirely on the radial profiles and the *assumption* of a specific functional form for the two components. The ability of the kinematics to uncover disks and bulges even at low inclination is illustrated in Fig. 4. For the plot we constructed a model with a Sersic bulge and an exponential disk. To have a representative model, we adopted the median ratio $R_e^{disk}/R_e^{bulge} \approx 5.2$ and the corresponding median Sersic index for the bulges $n \approx 1.7$ (from table C1 of Paper XVII). We used the MGE_FIT_1D procedure¹ of Cappellari (2002) to obtain an accurate MGE fit to the one-dimensional profiles of both the bulge and the disk. We then assigned a typical intrinsic axial ratio $q_d = 0.2$ to the disk and $q_b = 0.7$ to the bulge. The Gaussian components of bulge and disk were then combined into a single MGE model that was used as input for a JAM dynamical model (Cappellari 2008), which provided a prediction for the second velocity moments $\langle v_{los}^2 \rangle^{1/2}$. It was shown in Cappellari (2008), Scott et al. (2009) and for the full ATLAS^{3D} sample in Paper XV, that the second moments calculated by the JAM method provide good predictions of the observed $V_{rms} \approx \sqrt{V^2 + \sigma^2}$ of real galaxies, where V is the mean stellar velocity and σ is the stellar velocity dispersion. The only free parameter in this model is the anisotropy in the (v_R, v_z) plane, which is parametrized by $\beta_z \equiv 1 - \sigma_z^2/\sigma_R^2$, for which we adopted a typical value $\beta_z = 0.25$ (Cappellari 2008). While keeping the bulge and disk parameters fixed, we varied their relative contribution D/T and we projected the models at different inclinations. Fig. 4 shows that the bulge fraction provides dramatic changes in the relative values of the V_{rms} in the galaxy centre with respect to its value at large radii. The presence of disks can still be inferred from the kinematics as a horizontal elongation in the V_{rms} maps, even for disk fractions as small as $D/T = 0.1$, and even for inclination as low as $i = 30^\circ$. Pure spheroids have V_{rms} maps roughly following the galaxy isophotes. Although the details of the V_{rms} maps are different for different galaxies, the qualitative trends shown in the figure are similar for quite different bulge and disk parameters.

In Fig. 5 we show the maps of V_{rms} actually observed with SAURON for a representative subsample of 40 galaxies in the ATLAS^{3D} sample, uniformly sampling the M-size plane (the full set of maps is shown in Paper XV). The plot shows that near the line $\sigma_e = 50 \text{ km s}^{-1}$ all maps present the minimum of V_{rms} , which is characteristic of pure disk galaxies. The minimum disappears around the line $\sigma_e = 100 \text{ km s}^{-1}$, where the maps show the butterfly-like shape expected for galaxies with small bulges and dominant disks. For larger σ_e the central peaks in V_{rms} become the norm and finally, near the line $\sigma_e = 200 \text{ km s}^{-1}$, all galaxies show prominent V_{rms} peaks, indicating bulges as massive or more massive than the disks. The situation is, once more, qualitatively different above the characteristic mass $M_{JAM} \gtrsim 2 \times 10^{11} M_\odot$. Above that mass the V_{rms} maps lack the horizontal V_{rms} elongation. This indicates they don't have disks and consist of pure spheroid, in agreement with the bulge-disk decompositions of Paper XVII.




Figure 4. Inferring bulge fraction from stellar kinematics. The top panels show the prediction for the second velocity moment $\langle v_{\text{los}}^2 \rangle^{1/2}$ using JAM dynamical models. This quantity approximates the observed $V_{\text{rms}} \equiv \sqrt{V^2 + \sigma^2}$, where V is the mean stellar velocity and σ the velocity dispersion. The top panels show the predicted kinematics for edge-on realistic galaxy models with different bulge fractions, quantified by the disk-over-total luminosity parameter D/T . The middle and bottom panels show the same models, projected at different inclinations. The plots show that for any inclination: (i) an increase of the bulge fraction produces a peak in the central V_{rms} ; (ii) the presence of a disk produces an horizontal elongation of the V_{rms} contours.

To try to generalize the qualitative results from the maps of Fig. 5 we need to encode the key information on the shape of the V_{rms} maps into a simple parameter that can be extracted for the full volume-limited ATLAS^{3D} sample. We found that the following simple approach is sufficient⁴. We quantified the variety of shapes in Fig. 4 by simply measuring the ratio between the central velocity dispersion $\sigma(R_e/8)$ (given in Table 1) measured with pPXF (with keyword MOMENTS=2) within the commonly used circular aperture of radius $R = R_e/8$ and our reference σ_e (given in table 1 of Paper XV), measured within the ‘effective’ isophote of major axis R_e^{maj} . The distribution of the $\sigma(R_e/8)/\sigma_e$ ratio on the MP projections is shown in Fig. 6. It shows that the ratio still approximately follows lines of constant σ_e (or even better lines parallel to the slightly more shallow ZOE $R_e^{\text{maj}} \propto M_{\text{JAM}}^{0.75}$). This confirms that the trends in the bulge fractions we saw from the maps of the galaxy subset in Fig. 5 are also valid in a statistical sense for the whole ATLAS^{3D} sample. It demonstrates that at given mass, σ_e traces the bulge fraction, which in turn appear as the main driver for the trends in M/L , population parameters and molecular gas content that we observe.

Interestingly the plot also shows a slight decrease in the $\sigma(R_e/8)/\sigma_e$ ratio above the characteristic mass $M_{\text{JAM}} \gtrsim 2 \times 10^{11} M_{\odot}$. This value coincides with the characteristic scale in ETGs above which galaxies start to be rounder (Tremblay & Merritt 1996; van der Wel et al. 2009b), nearly all have flat (core/deficit) central

surface brightness profiles (Faber et al. 1997; Graham & Guzmán 2003; Ferrarese et al. 2006a; Lauer et al. 2007a), they deviate from power-law colour-magnitude relations (Baldry et al. 2004; Ferrarese et al. 2006a; Bernardi et al. 2011) and are embedded in massive X-ray halos (Kormendy et al. 2009; Sarzi et al. 2013, hereafter Paper XIX).

In the bottom panel of Fig. 6 we show for comparison the distribution of the Sersic (1968) index n (from Paper XVII) on the MP projection. These values were obtained by fitting a single Sersic function $I(R) \propto \exp(-kR^{1/n})$ to the entire galaxy profile and do not necessarily provide good descriptions of the galaxies. However this kind of approach is routinely used to characterize sizes and morphologies of large galaxy samples (e.g. Blanton et al. 2003; Shen et al. 2003) or to study galaxies scaling relations at high-redshift (e.g. Trujillo et al. 2007; Cimatti et al. 2008; van Dokkum et al. 2008). The plot clarifies that the Sersic index behaves quite differently from the others galaxy indicators and in particular it is weakly sensitive to the bulge fraction. The global Sersic index tends to vary along lines of nearly constant mass rather than σ_e . Unlike σ_e , the global Sersic index is not a good predictor of galaxy properties. The plot also clarifies that defining ETGs as having $n > 2.5$ (e.g. Shen et al. 2003) tends to select ETGs above the characteristic mass $M_{\text{JAM}} \approx 3 \times 10^{10} M_{\odot}$, while defining genuine nearby elliptical galaxies as having $n > 4$ tends to select objects above the characteristic mass $M_{\text{JAM}} \approx 2 \times 10^{11} M_{\odot}$.

3.4 Variations of galaxy intrinsic flattening and rotation

Our dynamical modelling effort provides the unique opportunity of having the galaxy inclination for the entire ATLAS^{3D} sample of galaxies (from table 1 of Paper XV). We verified that the JAM inclination agrees with the one inferred from the geometry of the dust,

⁴ For more quantitative results, one could try to break the degeneracy of the photometric bulge-disk decomposition approach by fitting simultaneously both the galaxy images, e.g. with a Sersic+exponential model, and the V_{rms} maps with the corresponding JAM prediction for different (i, β_z) . Although interesting and in principle straightforward, this goes beyond the scope of the present paper




Figure 5. Observed $V_{\text{rms}} \equiv \sqrt{V^2 + \sigma^2}$ maps on the M-size plane. All galaxies are oriented with their kinematical major axis PA_{kin} (from Paper II) horizontal. A characteristic isophote is indicated by the thick black contour. The plot shows that: (i) near the $\sigma_e \approx 50 \text{ km s}^{-1}$ line, galaxies show a drop in the central V_{rms} characteristic of pure disks (see Fig. 4); (ii) near the $\sigma_e \approx 100 \text{ km s}^{-1}$ line the maps show a butterfly-like shape characteristic of galaxies with small bulges; (iii) near the $\sigma_e \approx 200 \text{ km s}^{-1}$ line the maps show strong peaks of V_{rms} , indicative of bulges as massive or more massive than the disks, but horizontal elongations due to disks are still present; (iv) above the characteristic mass $M_{\text{JAM}} \gtrsim 2 \times 10^{11} M_{\odot}$ no horizontal elongations of V_{rms} are seen any more, indicating galaxies are pure spheroids. The latter finding agrees with the classification of the objects as slow rotators (Paper III) and the lack of disks in their bulge-disk decomposition (Paper XVII).

for 26 galaxies with regular dusts disks (see also Cappellari 2008). We further verified that we can recover with JAM models the inclination of simulated galaxies that resemble typical objects in our sample (Paper XII). Although one should not expect the inclination to be reliable for every individual galaxy, it is expected to be accurate for most of our sample. The inclination allows us to recover the intrinsic shape of individual objects. The deprojection is done using the ellipticity ε of the outer isophote, measured in Paper II at radii typically around $4R_e$, at the depth limit of the SDSS photometry. At these radii, when a disk is present, the ellipticity is representative of the outer disks and is insensitive to the possible presence of a bulge. We then computed the intrinsic ellipticity assuming an oblate spheroid geometry as (Binney & Tremaine 2008)

$$\varepsilon_{\text{intr}} = 1 - \sqrt{1 + \varepsilon(\varepsilon - 2)/\sin^2 i}, \quad (6)$$

The results of the deprojection is shown in the top panels of Fig. 7. Contrary to all previous diagrams presented so far, the distribution of the intrinsic ellipticity for ETGs on the MP shows a completely different trend. Few galaxies stand out in the top-right corner for being nearly round, with average intrinsic ellipticity $\varepsilon_{\text{intr}} \approx 0.3$.

These galaxies are all located above the same characteristic mass $M_{\text{JAM}} \gtrsim 2 \times 10^{11} M_{\odot}$, where other observables indicated a transition in galaxy properties. This result confirms previous statistical studies of ETGs shapes (Tremblay & Merritt 1996; van der Wel et al. 2009b; Bernardi et al. 2011). Below this characteristic mass, the mean ellipticity drops dramatically and sharply to about $\varepsilon_{\text{intr}} \approx 0.6$, characteristics of disks. This galaxy-by-galaxy shape deprojection confirms a similar result on their shape distribution inferred via Monte Carlo simulations from the distribution of fast rotators ETGs on the (λ_R, ε) diagram, suggesting fast rotators have a typical intrinsic ellipticity $\varepsilon_{\text{intr}} \approx 0.7$ (fig. 15 in Paper III). It also agrees with a statistical inversion of their shape (Weijmans et al. 2013, hereafter Paper XXIII). This agreement also seems to confirm the reliability of the inclinations derived via the JAM models. The result strongly confirms our previous conclusions that fast rotators as a class are disk like (Paper II, III, VII). At the lowest mass range ($M_{\text{JAM}} \lesssim 10^{10} M_{\odot}$), near the ZOE, there seems to be a marginal decrease of $\varepsilon_{\text{intr}}$. The significance or reality of this feature is however unclear.




Figure 6. Galaxy concentration on the M-size plane. *Top Panels:* Same as in Fig. 1 for the ratio $\sigma(R_e/8)/\sigma_e$, between the velocity second moment measured within a circular aperture of radius $R = R_e/8$ and the second moment σ_e (which includes both rotation and random motions) within the ‘effective’ isophote of semi-major axis R_e^{maj} . As shown in Fig. 4, this quantity is related to the bulge fraction, and is weakly sensitive to projection effects. Like for $(M/L)_{\text{JAM}}$, the variation of this parameter tends to follow lines of constant σ_e , or equivalently $R_e^{\text{maj}} \propto M_{\text{JAM}}$, or even better, it follows lines parallel to the ZOE above its break. Note the slight decrease of $\sigma(R_e/8)/\sigma_e$ above the characteristic mass $M_{\text{JAM}} \gtrsim 2 \times 10^{11} M_\odot$. *Bottom Panels:* for comparison this shows the variation of the global Sersic index n for the galaxies, as measured in Paper XVII. n does not follow lines of constant σ_e and it is not a good empirical predictor of galaxy properties. Above $M_{\text{JAM}} \gtrsim 2 \times 10^{11} M_\odot$ galaxies tend to have the value $n \gtrsim 4$, which is often associated to pure spheroids.

A similar trend as for the intrinsic shape is seen in the bottom panel of Fig. 7 for the optical morphology, as quantified by the T-type parameter from Hyperleda (Paturel et al. 2003). Above $M_{\text{JAM}} \gtrsim 2 \times 10^{11} M_\odot$ nearly all galaxies are classified as ellipticals ($T < -3.5$), consistently with the fact that galaxies in this mass range are genuine spheroids. While below this mass both E and S0 classification are present, due to the morphological misclassification of inclined disks for genuine spheroidal ellipticals.

The round and massive objects that stand out from the shape distribution above $M_{\text{JAM}} \gtrsim 2 \times 10^{11} M_\odot$ are the prototypical slow rotators, which also stand out for their extremely low specific stellar angular momentum λ_R (Emsellem et al. 2007), as shown in Paper III. In the top panel of Fig. 8 we plot the distribution of the intrinsic λ_R , which was approximately obtained from the observed one given in Paper III by deprojecting the observed velocity to an edge-on view, using the best-fitting inclination from the JAM mod-

els. The plot shows a clear transition in the average λ_R of the galaxy population at the mass $M_{\text{JAM}} \gtrsim 2 \times 10^{11} M_\odot$. Above this mass essentially all ETGs are slow rotators, while below this mass the population is dominated by fast rotators, although some slow rotators are still present. The bottom panel of Fig. 8 plots the angular momentum parameter λ_R for the slow rotator ETGs only. The slow rotators are found to populate a rather narrow sequence on the MP, with a small range of Σ_e . This plot shows that not all slow rotators are the same. In fact there is a clear trend of decreasing rotation with increasing mass as noted in Paper III, or possibly a transition around $M_{\text{JAM}} \approx 10^{11} M_\odot$, which is similar to the characteristic mass defined by other observables. This confirms that the massive slow rotators above our characteristic mass are indeed special and may have formed differently from the rest. Although we tend to cover a smaller field with our kinematics for the largest galaxies, the qualitative difference in the kinematics for the most massive slow




Figure 7. Intrinsic shape and morphology. *Top Panels:* Same as in Fig. 1, with colours indicating the intrinsic ellipticity ϵ_{intr} of the galaxies outer regions, computed by de-projecting the observed ellipticity ϵ at large radii (Paper II) using the inclination inferred from the JAM models (Paper XV). Except again for the special region above $M_{\text{JAM}} \gtrsim 2 \times 10^{11} M_{\odot}$, all the rest of the ETGs have on average the flattening of disks $\epsilon \approx 0.6$ close to the result from the Monte Carlo approach of Paper III, which suggests fast rotators have a typical intrinsic ellipticity $\epsilon_{\text{intr}} \approx 0.7$ and the statistical inversion in Paper XXIII. *Bottom Panels:* Same as in the top panel, with colours indicating the galaxy morphology as quantified by the T-type parameter from HyperLeda. For $M_{\text{JAM}} \gtrsim 2 \times 10^{11} M_{\odot}$ galaxies are invariably classified as E ($T < -3.5$), while both E and S0 classifications are present at lower masses.

rotators is already clearly visible from the velocity fields (Paper II) and it is not due to the difference in the field coverage. A strong indication that indeed the most massive slow rotators are special also comes from the fact that they all appear to have cores, while the less massive ones are core-less (Krajinović et al. 2013, hereafter paper XXIV).

In summary the distribution of galaxy properties on the $(M_{\text{JAM}}, \sigma_e)$ and $(M_{\text{JAM}}, R_e^{\text{maj}})$ projections of the MP can be understood as due to a smooth variation in the bulge mass as already suggested in figure 4 of Paper I. The connection between bulge mass and galaxy properties, and the close link between ETGs and spiral galaxies is further illustrated in Fig. 9, which includes the location of the spiral galaxies of the ATLAS^{3D} parent sample (Paper I) together with the ETGs. The masses of spiral galaxies was approximately estimated assuming a fixed $M/L_K = 0.8 M_{\odot}/L_{\odot}$, which ensures agreement between M_{JAM} and the K -band luminosity at the lowest masses. The plot shows that ETGs properties

vary smoothly with those of spiral galaxies. Galaxies with negligible bulges are almost invariably star forming and classified as late spirals. Galaxies with intermediate bulges can still form stars and be classified as early-spirals, or can be fast rotator ETGs. But the galaxies with the most massive bulges, as indicated by their largest concentration or σ_e are invariably ETGs. They have the largest M/L (Fig. 1), the reddest colours, smallest $H\beta$ and lowest molecular gas fraction (Fig. 3), but are still flat in their outer parts, indicating they have disks (Fig. 7) and generally still rotate fast (Fig. 8). An exception are galaxies with masses $M_{\text{JAM}} \gtrsim 2 \times 10^{11} M_{\odot}$, which stand out for being nearly round, and non rotating. Above this characteristic mass essentially no spirals are present.

Also included in Fig. 9 is the approximate location of dwarf ETGs (spheroidals) and dwarf irregulars (Im) galaxies in the mass-size diagram. The dwarf ETGs parameters were taken from Gavazzi et al. (2005), Ferrarese et al. (2006b), Misgeld, Mieske & Hilker (2008) and Misgeld, Hilker & Mieske (2009), while the Im pa-




Figure 8. Galaxy stellar rotation on the MP projections. Same as in Fig. 1, with colours indicating the specific angular momentum parameter λ_R of Emsellem et al. (2007), as given in Paper III, deprojected to an edge-on view according to the JAM inclination (Paper XV). *Top Panels:* all ATLAS^{3D} galaxies are included. A clear transition in the mean galaxy rotation is evident above $M_{\text{JAM}} \gtrsim 2 \times 10^{11} M_{\odot}$, with the population becoming entirely dominated by slow rotators. *Bottom Panels:* Only the slow rotators are shown. They seem to define a sequence in the MP, and they show a decrease in the rotation especially above $M_{\text{JAM}} \gtrsim 2 \times 10^{11} M_{\odot}$, where only the slow rotators with the lowest λ_R are present.

parameters come from Kirby et al. (2008). The plot shows that the spheroidal galaxies lie along the continuation of the M-size relation for spiral galaxies and naturally connect to the region of fast rotator ETGs with the lowest M/L , young ages and small bulges. One should keep in mind that, while the galaxies parameter space is fully sampled above the mass selection limit $M_{\text{stars}} \gtrsim 6 \times 10^9 M_{\odot}$ of the ATLAS^{3D} survey, below this limit the data are heterogeneous. A much improved coverage of these parameters will soon be provided by the Next Generation Virgo Cluster Survey (Ferrarese et al. 2012) as shown in Ferrarese (2012).

4 SYSTEMATIC VARIATION OF THE IMF

Although the IMF variation could have been included among the other galaxy observables described in Section 3, we keep the IMF in a separate section for a more detailed coverage than for the other observables. In recent months there has been a large amount of interest

on the IMF variation. Before describing our new ATLAS^{3D} results, in the following section we provide a summary of this rapidly evolving field. To clarify the significance and robustness of the various recent claims, including our own, we place particular emphasis on the modelling assumptions that went into the various studies.

4.1 Summary of previous IMF results

4.1.1 IMF of the Milky Way

The stellar initial mass function (IMF) describes the distribution of stellar masses when the population formed. Nearly every aspect of galaxy formation studies requires a choice of the IMF to calculate predictions for galaxy observables (Kennicutt 1998). For this reason the IMF has been subjected to numerous investigations since the first determination more than half a century ago finding it has the form (Salpeter 1955)




Figure 9. The Mass-size distribution for dwarfs, spiral galaxies and ETGs. The ETGs of the ATLAS^{3D} sample are coloured according to their $(M/L)_{\text{JAM}}$ as in Fig. 1. This plot however also includes the approximate location of spiral galaxies in the ATLAS^{3D} parent sample, where masses are obtained from K -band luminosities, scaled to match M_{JAM} (a plot using luminosities rather than masses for both ETGs and spirals was shown in fig. 4 of Paper I). Early spirals (Sa-Sb: $T \leq 4$) are indicated by the black filled diamonds, while late spirals (Sc-Irr: $T > 4$) are shown with the filled green squares. Late spirals are larger than ETGs, while early spirals overlap with the ETGs with low M/L . There are essentially no spirals in the region of the oldest and reddest ETGs, which have the largest M/L . This illustrates the fact that the increase of the spheroid is required to make a galaxy old and red and consequently produce the largest M/L . No spirals are present also above the characteristic mass $M_{\text{JAM}} \gtrsim 2 \times 10^{11} M_{\odot}$. Also included in the plot is the approximate location of dwarf early-type galaxies (Sph: magenta filled up triangles. From Gavazzi et al. 2005, Ferrarese et al. 2006b Misgeld, Mieske & Hilker 2008 and Misgeld, Hilker & Mieske 2009) and dwarf irregulars (Irr: blue filled down triangles. From Kirby et al. 2008). These are only shown below the mass limit of the volume-limited ATLAS^{3D} survey (vertical dashed line). For reference the dash-dotted blue line shows the relation $(R_e/\text{kpc}) = 8 \times [M_{\text{stars}}/(10^{10} M_{\odot})]^{0.24}$.

$$\zeta(m) \propto m^x = m^{-2.3}, \quad (7)$$

for $m > 0.4M_{\odot}$, where m is the stellar mass.

In the Milky Way the IMF can be measured via direct stellar counts. Various determination in different environments lead to the finding of a remarkable universality in the shape of the IMF (Kroupa 2002), with the IMF being well described by the Salpeter power slope $x = -2.3$ for $m \geq 0.5M_{\odot}$ and a shallower one at smaller masses $x = -1.3$ for $m < 0.5M_{\odot}$ (Kroupa 2001). The Milky Way IMF can also be described by a log-normal distribution (Chabrier 2003), which has some theoretical justification. However the Kroupa and Chabrier IMFs are essentially indistinguishable from an observational point of view (Kroupa et al. 2013). Recent IMF results were reviewed by Bastian, Covey & Meyer (2010) who concluded no clear evidence existed for a non-universal IMF in our galaxy and among different galaxies.

4.1.2 IMF from ionized gas emission or redshift evolution

In external galaxies individual stars cannot yet be resolved down to sufficiently small masses for IMF studies. Some indirect constraints on the slope of the IMF above $\sim 1M_{\odot}$ can be obtained by combining observations of the $H\alpha$ equivalent widths, which is related to the number of ionizing photons, with the galaxy colour, which is

a function of the galaxy stellar population (Kennicutt 1998). Using this technique Hoversten & Glazebrook (2008) inferred a variation of the IMF with more massive galaxies having a more top-heavy IMF than Salpeter, a result confirmed under the same assumptions (Gunawardhana et al. 2011) by the GAMA survey (Driver et al. 2011). A similar result was found using the ratio between the $H\alpha$ and ultraviolet fluxes by Meurer et al. (2009), who further pointed out that surface density is more important than mass in driving the IMF trend.

Constraints on the IMF can also be obtained by comparing the local ($z = 0$) stellar mass density to the integral of the cosmic star formation history. Wilkins et al. (2008) and Davé (2008) find that the local stellar mass density is lower than the value obtained from integrating the cosmic star formation history (SFH), assuming that all the stars formed with a Salpeter IMF. They propose a time-variable IMF to reconcile the observations.

Alternative constraints on the IMF can be obtained by studying the evolution of the M/L normalization in samples of galaxies as a function of redshift, for example from the evolution of the tilt and normalization of the Fundamental Plane with redshift (Renzini & Ciotti 1993). From these constraints and from the metal abundance in the clusters interstellar medium Renzini (2005) concluded that elliptical galaxies as a class must have an IMF close to Salpeter

for stellar masses $m > 1M_{\odot}$, while the IMF must flatten at lower masses, similar to the Kroupa IMF.

A related approach was employed by van Dokkum (2008) who used the ratio of luminosity evolution to colour evolution of massive galaxies in clusters to constrain the IMF. He originally concluded that the IMF must be top heavy at high redshift in apparent agreement with some theoretical model predictions (Zepf & Silk 1996; Chiosi et al. 1998). However this result was recently revised with better data and models to conclude that the IMF above $m > 1M_{\odot}$ is actually not inconsistent with a Salpeter slope (van Dokkum & Conroy 2012).

4.1.3 *Lensing or dynamics IMF constraints for external spirals*

A more direct approach to constraining the variation of the IMF in galaxies consists of measuring the mass of the stars, and comparing this with the predictions of stellar population models. It is important to stress however that this method does not measure the shape of the IMF directly, but only its overall mass normalization. The method can only verify whether the stellar mass in a certain region within a galaxy is consistent with that due to a stellar population with a certain assumed IMF. Moreover these methods do not measure the same IMF one can infer via direct stellar counts in star clusters. They measure instead the stellar mass distribution due to the superposition of the IMFs from a large number of star clusters in the galaxy as well as the IMFs coming from the accretion of satellite galaxies during the galaxy hierarchical growth. As pointed out by Kroupa & Weidner (2003), even if the IMF was universal in every cluster in a galaxy, the integrated galactic IMF (IGIMF) would be different from the universal one, due to the non-uniform mass distribution of star clusters (see Kroupa et al. 2013, for an in-depth discussion). Hierarchical evolution complicates the picture even further. For these reason, when we state e. g. that “a galaxy is consistent with the Salpeter IMF”, we simply imply, consistently with all previous similar studies, that “the galaxy has the same stellar M/L of a stellar population with the given age and metallicity (and abundances), and the Salpeter IMF”.

The first convincing constraint on the IMF of external galaxies was obtained for a sample of 21 spiral galaxies, using the kinematics of the gas. Bell & de Jong (2001) concluded that *if the IMF is universal*, a sensible assumption at the time, it cannot have the Salpeter form, but it must be lighter, consistently with Kroupa-type. The result was later confirmed, still using gas kinematics, for a sample of 34 bright spiral galaxies by Kassin, de Jong & Weiner (2006).

The need for a light IMF was inferred for the Einstein Cross spiral galaxy gravitational lens system (Huchra et al. 1985) by both van de Ven et al. (2010) and Ferreras et al. (2010), and for another spiral lens galaxy by Suyu et al. (2012). As part of the DiskMass Survey (Bershady et al. 2010), which obtained integral-field stellar kinematics of a sample of 30 spiral galaxies, a light IMF was indirectly confirmed from the sub-maximality of the disks (Bershady et al. 2011). As part of the SWELLS survey of spiral lens galaxies (Treu et al. 2011) the need for a light IMF for most spiral galaxies was also found from the analysis of 20 strong gravitational lens (Brewer et al. 2012).

An inconsistency of the Salpeter IMF normalization for low mass galaxies was also inferred by Dutton et al. (2011b), using simple galaxy models with a bulge and a disk, trying to reproduce global trends for a large sample of galaxies extracted from the SDSS.

In summary there is a good agreement on the fact that spiral galaxies as a class must have a normalization lighter than Salpeter

and similar to Kroupa/Chabrier. This is a robust result due to the fact that a Salpeter IMF would over-predict the total mass in the galaxy centres. It is unclear whether some spiral galaxies have a Salpeter IMFs, and of course whether the IMF varies within the galaxies themselves, as claimed by Dutton et al. (2013).

4.1.4 *Lensing or dynamics IMF constraints on early-types*

An early attempt at constraining the IMF of 21 elliptical galaxies, using detailed spherical dynamical models including dark matter, found a general consistency between the stellar $(M/L)_{\text{stars}}$ and the one of the population $(M/L)_{\text{pop}}$, using the Kroupa IMF, but could not accurately constrain the IMF normalization due to large observational errors (Gerhard et al. 2001). A large study of SDSS elliptical galaxies using fixed spherical Hernquist galaxy models with dark halos (Padmanabhan et al. 2004) found a mass excess over the predictions of stellar population models with a fixed IMF, increasing with luminosity. This was interpreted as a increase of the dark matter fraction. A similar conclusion was reached by Zaritsky, Gonzalez & Zabludoff (2006) while studying the fundamental manifold. A caveat of those studies was the use of homologous stellar profiles or approximate assumptions to study systematic variations in the heterogeneous galaxy population.

To overcome these limitations, as part of the SAURON project (de Zeeuw et al. 2002), we constructed self-consistent axisymmetric models reproducing in detail both the photometry and the state-of-the-art integral-field stellar kinematics data (Emsellem et al. 2004) for 25 early-type galaxies (Cappellari et al. 2006). The assumption that mass follows light is not accurate at large radii, but it is a good assumption for the region where kinematics is available ($\lesssim 1R_e$), and provides accurate measurements of the total (luminous plus dark) M/L within a sphere of radius $r \approx R_e$ (see Paper XV). The resulting improvement in the M/L accuracy and the removal of systematic biases allowed us to strongly confirm that “the total and stellar M/L clearly do not follow a one-to-one relation. Dark matter is needed to explain the differences in M/L (if the IMF is not varying)” (Cappellari et al. 2006, see fig. 17 there). Although dark matter seemed, at the time, still a more natural explanation of our observations, if the measured trends are re-interpreted as an IMF variation, they would imply an IMF heavier than Salpeter for some of the oldest objects. That study also concluded that *if the IMF is universal* it must have a mass normalization as low as Kroupa IMF, consistently with the results for spiral galaxies, otherwise the stellar mass would overpredict the total one for a number of galaxies. A similar finding was obtained by Ferreras, Saha & Burles (2008) using gravitational lensing of 9 elliptical galaxies extracted from the SLACS survey (Bolton et al. 2006).

A number of subsequent studies used approximate models reproducing the velocity dispersion of large galaxy samples to study dark matter and IMF variations in galaxies. Tortora et al. (2009) used spherical isotropic models with Sersic (1968) profiles to reproduce the large velocity dispersion compilation by Prugniel & Simien (1996). Graves & Faber (2010) used the Fundamental Plane of early-type galaxies from SDSS data. Schulz, Mandelbaum & Padmanabhan (2010) combined weak lensing measurements in the outer parts to SDSS velocity dispersion determination, using spherical Jeans models with a Hernquist profile. All three studies confirmed the existence of a mass excess which increases with mass. They all preferred a dark matter trend to explain the observations, although they could not exclude the IMF variation alternative.

Grillo et al. (2009) compared stellar population masses, derived from multicolour photometry, to the total masses, inside the

Einstein radius of the lenses, published by the SLACS team (Bolton et al. 2008a). He assumed for all galaxies the average dark matter fraction determined for some of the galaxies by Koopmans et al. (2006) and Gavazzi et al. (2007), which were based on spherical models assuming a single power-law total mass and a Hernquist (1990) (or Jaffe 1983) stellar profile. From this comparison Grillo et al. (2009) concluded that elliptical galaxies prefer a Salpeter rather than Kroupa/Chabrier IMF. The same result was found when comparing stellar dynamical masses for galaxies in the Coma cluster by Thomas et al. (2009), which includes dark matter halos with fixed NFW profiles, to stellar population masses (Grillo & Gobat 2010). However, given that the dynamical masses were determined assuming a fixed NFW profile, the conclusion depended on the correctness of that assumption. A recent reanalysis of the 16 Coma galaxies by Thomas et al. (2011), still assuming a NFW dark halo profile, interpreted the M/L excess in massive galaxies as more likely due to dark matter, although their results are not inconsistent with a Salpeter IMF instead. The same conclusions were reached by Wegner et al. (2012) for 8 galaxies in Abell 262.

The SLACS team analysed their data assuming all galaxies in their sample can be approximated by homologous spherical and isotropic systems with a Hernquist (1990) (or Jaffe 1983) profile. In Treu et al. (2010) they further assumed a spherical NFW profile for the dark matter, with fixed slope and only mass as free parameter. Given that the enclosed total (luminous plus dark) mass inside the Einstein radius is essentially fixed by the lens geometry, under these assumptions the central velocity dispersion is a unique function of the profile of the total mass distribution, which in this case is defined by a single parameter: the ratio between the stellar and dark matter components. Comparing the stellar mass from the lens model to the one from population, based on multicolour photometry (Auger et al. 2009), they found that the data prefer a Salpeter-like normalization of the IMF. Auger et al. (2010b) used the SLACS data and the same spherical Hernquist models but, instead of allowing for free halos for different galaxies, they assumed the same IMF and the same halos, following the trends predicted by the abundance matching technique (Moster et al. 2010), for the entire ensemble population. They still assumed NFW halos but explored both adiabatically contracted (Gnedin et al. 2004) and not contracted halos. They concluded that the data favour a Salpeter-like normalization of the IMF over a lighter Kroupa/Chabrier form.

Although ground-breaking, the analyses of the SLACS data depended on some non-obvious assumptions. Grillo et al. (2009) had assumed a fixed dark matter fraction, as well as a power-law form for the total mass. Treu et al. (2010) result depended on the assumption of a fixed NFW halo slope, as the authors acknowledged concluding that “the degeneracy between the two [IMF or dark matter] interpretations cannot be broken without additional information, the data imply that massive early-type galaxies cannot have both a universal IMF and universal dark matter halos”, in agreement with previous dynamical analyses. Auger et al. (2010b) result depended on assuming the halo mass from the abundance matching techniques. This led the authors to conclude that “better constraints on the star formation efficiency must be obtained from the data in order to draw definitive conclusions about the role of a mass-dependent IMF relative to CDM halo contraction”.

Other independent studies of the same SLACS data did not exclude an IMF variation, but did not confirm the need for a Salpeter IMF to explain the observations. Tortora et al. (2010) reached this conclusion using spherical Hernquist isotropic models like Treu et al. (2010) and Auger et al. (2010b), but could not explain the reason for the disagreement. A similar study was also performed

by Deason et al. (2012). They showed that the trend in the galaxies enclosed masses can be explained by a toy model similar to the one adopted by Auger et al. (2010b), but with a universal Kroupa IMF. Deason et al. (2012) study however did not fit the galaxies stellar velocity dispersion, which could be a reason for the difference in the results. Barnabè et al. (2011) re-analysed 16 SLACS galaxies using axisymmetric, rather than spherical, dynamical models, and describing the galaxy images in detail, rather than assuming fixed Hernquist profiles. Their models are fitted to integral-field stellar kinematics, instead of a single velocity dispersion. They find that the data are consistent with both a Kroupa or a Salpeter IMF, in agreement with the early study by Ferreras, Saha & Burles (2008).

An additional limitation all previous analyses of the full SLACS sample was the assumption that all galaxies in their sample could be described by homologous spherical Hernquist distributions. These models ignore known systematic variations of galaxy morphology with mass (e.g. Caon, Capaccioli & D’Onofrio 1993; Kormendy et al. 2009, see also Fig. 6). Moreover real early-type galaxies, even in the mass range of the SLACS survey, are dominated by fast rotating disks (Paper II, Paper III), some of which are clearly visible in the SLACS photometry. Disk galaxies are not well described by spherical single-component models. It is unclear whether galaxy models that do not reproduce neither the kinematics nor the photometry of the real galaxies under study can be trusted at the $\sim 10\%$ level that is required for IMF studies. Possible systematic biases in the results are however difficult to estimate. A reanalysis of the excellent SLACS dataset using more realistic and flexible models would seem the best way to clarify the situation and provide a consistent picture. This approach has already been demonstrated by Barnabè et al. (2012) for a single galaxy using the JAM method (Cappellari 2008) in combination with the lensing analysis. One should finally consider that lensing studies measure the mass within cylinders along the line-of-sight. The recovered stellar mass depends on both the assumed dark matter profile in the centre, as well as at large radii (Dutton et al. 2011a).

A simple assumption on the dark matter content of early-type galaxies was made by Dutton, Mendel & Simard (2012). They selected the most dense galaxies from a large sample of SDSS galaxies and constructed spherical isotropic models to reproduce their stellar velocity dispersion. They assumed the total dynamical M/L of this set of galaxies is the same as the stellar one. Comparing the inferred mass to the one from stellar population they concluded the galaxies require on average a Salpeter IMF normalization. This study still depends on the spherical approximation. Moreover it is unclear to what accuracy the zero dark matter assumption is verified even for the densest galaxies. Nonetheless this study confirms that, unlike spiral galaxies, the densest early-type galaxies are not inconsistent with a Salpeter IMF.

In summary, the two result on which nearly all previously discussed studies agree are: (i) the *total* M/L in the central regions of galaxies ($r \lesssim R_e$) does not follow the M/L inferred assuming a universal IMF; (ii) less massive galaxies, especially spiral ones, *require* an IMF normalization lighter than Salpeter and consistent with Kroupa-like, while more massive and dense ones allow for a Salpeter IMF. Indications were found for the Salpeter IMF to be actually *required* for massive ellipticals (Grillo et al. 2009; Treu et al. 2010; Auger et al. 2010b; Dutton, Mendel & Simard 2012). However these conclusions depend on assuming the knowledge of either the dark matter fraction or the slope, or the mass of the dark halo. An exception is the recent work by Sonnenfeld et al. (2012), which modelled a rare elliptical with two concentric Einstein rings, and concluded for a Salpeter IMF. More similar ob-

jects would be needed to draw solid conclusions. Not all studies agreed on the requirement for a Salpeter IMF, even when analysing the same SLACS data (Ferreras, Saha & Bures 2008; Tortora et al. 2010; Barnabè et al. 2011; Deason et al. 2012). It seems that this situation could be resolved by the use of more realistic models.

4.1.5 Systematic IMF variation in galaxies

A breakthrough in IMF studies was provided by the work by van Dokkum & Conroy (2010), further strengthened in van Dokkum & Conroy (2011). Contrary to the previously described set of dynamical and lensing works, they looked for evidences of IMF variation in subtle IMF-sensitive spectral features of the near-infrared region of galaxy spectra, in particular the Wing-Ford band. In this way their study did not suffer from the dependency on the halo assumptions. Although the spectral technique has been around for decades (e.g. Spinrad & Taylor 1971), only the availability of reliable stellar population models has made the approach sufficiently accurate for IMF studies (Schivon, Barbuy & Bruzual A. 2000; Cenarro et al. 2003). van Dokkum & Conroy (2010) analysed the IMF of eight massive ellipticals, from stacked spectra. They used new population models that allow for a variation of the detailed abundance patterns of the stars (Conroy & van Dokkum 2012a) to distinguish abundance from IMF variations. They concluded that the observed spectra required a bottom-heavy, dwarf-rich IMF. Combining their finding with the previous results on the IMF of spiral galaxies they tentatively concluded that “Taken at face value, our results imply that the form of the IMF is not universal but depends on the prevailing physical conditions: Kroupa-like in quiet, star-forming disks and dwarf-rich in the progenitors of massive elliptical galaxies”.

Our relatively large and well-selected ATLAS^{3D} sample and high-quality integral-field stellar kinematics appeared well suited to resolve the halo degeneracies of previous dynamical and lensing analyses and test the claims from the spectral analysis. Contrary to previous large studies, we adopted an axisymmetric modelling method which describes in detail both the individual galaxy images using the MGE technique (Emsellem, Monnet & Bacon 1994; Cappellari 2002) and the richness of our two-dimensional kinematics (Cappellari 2008), thus avoiding the possible biases of previous more approximate approaches. For the first time, thanks to the tighter constraints to the models provided by the two-dimensional data, our study could leave *both* the halo slope and its mass as free parameters. The halo slope is allowed to vary in a range which includes both the flat inner halos predicted by halo expansion models (e.g. Governato et al. 2010), and the steep ones predicted by adiabatic halo contraction (e.g. Gnedin et al. 2004). Our models also explicitly include the galaxy inclination and anisotropy as free parameters, although the latter is still assumed to be constant in the region where we have data. The parameters are estimated in a Bayesian framework with a maximum ignorance (flat) prior on the parameters.

We showed that even in this relatively general case the models required a variation of the IMF to reproduce the data (Cappellari et al. 2012), unless there are major flaws affecting all available stellar population models. We additionally tested a variety of sensible modelling assumptions on the halo, some of which had already been employed by previous studies. However this time our models accurately described the photometry and kinematics of the real galaxies. We confirmed that a non-universal IMF is required under all those halo assumption. The similarity is due to the fact that in all cases dark matter contributes just about 10-20% to the total mass within

$1R_e$, so that it cannot explain an observed M/L excess of up to a factor 2-3.

Our ATLAS^{3D} sample includes a larger range of masses than previous lensing studies. Not only we could quantify the IMF normalization of early-type galaxies as a class, but we showed that the IMF normalization varies systematically within the early-type galaxy population, as a function of the stellar $(M/L)_{\text{stars}}$. The IMF was found consistent with Kroupa/Chabrier-like at low $(M/L)_{\text{stars}}$ and heavier than Salpeter at large $(M/L)_{\text{stars}}$. This finding bridges the gap between the Kroupa-like IMF determinations in spirals and the evidence for a Salpeter-like or heavier in early-type galaxies. This should be expected given the parallelism and continuity in physical parameters between early-type and spiral galaxies as emphasized in our ‘comb’ morphological diagram (Paper VII). Early-type galaxies at low $(M/L)_{\text{stars}}$ in fact completely overlap with the region populated by spiral galaxies on the mass-size projection of the MP (Fig. 9). Our result also reconciled the apparent disagreement between studies claiming that early-type galaxies include cases with Kroupa/Chabrier normalization and those claiming that Salpeter is required.

Although, unsurprisingly, the cleanest IMF trend in Cappellari et al. (2012) was obtained by simply comparing the dynamical and population M/L , a trend with velocity dispersion σ_e was also presented. This is a natural consequence of existence of the $(M/L) - \sigma$ correlation (Cappellari et al. 2006; van der Marel & van Dokkum 2007). In particular Cappellari et al. (2012) (their fig. 2) showed an IMF trend, with significant scatter, going from a Kroupa/Chabrier to a Salpeter IMF within the range $\log_{10}(\sigma_e/\text{km s}^{-1}) \approx 1.9 - 2.5$, with a gradual variation in between.

A flurry of papers have appeared in subsequent months from independent groups, all in agreement on the existence of a systematic IMF variation in galaxies. Spiniello et al. (2012) used the models of Conroy & van Dokkum (2012a) in combination with SDSS spectra and inferred a variation of the IMF from sodium and titanium-oxide absorptions, which correlate with velocity dispersion (also illustrated by Zhu, Blanton & Moustakas 2010). A similar result was inferred by Ferreras et al. (2013), also from SDSS spectra, but using the population models by Vazdekis et al. (2012). Smith, Lucey & Carter (2012) studied the Wing-Ford absorption for galaxies in the Coma Cluster. They find that their galaxy spectra are best reproduced by a Salpeter IMF and detect a weak IMF trend with metallicity, but do not find an IMF trend with velocity dispersion. Conroy & van Dokkum (2012b) detected a systematic trend of the IMF, versus either σ or metallicity, using both optical and near-infrared spectral features. A key difference from other spectral studies is that they were able to rule out, in a Bayesian framework, a variation of element abundances as the origin of the empirical trends.

Dutton et al. (2012) showed that it is sufficient to assume the knowledge of the halo mass and fit galaxy scaling relations via approximate spherical galaxy models, to infer a systematic variation of the IMF. Similarly Tortora, Romanowsky & Napolitano (2013) tested a set of different assumption on the dark halo, using approximate spherical Sersic (1968) galaxy models to fit the SDSS velocity dispersion of a large galaxy sample. Some of the halo assumptions were also included in Cappellari et al. (2012) with more detailed dynamical models, producing consistent results. Although these results still depend on simple approximations and on assumption on the halo, they illustrate that many different halo assumptions, motivated by theoretical models, all lead to the same conclusion of a systematic IMF variation.

In summary, after years of debate on whether the IMF or the

dark matter were responsible for the observed disagreement between stellar population and dynamical or lensing indicators, a consensus seems to have emerged, from dynamical, lensing and spectral arguments, on a systematic IMF variation with the IMF becoming heavier with M/L , mass or velocity dispersion. It is time to investigate IMF trends against other observables, which we do in the following section.

4.2 IMF variation on the Mass Plane

In Cappellari et al. (2012) we presented a systematic trend between the IMF and the stellar $(M/L)_{\text{stars}}$ derived using axisymmetric JAM models including a dark halo. The IMF was parametrised by the IMF mismatch parameter $\alpha \equiv (M/L)_{\text{stars}}/(M/L)_{\text{Salp}}$ (in the notation of Treu et al. 2010), where $(M/L)_{\text{Salp}}$ was derived from full spectral fitting as summarized in Section 2.3. Different population approaches were also tested there, producing less clean trends but consistent results. The $(M/L)_{\text{Salp}}$ normalization depends on the adopted lower and upper mass cut-offs for the IMF in the population models. The models we use (Vazdekis et al. 2012) adopt standard lower and upper mass cut-offs for the IMF of 0.1 and 100 M_{\odot} , respectively. $(M/L)_{\text{Salp}}$ further depends on whether the gas lost by the stars during the early stages of their evolution is retained in the central regions we observe or is recycled into stars or expelled at large radii. Evidence suggests the gas is not retained in significant quantities within $1R_e$, neither in ionized or hot X-ray emission form (Sarzi et al. 2010), nor as a cold component (Paper IV). If all the gas was retained it would increase $(M/L)_{\text{Salp}}$ by up to 30–40% (Maraston 2005), making population M/L overpredicting total dynamical M/L even for a light Kroupa/Chabrier IMF, for a number of galaxies.

A careful determination of the population M/L for a fixed IMF was recently provided for 35 galaxies in our sample by Conroy & van Dokkum (2012b). They employed different models (Conroy & van Dokkum 2012a) from the ones we adopted (Vazdekis et al. 2012) and used them to fit an independent set of spectra, spanning a larger wavelength range than the SAURON ones. In Fig. 10 we compare the two determinations of M/L (allowing for an arbitrary offset in the absolute normalization, which can differ by up to 10% between different authors). Assuming all measurements have comparable errors, the observed scatter between the 35 values is consistent with an error of just 6% in $(M/L)_{\text{Salp}}$. This error is consistent with the error of 7% predicted by Gallazzi & Bell (2009) when optimal spectra information is available. This gives confidence that our population $(M/L)_{\text{Salp}}$ are robust and any trend we observe is not due to the details of our population modelling approach.

Cappellari et al. (2012) showed that consistent IMF trends are found for a variety of different assumptions on the dark halo, as well as for the most general one (model D) which leaves both the halo slope and mass as free parameters, with only an upper limit on the halo slope, derived from model predictions of halo contraction (Gnedin et al. 2011) (see Paper XV for a description of the models). The similarity in $(M/L)_{\text{stars}}$ for the different models is due to the fact that in all cases the data allow for a small fraction of dark matter within the region where the kinematics are available ($\sim 1R_e$), with the most general model implying a median dark matter fractions of just 10%. Given the similarity of the different approaches, here we adopt as reference the $(M/L)_{\text{stars}}$ values obtained with a NFW halo, with halo mass as free parameter (model B). This choice also makes it easy to compare our results with those of others authors that make the same assumption.

In Fig. 11 we show a different rendition of the similar




Figure 10. Testing the accuracy of $(M/L_r)_{\text{Salp}}$ against the completely independent determination of $(M/L_K)_{\text{MW}}$ from Conroy & van Dokkum (2012b), which use a different population modelling software and different spectra with a longer wavelength range than the SAURON ones. Assuming all measurements have comparable errors, the observed scatter between the 35 values is consistent with a 1σ error $\Delta/\sqrt{2}$ of 6% in $(M/L_r)_{\text{Salp}}$. The $(M/L_K)_{\text{MW}}$ values were multiplied by a constant factor to approximately account for the differences in the photometric band and IMF.

fig. 2b of Cappellari et al. (2012). Here we plot $(M/L)_{\text{Salp}}$ versus $(M/L)_{\text{stars}}$. We still exclude galaxies with very young stellar populations, selected as having stellar absorption line strength index $H\beta > 2.3 \text{ \AA}$. We found that those galaxies have strong gradient in the stellar population and this breaks our approximate assumption of a constant $(M/L)_{\text{stars}}$ within the region where we have kinematics, making both $(M/L)_{\text{stars}}$ and $(M/L)_{\text{Salp}}$ inaccurate and ill-defined. Different diagonal lines illustrate the expected trends if the IMF was universal or either the Chabrier, Kroupa or Salpeter ($\zeta(m) \propto m^{-2.3}$) forms. Also shown is the expected trend for two IMFs heavier than Salpeter: a top-heavy one, dominated by stellar remnants ($\zeta(m) \propto m^{-1.5}$), and a bottom-heavy one, dominated by dwarfs ($\zeta(m) \propto m^{-2.8}$). This figure clearly illustrates the fact that the two indicators of M/L do not follow a one-to-one relation, but deviate systematically, with the IMF being consistent on average with Kroupa/Chabrier at the lowest $(M/L)_{\text{stars}}$ and with Salpeter or heavier at the largest $(M/L)_{\text{stars}}$, as already concluded in Cappellari et al. (2012). Another way to interpret this plot is by noting that galaxies with the largest $(M/L)_{\text{Salp}}$, which are characterized by the oldest populations, have a Salpeter or heavier IMF, while those with the lowest $(M/L)_{\text{Salp}}$, which have younger populations, have on average a Kroupa/Chabrier IMF. Detailed trends of the IMF with other population indicators will be presented in McDermid et al. (in preparation).

As in the Cappellari et al. (2012) we also show with colours the galaxies velocity dispersion. However, to emphasize the trend, here we apply the same two-dimensional LOESS smoothing approach (Cleveland & Devlin 1988) introduced in Section 3, instead of showing the individual σ_e values. One can still see the trend for the IMF to vary between Kroupa/Chabrier to Salpeter, albeit with




Figure 11. The systematic variation of the IMF. The $(M/L_r)_{\text{stars}}$ of the stellar component, determined via dynamical models, is compared to the $(M/L_r)_{\text{Salp}}$ determined from spectral fitting using stellar population models and assuming for reference a fixed Salpeter IMF. The colours of the symbols show the galaxies σ_e , which was LOESS smoothed in two-dimension to emphasize the trends. The plot only includes the subsets of ATLAS^{3D} galaxies with $H\beta < 2.3 \text{ \AA}$, for which $(M/L)_{\text{Salp}}$ is sufficiently reliable. Diagonal lines illustrate the expected trends if the IMF was universal with either the Chabrier (red dash-dotted line), Kroupa (green dashed line) or Salpeter ($\zeta(m) \propto m^{-2.3}$, solid magenta line) forms. Also shown is the expected trend for two IMFs heavier than Salpeter (blue dotted line): a top-heavy one, dominated by stellar remnants ($\zeta(m) \propto m^{-1.5}$), and a bottom-heavy one, dominated by dwarfs ($\zeta(m) \propto m^{-2.8}$). The black solid curve is a LOESS smoothed version of the data. A clear systematic trend is evident, with the IMF being closer to Kroupa/Chabrier at the lowest M/L , which also have the lowest σ_e , and closer to Salpeter or heavier at the largest M/L or largest σ_e . A different rendition of this plot was presented in fig. 2b of Cappellari et al. (2012).

large scatter, within the interval $\log_{10}(\sigma_e/\text{km s}^{-1}) \approx 1.9 - 2.5$, with a smooth variation in between. This trend will be more precisely quantified in Section 4.3.

In Fig. 12 we show the variation of the IMF mismatch parameter $(M/L)_{\text{stars}}/(M/L)_{\text{Salp}}$ on the $(M_{\text{JAM}}, \sigma_e)$ and (M_{JAM}, R_e) projections of the MP. The trends in IMF are necessarily noisier than the ones in $(M/L)_{\text{stars}}$, as here the errors in both M/L combine. However the structure in this figure closely resembles the one in the previous Fig. 1, 3 and the top panel of Fig. 6. The systematic variation in the IMF follows on average the variations in the total $(M/L)_{\text{JAM}}$ and its corresponding stellar population indicators $H\beta$ and galaxy colour, as well as the molecular gas fraction. Like the other quantities, also the IMF appears to roughly follow lines of constant σ_e on the MP, which we showed is tracing the bulge mass at given galaxy size and mass. In both projections there is indication for some extra substructure, with IMF variation at constant σ_e . Moreover the proximity to the ZOE (thick red line) seems to be

an even better indicator of IMF than σ_e (blue dashed lines). The (M, R_e) projection also makes clear why one should expect spirals to have on average light IMF: spirals populate the empty zone above our galaxies (Fig. 9 here and fig. 4 in Paper I) and overlap with the distribution of ETGs with the Kroupa/Chabrier IMF. However our plot suggests that, at fixed galaxy stellar (or M_{JAM}) mass and below $M_{\text{JAM}} \lesssim 10^{11} M_{\odot}$, the bulge mass (as traced by σ_e) is the main driver of the IMF variation, rather than morphological type, and one may expect IMF variations within the spiral population as well. Our finding explains why Dutton, Mendel & Simard (2012) inferred a Salpeter or heavier IMF when selecting the smallest (or densest) galaxies at fixed mass, and assuming no dark matter. Dark matter appears to provide a small contribution to the M/L for our entire sample and not just the smallest ones (Paper XV), but the densest galaxies are precisely the ones with the heaviest IMF. A recent claim has been made for a Salpeter IMF in the bulge of 5 massive spirals (Dutton et al. 2013), which would go in the suggested direction. However the IMF is currently already difficult to infer when the M/L can be assumed to be spatially constant and the stellar population homogeneous. Relaxing these assumptions makes the results more uncertain unless very good population data from galaxy spectra and accurate non-parametric models for the photometry are employed.

4.3 IMF versus σ_e correlation

In the previous section we illustrated the systematic IMF trends already presented in Cappellari et al. (2012) and we additionally presented the variation of the IMF on the projections of the MP. We pointed out that, like the dynamical M/L , its population indicators, colour and $H\beta$, the cold gas fraction and the galaxy concentration which traces to the bulge mass, also the IMF broadly follows lines of nearly constant σ_e on the $(M_{\text{JAM}}, \sigma_e)$ MP projection and consequently $R_e^{\text{maj}} \propto M_{\text{JAM}}$ on the $(M_{\text{JAM}}, R_e^{\text{maj}})$ projection. Interestingly a ‘conspiracy’ between the IMF variation and other galaxies properties is required for galaxy scaling relations to still have a small scatter (Renzini & Ciotti 1993).

In Fig. 13 we present a direct correlation between the logarithm of the IMF mismatch parameter $(M/L)_{\text{stars}}/(M/L)_{\text{Salp}}$ and the logarithm of σ_e , using the robust linear fitting routine `LTS_LINEFIT` presented⁵ in (Paper XV), which explicitly allows and fits for intrinsic scatter and removes outliers. Like earlier we exclude galaxies with very young stellar populations, selected having stellar absorption line strength index $H\beta > 2.3 \text{ \AA}$, leading to a sample of 223 out of 260 galaxies. In the fits we quadratically co-added JAM modelling errors of 6% (Paper XV) plus distance errors (Paper I) for $(M/L)_{\text{stars}}$ plus population models errors of 6% for $(M/L)_{\text{Salp}}$ (Fig. 10) plus 10% errors for our photometry (Paper XXI). We present fits for (i) the full sample, (ii) for the subsample of galaxies with SBF distances (mostly from Tonry et al. 2001 and Mei et al. 2007: see Paper I) and (iii) for galaxies with $M_{\text{JAM}} < 2 \times 10^{11} M_{\odot}$, to eliminate the galaxies above the characteristic mass where the IMF trend, as well as other galaxy properties, seems to change (see Fig. 7, 8 and 12). All three relations have a comparable observed scatter $\Delta \approx 0.10$ dex (26%) in $(M/L)_{\text{stars}}/(M/L)_{\text{Salp}}$ and imply a significant intrinsic scatter of about 20%. When the same outliers are removed, we verified that indistinguishable values for the parameter, their errors and the inferred intrinsic scatter are obtained with the Bayesian approach of

⁵ Available from <http://purl.org/cappellari/idl>




Figure 12. Variation of the IMF on the MP projections. Same as in Fig. 1 for the variation of the IMF mismatch parameter $(M/L)_{\text{stars}}/(M/L)_{\text{Salp}}$, which measure the ratio between the stellar M/L from dynamical models and the one from population models, with a fixed Salpeter IMF for reference. Like in Fig. 11 only galaxies with $H\beta < 2.3 \text{ \AA}$ are included. The trend follows lines of roughly constant σ_e , or even better, follows lines parallel to the ZOE (thick red line) above its break. Like in other diagrams the trend looks different at large masses $M_{\text{JAM}} \gtrsim 10^{11} M_{\odot}$, suggesting these objects are different. The heaviest IMF (red colour) is not reached by the most massive galaxies, but by the fast rotators with the biggest bulges.

Kelly (2007), as implemented in his IDL routine `LINMIX_ERR`. Our three fits provide consistent values for the best-fitting slopes, within the errors, and nearly consistent normalizations. The fitted relation has the form

$$\log_{10} \frac{(M/L)_{\text{stars}}}{(M/L)_{\text{Salp}}} = a + b \times \log_{10} \frac{\sigma_e}{130 \text{ km s}^{-1}}, \quad (8)$$

and our preferred values (bottom panel of Fig. 13) have best fitting parameters and formal errors $a = -0.12 \pm 0.01$ and $b = 0.35 \pm 0.06$ (parameters and errors for the other fits are given inside the figures). The observed trend of IMF with σ appears to account for about half of the total trend in the $(M/L) - \sigma_e$ relation $(M/L)_{\text{JAM}} \propto \sigma_e^{0.72}$ (see Paper XV), the remaining one being due to stellar population variations.

Our trend implies a transition of the mean IMF from Kroupa to Salpeter in the interval $\log_{10}(\sigma_e/\text{km s}^{-1}) \approx 1.9 - 2.5$ (or $\sigma_e \approx 90 - 290 \text{ km s}^{-1}$), with a smooth variation in between, consistently with what can be seen in Cappellari et al. (2012) and in Fig. 11. The fact that this trend is slightly weaker than the one implied by Fig. 11 seems to confirm the intrinsic differences in the IMF of individual galaxies. However, part of this difference could also be explained if the distance errors were underestimated. One way to address this issue would be to repeat our analysis for galaxies in a clusters at intermediate distance like the Coma cluster, for which relative distance errors can be neglected.

Our slope is a factor ≈ 3.5 smaller than the “tentative” trend reported in Treu et al. (2010) for the same quantities, and making the same assumption for the dark halo. A reason for this large difference must be due to the fact that their sample only included galaxies with $\sigma \gtrsim 200 \text{ km s}^{-1}$ ($\log_{10}(\sigma/\text{km s}^{-1}) \gtrsim 2.3$). Our sample is too small to reliably study trend in IMF for the galaxies above that σ_e , but their reported trend would exceed the slope of the $(M/L) - \sigma$ relation (Paper XV), making it difficult to explain. Their steep inferred trend may then be due to their use of spherical and homologous models for all the galaxies, which may introduce systematic trends. Moreover they used single stellar population M/L based on colours is expected to be less reliable than our determinations

based on spectra (Gallazzi & Bell 2009). A reanalysis of the unique SLACS dataset seems required to clarify this issue. Our trend with σ_e is also smaller than the one reported by Ferreras et al. (2013) from spectral analysis, who find a rapid change from a Kroupa to a Salpeter IMF in the narrow interval $\sigma \approx 150 - 200 \text{ km s}^{-1}$. Our relation is not inconsistent with the values presented in Conroy & van Dokkum (2012b) also from spectral analysis, or with the result reported by Spiniello et al. (2012), with the IMF becoming steeper than Salpeter above $\sigma \gtrsim 200 \text{ km s}^{-1}$. We are also broadly consistent with the IMF variation implied by the non-contracted halo spherical dynamical models of Tortora, Romanowsky & Napolitano (2013). Overall there is a qualitative agreement between different approaches and the still significant systematic differences in the various methods could account for the differences.

5 DISCUSSION

5.1 Previous relations as seen on the Mass Plane

We have shown in Paper XV that the galaxy Fundamental Plane (Dressler et al. 1987; Djorgovski & Davis 1987) can be accurately explained by virial equilibrium combined with a smooth variation of galaxy properties, mainly the total mass-to-light ratio M/L , with velocity dispersion, with galaxies lying on a tight MP $(M_{\text{JAM}}, \sigma_e, R_e^{\text{maj}})$, for a large volume-limited sample of ETGs (Paper I). Once this has been established, the interesting information on galaxy formation is then all contained in the distribution and the physical properties of galaxies on this plane, which we presented in this paper.

We find that on the MP: (i) galaxies sizes are delimited by a lower-boundary, which has a minimum at a characteristic mass $M_{\text{JAM}} \approx 3 \times 10^{10} M_{\odot}$; (ii) A number of key galaxy properties: dynamical M/L , and its population indicators $H\beta$ and colour, as well as the molecular gas fraction, which are mainly related to age, the normalization of the IMF and the prominence of the bulge, all tend to be constant along lines of constant σ_e , on the MP;




Figure 13. IMF versus σ_e correlations. The plots show as blue filled circles with error bars the logarithm of the IMF mismatch parameter $(M/L)_{\text{stars}}/(M/L)_{\text{Salp}}$ versus the effective velocity dispersion σ_e , for the subsets of ATLAS^{3D} galaxies with $H\beta < 2.3 \text{ \AA}$. Green symbols are outliers automatically removed from the fits. Dashed and dotted red lines indicate the 1σ and 2.6σ (99%) observed scatter around the best-fitting relation (black solid line). The top panel includes all galaxies. The middle panel shows galaxies with accurate SBF distances. The bottom panel shows galaxies with $M_{\text{JAM}} < 2 \times 10^{11} M_{\odot}$.

(iii) Another characteristic mass for galaxy properties is the value $M_{\text{JAM}} \approx 2 \times 10^{11} M_{\odot}$ which separates a region dominated by the round or weakly triaxial slow rotators at large masses from one dominated by fast-rotators ETGs, flattened in their outer parts and with embedded exponential disks (Paper XVII), whose characteristics merge smoothly with the ones of spiral galaxies. A transition in the bulge fraction at this galaxy mass appears required in our models for the formation of fast and slow rotators (Khochfar et al. 2011, hereafter Paper VIII).

Although our ATLAS^{3D} sample is limited to a minimum mass $M_{\text{JAM}} \gtrsim 6 \times 10^9 M_{\odot}$, our picture naturally extends to lower masses. As shown in Fig. 9 our trends for fast rotator ETGs continues with the dwarf spheroidal (Sph) sequence at lower masses (see also fig. 7 of Binggeli, Sandage & Tarengi 1984; fig. 38 of Kormendy et al. 2009; fig. 12 of Chen et al. 2010; fig. 4 of Misgeld & Hilker 2011; fig. 20 of Kormendy & Bender 2012), while the spirals sequence continues with a sequence of low-mass late spirals or irregulars (Sc-Irr), as independently noted also by Kormendy & Bender (2012). Interestingly the approximate stellar mass $M_{\star} \approx 2 \times 10^9 M_{\odot}$ where there is a break in the M -size relation of dwarf galaxies and where galaxies with bulge starts to appear (Fig. 9) corresponds to the threshold for quenching of field galaxies recently discovered by Geha et al. (2012). Below that mass only the cluster or group environment can strip spirals of their gas. Perhaps below that mass bulges cannot grow and star formation cannot be quenched by internal processes.

The parallelism between fast rotator ETGs and spiral galaxies in scaling relations was one of the driver for our proposed revision (Paper VII) to the tuning-fork diagram by Hubble (1936), in the spirit of van den Bergh (1976) proposed parallelism between S0s and spirals. In Fig. 14. we summarize our results on morphology, kinematics, population and scaling relations in a single diagram, using the same galaxy symbols as fig. 2 of Paper VII, to provide a link between the two papers. This picture allows us to provide a new perspective and a clean empirical view of a number of classic scaling relation and known trends in galaxy properties.

5.1.1 $L - \sigma$ trends

The classic Faber & Jackson (1976) relation $L \propto \sigma^4$ is well known to be a projection of the FP. We study it here using dynamical mass instead of light. Our study for the first time uses σ_e values integrated within R_e , which properly account for both velocity dispersion and stellar rotation, in combination with accurate dynamical masses which, unlike luminosity, can be properly related to the stellar kinematics. We find that the relation is not well described by an approximately linear trend, but it shows a break which is a reflection of the one in the ZOE (equation 4). The slope of the mean $\sigma - M$ relation (equation 5) changes from $M_{\text{JAM}} \propto \sigma_e^{2.3}$ below $M_{\text{JAM}} \approx 5 \times 10^{10} M_{\odot}$ ($\sigma_e \approx 140 \text{ km s}^{-1}$) to $M_{\text{JAM}} \propto \sigma_e^{4.7}$ at larger masses. Thanks to our use of dynamical masses and σ_e we find for the first time that the bend in the relation is quantitatively consistent with the one we observe in the M -size relation.

The slope we find at the high-mass range is consistent with the original Faber & Jackson (1976) relation, while the low-mass slope in our $\sigma - M$ relation is consistent with claims of a change in the slope of the $\sigma - L$ relation of elliptical galaxies at low luminosities, where the relation was reported to become $L \propto \sigma^2$ (Davies et al. 1983; Held et al. 1992; Matković & Guzmán 2005; de Rijcke et al. 2005; Lauer et al. 2007b; Forbes et al. 2008; Tortora et al. 2009). However previous studies differ from ours because of M/L variations, they ignore stellar rotation, suffer from small number




Figure 14. Schematic summary of the results presented in Section 3 and Section 4. ETGs properties, dynamical M/L (Fig. 1) or its population proxies, $H\beta$ and galaxy colour, as well as the molecular gas fraction (Fig. 3), kinematical concentration (Fig. 6), which traces bulge mass, and IMF mass normalization (Fig. 12), all tend to vary along lines with roughly $R_e \propto M$ (or even better $R_e \propto M^{0.75}$), where σ_e is nearly constant. This sequence of ETGs properties merges smoothly with the one of spiral galaxies, with little overlap between late spirals (Sc-Irr) and ETGs, a significant overlap between early spirals (Sa-Sb) and fast-rotator ETGs with low M/L and no overlaps between spirals and fast-rotators with high M/L . Three characteristic masses are emphasized in this diagram: (i) below $M_* \approx 2 \times 10^9 M_\odot$ there are no regular ETGs and the mass-size lower boundary is increasing; (ii) $M_* \approx 3 \times 10^{10} M_\odot$ is the mass at which ETGs reach their minimum size (or maximum stellar density), before a sudden change in slope $R_e \propto M^{0.75}$ at larger masses (see also fig. 4 in Paper I); (iii) Below $M_* \approx 2 \times 10^{11} M_\odot$ ETGs are dominated by flat fast rotators, showing evidence for disks (Paper XVII), while slow rotators are rare. Above this mass there are no spirals and the population is dominated by quite round or weakly triaxial slow rotators (paper III) with flat (core/deficit) central surface brightness profiles (Paper XXIV). These smooth trends in scaling relations motivated our proposed parallelism between spirals and ETGs. To emphasize this connection we use the same morphology symbols here as in our ‘comb’ diagram in fig. 2 of Paper VII.

statistics and selection biases. This may be the reason why previous studies suggested a value $M_B \approx -20.5$ mag for the break in the $\sigma - L$ relation (see Sec. 3.3.3 of Graham 2013, for a review), which corresponds to the characteristic mass $M \approx 2 \times 10^{11} M_\odot$, using the $(M/L) - \sigma$ relation (Cappellari et al. 2006) and considering a typical galaxy colour $B - I \approx 2.2$ mag. The previously reported mass is clearly inconsistent with our accurate value $M_{\text{JAM},b} \approx 5 \times 10^{10} M_\odot$. Our new result shows that the break in the relation is associated to the characteristic mass $M_{\text{JAM}} \approx 3 \times 10^{10} M_\odot$ of the break in the ZOE and not with the other characteristic mass $M_{\text{JAM}} \approx 2 \times 10^{11} M_\odot$ above which round, slow rotator, cored galaxies dominate.

Observations from the much larger SDSS sample have failed to find evidence for a clear break in the $\sigma - L$ relation (Bernardi et al. 2003; Gallazzi et al. 2006; Hyde & Bernardi 2009a). For this reason the relation is still often assumed to be a power-law. The accuracy and homogeneity of our data, the size of our sample and especially the quantitative consistency between the $\sigma - M$ and $R_e - M$ relation provides an unambiguous confirmation for the break and an accurate determination of the transition mass.

5.1.2 $L - R_e$ trends

Another well known projection of the FP is the Kormendy (1977) relation. When using mass instead of light it becomes clear it represents the analogue of the Faber & Jackson (1976), but this time in the $(M_{\text{JAM}}, R_e^{\text{maj}})$ projection of the MP. Also in this case, when samples are morphologically selected to consist of ellipticals, they tend to populate mostly the region of the diagram near the ZOE, defining a relatively narrow sequence (Graham & Worley 2008; Ko-

rmendy et al. 2009; Chen et al. 2010; Misgeld & Hilker 2011). Although the sequence is useful for a number of studies, it is important to realize that it is not a real sequence in galaxy space on the MP. It is due to the sample selection and it represents essentially one of the contour levels of a continuous trend of galaxy properties, spanning from spiral galaxies, to ETGs, and only terminating on the well defined ZOE (Fig. 9 here and figure 4 in Paper I).

Given that photometry is much easier to obtain than stellar kinematics, a change of slope in the luminosity-size relation has been known for long. It was pointed out by Binggeli, Sandage & Tarengi (1984, their fig. 7) when combining photometry measurements of dwarf spheroidals and ordinary ellipticals: the dwarf spheroidals sequence appears to sharply bend from the ellipticals sequence. Similar differences in the slope of the luminosity-size relation of dwarfs and ellipticals were presented by a number of authors (e.g. Kormendy 1985; Graham & Guzmán 2003; Kormendy et al. 2009; Misgeld & Hilker 2011). The change of slope has been interpreted in different ways. Kormendy (1985) and Kormendy et al. (2009) interpreted dwarf spheroidal as constituting a separate family, of gas-stripped dwarf spirals/irregulars, while Graham & Guzmán (2003) and Graham & Worley (2008) explain the change of slope or curvature in the relation as a natural consequence of the variation of the Sersic (1968) index with luminosity (e.g. Young & Currie 1994; Graham & Guzmán 2003) in a homogeneous class of elliptical galaxies with a range of masses (see Graham 2013 and Kormendy & Bender 2012 for two complementary reviews of this subject).

Our results cannot be compared to theirs in a statistical sense, as galaxies in their diagrams are, by design, not representative of the population in the nearby Universe, and certain classes of ob-

jects (e.g. M32) are over-represented. Our sample is volume-limited and for this reason it gives a statistically representative view of the galaxy population above a certain mass. Still the fact that the sequence of dwarf spheroidals and low-mass spirals/irregulars, lie on the continuation of our trends for fast rotator ETGs and spiral galaxies respectively (Fig. 9), below the $M_* \lesssim 6 \times 10^9 M_\odot$ mass limit of our survey, suggests a continuity between dwarf spheroidals and the low-mass end of our disk-dominated fast-rotator ETGs population, which in turns we showed are closely related to spiral galaxies. For this reason our results reconciles the apparent contrast between the findings of a Sph-E dichotomy (Kormendy 1985; de Rijcke et al. 2005; Janz & Lisker 2008; Kormendy et al. 2009) and the ones of a continuity (Graham & Guzmán 2003; Gavazzi et al. 2005; Ferrarese et al. 2006b; Côté et al. 2006; Janz & Lisker 2009; Boselli et al. 2008; Graham & Worley 2008; Forbes et al. 2011). Our finding in fact agrees with the proposed common origin of dwarf spheroidal and low-mass spiral galaxies and irregulars (Kormendy 1985; Dekel & Silk 1986), but also shows an empirical continuity between dwarf spheroidals and a subset of the ellipticals family. The missing link between Sph and E is constituted by disk-dominated fast rotator ETGs (Emsellem et al. 2007; Cappellari et al. 2007). In fact the continuity we find is not between “true” ellipticals, namely the slow rotator, and dwarf spheroidals, but between “misclassified” ellipticals with disks and S0, namely the fast rotators, and dwarf spheroidals. After this text was written and the parallelism between spiral galaxies and early-type galaxies were presented (Paper VII) to interpret the trends we observed in galaxy scaling relations (Paper I and Cappellari 2011a), an independent confirmation of this picture, including its extension to low mass systems was also provided by Kormendy & Bender (2012).

5.1.3 Population trends with σ or R_e

The characteristic mass $M_{\text{JAM}} \approx 3 \times 10^{10} M_\odot$ is the same transition mass discovered by Kauffmann et al. (2003a) who state that “low-redshift galaxies divide into two distinct families at a stellar mass of $3 \times 10^{10} M_\odot$. Lower-mass galaxies have young stellar populations, low surface mass densities and the low concentrations typical of discs. Their star formation histories are more strongly correlated with surface mass density than with stellar mass.” A similar trend involving colours and also better correlated with surface density (or with the velocity dispersion estimated from the photometry) than with mass, was found to extend to redshift up to $z \approx 3$, with red galaxies being systematically small, and blue galaxies being large at a given mass (Franx et al. 2008). This was recently confirmed, still using photometric data alone, by Bell et al. (2012). The correctness of all these statements can now be easily and accurately verified for the nearby ETGs subset in Fig. 1 and 3.

The novelty of our work, with respect to all previous studies, is that we have unprecedentedly accurate and unbiased dynamical masses, and stellar velocity dispersions σ_e integrated within a large aperture ($1R_e$), instead of σ values inferred from photometry. This allow us to conclusively state that neither dynamical mass nor stellar surface density are actually the best descriptor of galaxy properties, the main trend being along the σ_e direction (which includes rotation in the case of disk galaxies). Our clean ATLAS^{3D} result was already presented in Cappellari (2011a) and subsequently confirmed with SDSS data and using virial mass estimates by Wake, van Dokkum & Franx (2012). The trend we observed for the ETGs can be extended to spiral galaxies, which fill the region of larger sizes above the ETGs in the $(M_{\text{JAM}}, R_e^{\text{maj}})$ projection, smoothly overlapping with the ETGs for the largest spiral bulge fractions (fig-

ure 4 of Paper I and Fig. 9). It has been known for long that in spirals luminosity-weighted ages are lower, star formation is larger and colours bluer on average than the ETGs, essentially by definition (e.g. Hubble 1936; van den Bergh 1976).

Similar trends between age and size (or surface brightness), with older objects being smaller at given age, were found to persist in ETGs. van der Wel et al. (2009a) state that “at a given stellar velocity dispersion, SDSS data show that there is no relation between size and age”. The same age-size trend was pointed out by Shankar & Bernardi (2009), and in different terms by Graves, Faber & Schiavon (2009) who state that “no stellar population property shows any dependence on R_e at fixed σ , suggesting that σ and not dynamical mass is the better predictor of past SFH”. These findings are another way of saying that age variations must follow lines of constant σ on the MP as we find here for M/L , $H\beta$, colour and molecular gas fraction, and confirm using age for our sample in McDermid et al. (in preparation), in agreement with Gallazzi et al. (2006, their figure 15). The age-size trend was also confirmed in different samples of nearby galaxies by Valentinuzzi et al. (2010) and Napolitano, Romanowsky & Tortora (2010), and a similar trend, with star forming galaxies being larger than passive ones was found in place from redshift $z \sim 2.5$ (Williams et al. 2010; Wuyts et al. 2011; Newman et al. 2012). The only contrasting view is the one by Trujillo, Ferreras & de La Rosa (2011), who find a lack of age-size trend both at low and high-redshift.

The same characteristics mass $M_* \approx 3 \times 10^{10} M_\odot$ of Kauffmann et al. (2003a), which constitute the location of the break in our ZOE, was found by Hyde & Bernardi (2009a) in the mass-size relation of 5×10^4 SDSS galaxies. Their trend is significant but quite subtle. The reason for the curvature they find becomes clear from what we find: the average radius of the objects on the MP at constant mass, bends upwards at low masses, due to the cusp in the ZOE, with the strength of the effect being dependent on the specific criterion adopted to select ETGs.

5.2 Implications for galaxy formation

Galaxy formation is the superposition of a number of complex events that happen in parallel. Here we sketch a tentative picture of some of the phenomena that can play a role in explaining what we see. We refer the reader to Bois et al. (2011, hereafter Paper VI), Paper VIII and Naab et al. (in preparation) for a more in-depth discussion.

The smoothness and regularity of the trends we observe and the fact that they extend to spirals, appears to indicate a close connection between the formation of the two classes of objects (Paper VIII). The same similarity between the fast rotator ETGs and spirals in terms of their morphology and degree of rotation, lead us to propose a revision (Paper VII) of the classic morphological classification (Hubble 1936) to emphasize the parallelism between the fast rotators and spirals, in the same way that van den Bergh (1976) proposed it for S0s and spirals. Only the most massive slow rotators appear to form an empirically separated class. The *kinematic* morphology-density relation (Paper VII), which applies to our kinematic classes the relation discovered by Dressler (1980), suggest that most spirals are being transformed into fast rotators due to environmental effects (Khochfar & Ostriker 2008), with a mechanism that is sufficiently ‘gentle’ to preserve the near axisymmetry of the disk (Paper II).

One process which is often mentioned in the context of cluster galaxy populations is stripping by the interstellar medium (Spitzer & Baade 1951; Abadi, Moore & Bower 1999), which may impact

the global morphology of the galaxy by removing a significant fraction (or most) of its gas content but mostly preserving the stellar disk component. This may explain some of the most flattened galaxies in the fast rotator class (e.g., NGC4762), as emphasised in Paper VII. This would work by removing some of the mass of the galaxy keeping a relatively constant effective size (of the stellar component), and may contribute to the scatter in the Mass-size plane for fast rotator as well as to their overlap with the more gas-rich spiral galaxies.

As to explain the intermediate to high mass end of the fast rotator class, we would need a process which is able to increase the bulge size, while at the same time removing the gas or shutting off star formation (Paper VIII). The empirical signatures of this phenomenon are visible in our data as an ‘apparent’ decrease of the galaxy size, and increase of σ_e , which is actually due to the bulge concentrating more light at smaller radii, accompanied by an increase in M/L , which is tracing a decrease of star formation or an age increase. The process appears to generally preserve the intrinsic flatness of the stellar disks at large radii (top panel of Fig. 7). A similar scenario was recently proposed by Bell et al. (2012) to interpret the relationship between rest-frame optical colour, stellar mass, star formation activity, and galaxy structure from $z \approx 2$ to the present day.

Intense gas-rich accretion events, mostly via cold streams (Kereš et al. 2005; Dekel et al. 2009), or major gas rich mergers (Paper VIII), will increase both the mass and σ_e . During the accretion the gas may sink toward the centre (Mihos & Hernquist 1994) until it becomes self-gravitating and starts forming stars. It is during this phase of rapid gas accretion that the $(M/L) - \sigma$ (Cappellari et al. 2006; van der Marel & van Dokkum 2007) and IMF – σ (Section 4.3) relations, the tilt of the FP (Dressler et al. 1987; Faber et al. 1987; Djorgovski & Davis 1987), the $Mgb - \sigma$ (Burstein et al. 1988; Bender, Burstein & Faber 1993) or the $Mgb - V_{esc}$ relation (Davies, Sadler & Peletier 1993; Scott et al. 2009, Paper XXI), will be imprinted in the ETGs population (Robertson et al. 2006; Hopkins et al. 2009b) and then mostly preserved in the following evolution.

The early progenitors of today’s fast rotator ETGs would be high-redshift spirals, which are different from local ones. They are characterized by giant gas clumps (Elmegreen et al. 2007; Genzel et al. 2011) have high gas fractions (Tacconi et al. 2010; Daddi et al. 2010), possess large velocity dispersion and are dominated by turbulent motions (Förster Schreiber et al. 2006, 2009; Genzel et al. 2006, 2008; Law et al. 2012). In that situation bulges may form naturally as the clumps collide and sink to the centre (Bournaud, Elmegreen & Elmegreen 2007; Dekel, Sari & Ceverino 2009), unless they are efficiently destroyed by stellar feedback (Genel et al. 2012; Hopkins et al. 2012). Secular effects (Kormendy & Kennicutt 2004) will also contribute to the bulge growth and σ_e increase, while keeping the mass unchanged, as will contribute the destabilizing effect of minor mergers.

During the bulge growth some process must be able to turn off star formation, without destroying the fast rotating disks that still dominate the local ETGs population (Paper II; Paper III) and that dominates the ETGs population already from $z \sim 2$ (van der Wel et al. 2011; see also van Dokkum 2011). The reduced efficiency of star formation by morphological quenching (Martig et al. 2009; Ceverino, Dekel & Bournaud 2010) may be one of the processes explaining why on average bigger bulges correspond to older ages and larger M/L . However we showed in Fig. 3 that the cold gas fraction is decreasing with the bulge fraction, which shows that less fuel is available for galaxies with more massive bulges. Bulge/spheroid growth seems also associated with AGN feedback, which would




Figure 15. Evolution scenario for ETGs. The symbols are the same as in Fig. 1, while the large arrows indicate the proposed interpretation of the observed distribution as due to a combination of two processes: (a) in-situ star formation: bulge or spheroid growth, which seems associated to the quenching of star formation, which moves galaxies to the right of towards the bottom, due to the increased concentration (decreasing R_e and increasing σ_e), and (b) external accretion: dry mostly minor merging, increasing R_e by moving galaxies along lines of roughly constant σ_e (or steeper), while leaving the population unchanged. A schematic illustration of these two processes is shown in Fig. 16.

provide another bulge-related quenching mechanism (Silk & Rees 1998; Granato et al. 2004; Bower et al. 2006; Hopkins et al. 2006; Croton et al. 2006). During the sequence of bulge/spheroid-growth followed by quenching, the original gas rich spiral will move from the left of the observed (M_{JAM}, R_e^{maj}) plane towards the right, or possibly the bottom right due to the increased concentration, while intersecting the constant σ lines (blue arrow in Fig. 15). At the end of the evens the galaxy will be a fast rotator ETG, generally more massive and with a bigger bulge (smaller R_e , larger σ_e and concentration) than the precursor clumpy spiral.

At any stage during the bulge-growth phase the galaxy may accrete purely stellar satellites (Khochfar & Silk 2006). In the case of these dry mergers the situation is quite different and one can predict the final configuration using energy conservation (Hernquist, Spiegel & Heyl 1993; Cole et al. 2000; Boylan-Kolchin, Ma & Quataert 2006; Ciotti, Lanzoni & Volonteri 2007). The predictions show that sizes will increase as the mass grows. For major mergers (equal mass) and typical orbital configurations, one can show that the mass and radius double, while σ remains constant in agreement with simulations (Nipoti et al. 2009; see Hilz et al. 2012 and Hilz, Naab & Ostriker 2013 for more accurate numerical simulations and detailed physical explanations of this process). While in the limit in which the same mass doubling happens via small satellites, as mostly expected from the shape of the Schechter (1976) mass function, the radius will increase by a factor of four and the dispersion will be twice smaller (Naab, Johansson & Ostriker 2009; Bezanon et al. 2009; Hopkins et al. 2009a; Hilz et al. 2012). As a result the galaxy will move along lines that are parallel to the constant σ




Figure 16. Schematic representation of the two main processes responsible for the formation of the observed distribution of galaxies on the MP. (a) In-situ star formation: bulge growth via cold accretion, secular evolution, or minor gas-rich mergers, followed by quenching by AGN or other mechanisms, leaving the galaxy more massive, more compact, and consequently with a larger σ_e , and gas poor (blue arrow in Fig. 15); (b) external accretion: major or minor dry mergers, increasing galaxy mass and sizes at nearly constant σ_e , or with a possible decrease, leaving the population mostly unchanged (red arrow in Fig. 15). (taken from Cappellari 2011b).

lines, or steeper. During these dry mergers, given that there is little gas involved, the stellar population, colour and M/L will remain unchanged (red arrow in Fig. 15).

When the galaxy grows more massive than the characteristic mass $M_* \gtrsim 2 \times 10^{11} M_\odot$, a transition seems to happen in their formation, possibly associated to the shock heating of the infalling gas, due to the halo mass (Dekel & Birnboim 2006). Above this mass galaxies are in fact embedded in massive and hot X-ray halos (Kormendy et al. 2009, Paper XIX) which prevents any further cold gas accretion onto the galaxy.

The gas-rich mergers/accretion scenario is generally consistent with the observed correlation between supermassive black holes (BH) and galaxy velocity dispersion or bulge mass (Di Matteo, Springel & Hernquist 2005). It is generally believed that the correlations indicate a joint evolution of galaxies and BHs, with BH growth happening at the same time as the bulge growth, and providing a self-regulation via feedback (Silk & Rees 1998; but see Peng 2007 and Jahnke & Macciò 2011 for a non-causal origin of the correlations). We demonstrate that indeed the σ variation directly traces the bulge growth (Kormendy & Gebhardt 2001). In fact σ traces the central galaxy concentration and the bulge size as estimated on optical morphology, while the outer galaxy disks remain flat. This implies that, if the BH indeed accretes from the same gas that grows the bulge, BH mass should correlate better with σ than with total mass as observed (Gebhardt et al. 2000; Ferrarese & Merritt 2000).

However, when ETGs experience dry mergers, their BH grows in proportion to the mass, but galaxies move along lines of nearly constant σ . For this reason, an expectation from this picture is that BHs at the high mass end should start to appear too massive with respect to the predictions of the $M_{\text{BH}} - \sigma$ relation (see also Nipoti, Londrillo & Ciotti 2003; Boylan-Kolchin, Ma & Quataert 2006; Ciotti 2009). The high-mass BH end is still not sufficiently populated to reliably test this prediction, but indirect evidence seems to support this possibility (Lauer et al. 2007b). Early direct evidences come from the recent detection of two giant black holes at the centre of two bright cluster galaxies, which are clearly more massive than the $M_{\text{BH}} - \sigma$ prediction (McConnell et al. 2011) or the related

observation that non-core ETGs follow a more shallow $M_{\text{BH}} - \sigma$ relation than core ETGs (Graham 2012).

In summary we propose that the distribution of galaxy properties on the MP, where M/L and age follows lines of constant σ on the MP, could be explained by the combination of two processes, which can happen multiple times during the evolution of a single galaxy: (i) They accrete gas, which grows the bulge and BH, shrinks their R_e , and increases σ_e and concentration, while some process which seems associated with the bulge growth (e.g. AGN feedback), quenches star formation; (ii) They experience mostly minor dry mergers that move them along lines of roughly constant σ (or steeper).

An open question in the scenario in which ETGs evolve relatively quietly from spirals comes from the comparison of our findings with the empirical scaling relations one observes at larger redshift. In fact at $z \gtrsim 1.5$ galaxies are found to be smaller than local ones with the same mass (Daddi et al. 2005; Trujillo et al. 2006, 2007; van Dokkum et al. 2008; Cimatti et al. 2008). They populate the region below the ZOE of local galaxies, although their σ tend to be consistent with our local observations (Cappellari 2013). This may indicate that the compact high-redshift ETGs follow a different and more violent evolutionary path than the more quiet majority of local ETGs, as suggested by other high-redshift observations (Barro et al. 2013). A way to reconcile this very mild (or lack of) σ evolution is by assuming that the compact primordial ETGs grow mostly by accretion of small satellites in their outer halos, while preserving the central structure (Naab, Johansson & Ostriker 2009; Hopkins et al. 2010; Oser et al. 2010, 2012). This seems consistent with the shape of the photometric profiles of the early ETGs (Hopkins et al. 2009a; Bezanson et al. 2009; van Dokkum et al. 2010; Hilz, Naab & Ostriker 2013). A caveat is that significant biases may still exist in the high-redshift photometry (Mancini et al. 2010), considering that systematic differences of up to a factor of two exists even on well observed ETGs in the nearby Universe (Kormendy et al. 2009, Chen et al. 2010, Paper I). Moreover comparisons of photometric profiles tend to be made against bona fide ellipticals, while the remnant of the high-redshift ETGs are likely disk-like fast rotators ETG, which have systematically different profiles. Finally the comparisons should ideally be done in mass density, instead of surface density, but kinematic information is available for only a handful of galaxies. This implies that there is perhaps still some room for the compact high- z ETGs to become more consistent with local ones, than currently assumed.

6 SUMMARY

In the companion Paper XV we describe in detail the axisymmetric dynamical models for all the ATLAS^{3D} ETGs that were introduced in Cappellari et al. (2012). We found that galaxies lie on a thin Mass Plane (MP) in the three-dimensional parameter space defined by $(M_{\text{JAM}}, \sigma_e, R_e^{\text{maj}})$.

Here we studied the inclined projection of the MP and find that: (i) the location galaxies define a clear zone of exclusion (ZOE), roughly described by two power-laws, joined by a break at a first characteristic mass $M_{\text{JAM}} \equiv L \times (M/L)_e \approx 3 \times 10^{10} M_\odot$, which corresponds to the regime where a number of global galaxy properties were reported to change; (ii) we find a second characteristic mass $M_{\text{JAM}} \approx 2 \times 10^{11} M_\odot$, which separates a region with only nearly round slow rotator ETGs at large masses from one dominated by fast rotator ETGs with disks and spiral galaxies at lower masses; (iii) The distribution of $(M/L)_e$, as well as population indicators of

$(M/L)_{\text{pop}}$ like $H\beta$ and colour, and the molecular gas fraction, all tend to be constant along lines of constant σ_e on the $(M_{\text{JAM}}, \sigma_e)$ plane, or correspondingly $R_e^{\text{maj}} \propto M_{\text{JAM}}$ (or even better parallel to the ZOE $R_e^{\text{maj}} \propto M_{\text{JAM}}^{0.75}$) on the $(M_{\text{JAM}}, R_e^{\text{maj}})$ plane; (iv) properties of ETGs merge smoothly and form parallel sequences to those of spiral galaxies; (v) below our survey mass limit, the distribution of fast rotators ETGs continues with that of dwarf spheroidal (Sph), while the sequence of late spirals continues with the dwarf irregulars (Im).

We show that, at fixed mass, σ_e in ETGs traces the galaxy bulge fraction, which appears the main driver for the changes in galaxy properties that we observe. These findings explains why σ_e (not L , M_* , R_e or Σ_e) has often been the most successful single descriptor of galaxy properties. We discuss a number of previously found galaxy correlation and we find that they are all consistent with what we find. In fact most observed relations turn out to be just special projections of the cleaner and more general view provided by the MP.

The stellar initial mass function was previously shown to vary systematically with $(M/L)_{\text{stars}}$ (Cappellari et al. 2012) and as expected it also follows the other population indicators. It also tend to vary along lines of nearly constant σ_e on the MP. This is a necessary expectation if one wants to preserve the tightness of the Fundamental Plane or the $(M/L) - \sigma_e$ relation. The trend of IMF with σ appears to account for about half of the total trend in the $(M/L) - \sigma_e$ relation (see Paper XV), the remaining one being due to stellar population variations.

The distribution of galaxy properties on the MP can be qualitatively interpreted as due to a combination of two main processes: (i) bulge growth, changing the galaxy population and decreasing R_e , while increasing σ_e , and (ii) dry merging, increasing R_e by moving galaxies along lines of approximately constant σ_e , while leaving the population unchanged. It is unclear where the reported dense ETGs fit in this picture. They may simply follow a different route, which forms the most massive galaxies and is not affecting the fast rotator population, which dominates ETGs in the nearby Universe. Or there may be some remaining biases in the photometric observations, producing an underestimation of the radius. Stellar kinematics of high-redshift ETGs is critically needed to address this question.

ACKNOWLEDGEMENTS

MC acknowledges support from a Royal Society University Research Fellowship. This work was supported by the rolling grants ‘Astrophysics at Oxford’ PP/E001114/1 and ST/H002456/1 and visitors grants PPA/V/S/2002/00553, PP/E001564/1 and ST/H504862/1 from the UK Research Councils. RLD acknowledges travel and computer grants from Christ Church, Oxford and support from the Royal Society in the form of a Wolfson Merit Award 502011.K502/jd. RLD also acknowledges the support of the ESO Visitor Programme which funded a 3 month stay in 2010. SK acknowledges support from the Royal Society Joint Projects Grant JP0869822. RMcD is supported by the Gemini Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., on behalf of the international Gemini partnership of Argentina, Australia, Brazil, Canada, Chile, the United Kingdom, and the United States of America. TN and MBois acknowledge support from the DFG Cluster of Excellence ‘Origin and Structure of the Universe’. MS acknowledges support from a STFC Advanced Fellowship ST/F009186/1. PS is a NWO/Veni fellow. (TAD) The

research leading to these results has received funding from the European Community’s Seventh Framework Programme (FP7/2007-2013/) under grant agreement No 229517. MBois has received, during this research, funding from the European Research Council under the Advanced Grant Program Num 267399-Momentum. The authors acknowledge financial support from ESO.

REFERENCES

- Abadi M. G., Moore B., Bower R. G., 1999, MNRAS, 308, 947
 Aihara H. et al., 2011, ApJS, 193, 29
 Auger M. W., Treu T., Bolton A. S., Gavazzi R., Koopmans L. V. E., Marshall P. J., Bundy K., Moustakas L. A., 2009, ApJ, 705, 1099
 Auger M. W., Treu T., Bolton A. S., Gavazzi R., Koopmans L. V. E., Marshall P. J., Moustakas L. A., Burles S., 2010a, ApJ, 724, 511
 Auger M. W., Treu T., Gavazzi R., Bolton A. S., Koopmans L. V. E., Marshall P. J., 2010b, ApJ, 721, L163
 Bacon R. et al., 2001, MNRAS, 326, 23
 Baldry I. K., Glazebrook K., Brinkmann J., Ivezić Ž., Lupton R. H., Nichol R. C., Szalay A. S., 2004, ApJ, 600, 681
 Barnabè M., Czoske O., Koopmans L. V. E., Treu T., Bolton A. S., 2011, MNRAS, 415, 2215
 Barnabè M. et al., 2012, MNRAS, 423, 1073
 Barro G. et al., 2013, ApJ, 765, 104
 Bastian N., Covey K. R., Meyer M. R., 2010, ARA&A, 48, 339
 Bell E. F., de Jong R. S., 2001, ApJ, 550, 212
 Bell E. F. et al., 2012, ApJ, 753, 167
 Bender R., Burstein D., Faber S. M., 1992, ApJ, 399, 462
 Bender R., Burstein D., Faber S. M., 1993, ApJ, 411, 153
 Bernardi M., Roche N., Shankar F., Sheth R. K., 2011, MNRAS, 412, L6
 Bernardi M. et al., 2003, AJ, 125, 1866
 Bershady M. A., Martinsson T. P. K., Verheijen M. A. W., Westfall K. B., Andersen D. R., Swaters R. A., 2011, ApJ, 739, L47
 Bershady M. A., Verheijen M. A. W., Swaters R. A., Andersen D. R., Westfall K. B., Martinsson T., 2010, ApJ, 716, 198
 Bezanson R., van Dokkum P. G., Tal T., Marchesini D., Kriek M., Franx M., Coppi P., 2009, ApJ, 697, 1290
 Binggeli B., Sandage A., Tarenghi M., 1984, AJ, 89, 64
 Binney J., Tremaine S., 2008, Galactic Dynamics: Second Edition. Princeton University Press
 Blanton M. R. et al., 2003, ApJ, 592, 819
 Bois M. et al., 2011, MNRAS, 416, 1654 (Paper VI)
 Bolton A. S., Burles S., Koopmans L. V. E., Treu T., Gavazzi R., Moustakas L. A., Wayth R., Schlegel D. J., 2008a, ApJ, 682, 964
 Bolton A. S., Burles S., Koopmans L. V. E., Treu T., Moustakas L. A., 2006, ApJ, 638, 703
 Bolton A. S., Treu T., Koopmans L. V. E., Gavazzi R., Moustakas L. A., Burles S., Schlegel D. J., Wayth R., 2008b, ApJ, 684, 248
 Boselli A., Boissier S., Cortese L., Gavazzi G., 2008, ApJ, 674, 742
 Bournaud F., Elmegreen B. G., Elmegreen D. M., 2007, ApJ, 670, 237
 Bower R. G., Benson A. J., Malbon R., Helly J. C., Frenk C. S., Baugh C. M., Cole S., Lacey C. G., 2006, MNRAS, 370, 645
 Boylan-Kolchin M., Ma C.-P., Quataert E., 2006, MNRAS, 369, 1081
 Brewer B. J. et al., 2012, MNRAS, 422, 3574
 Burstein D., Bender R., Faber S., Nolthenius R., 1997, AJ, 114, 1365
 Burstein D., Davies R. L., Dressler A., Faber S. M., Lynden-Bell D., 1988, in Astrophysics and Space Science Library, Vol. 141, Towards Understanding Galaxies at Large Redshift, Renzini R. G. K. . A., ed., pp. 17–21
 Caon N., Capaccioli M., D’Onofrio M., 1993, MNRAS, 265, 1013
 Cappellari M., 2002, MNRAS, 333, 400
 Cappellari M., 2008, MNRAS, 390, 71
 Cappellari M., 2011a, in Paper presented at the conference on Galaxy Formation held 18–22 July, 2011 at Durham University, Durham, UK. Online at <http://astro.dur.ac.uk/Gal2011/talks.php>
 Cappellari M., 2011b, Nature, 480, 187

- Cappellari M., 2013, in IAU Symposium, Vol. 295, The Intriguing Life of Massive Galaxies, Thomas D., Pasquali A., Ferreras I., eds., p. (arXiv:1210.7742)
- Cappellari M. et al., 2006, MNRAS, 366, 1126
- Cappellari M., Emsellem E., 2004, PASP, 116, 138
- Cappellari M. et al., 2007, MNRAS, 379, 418
- Cappellari M. et al., 2011a, MNRAS, 413, 813 (Paper I)
- Cappellari M. et al., 2011b, MNRAS, 416, 1680 (Paper VII)
- Cappellari M. et al., 2012, Nature, 484, 485
- Cappellari M. et al., 2013, MNRAS in press (arXiv:1208.3522), (Paper XV)
- Carlberg R. G., 1986, ApJ, 310, 593
- Cenarro A. J., Gorgas J., Vazdekis A., Cardiel N., Peletier R. F., 2003, MNRAS, 339, L12
- Ceverino D., Dekel A., Bournaud F., 2010, MNRAS, 404, 2151
- Chabrier G., 2003, PASP, 115, 763
- Chen C., Côté P., West A. A., Peng E. W., Ferrarese L., 2010, ApJS, 191, 1
- Chiosi C., Bressan A., Portinari L., Tantalo R., 1998, A&A, 339, 355
- Cimatti A. et al., 2008, A&A, 482, 21
- Ciotti L., 1991, A&A, 249, 99
- Ciotti L., 2009, Nuovo Cimento Rivista Serie, 32, 1
- Ciotti L., Lanzoni B., Volonteri M., 2007, ApJ, 658, 65
- Cleveland W., 1979, Journal of the American statistical association, 829
- Cleveland W., Devlin S., 1988, Journal of the American Statistical Association, 596
- Cole S., Lacey C. G., Baugh C. M., Frenk C. S., 2000, MNRAS, 319, 168
- Conroy C., Gunn J. E., White M., 2009, ApJ, 699, 486
- Conroy C., van Dokkum P., 2012a, ApJ, 747, 69
- Conroy C., van Dokkum P. G., 2012b, ApJ, 760, 71
- Côté P. et al., 2006, ApJS, 165, 57
- Croton D. J. et al., 2006, MNRAS, 365, 11
- Daddi E. et al., 2010, ApJ, 713, 686
- Daddi E. et al., 2005, ApJ, 626, 680
- Davé R., 2008, MNRAS, 385, 147
- Davies R. L., Efstathiou G., Fall S. M., Illingworth G., Schechter P. L., 1983, ApJ, 266, 41
- Davies R. L., Sadler E. M., Peletier R. F., 1993, MNRAS, 262, 650
- de Rijcke S., Michielsen D., Dejonghe H., Zeilinger W. W., Hau G. K. T., 2005, A&A, 438, 491
- de Vaucouleurs G., 1959, Handbuch der Physik, 53, 311
- de Zeeuw P. T. et al., 2002, MNRAS, 329, 513
- Deason A. J., Belokurov V., Evans N. W., McCarthy I. G., 2012, ApJ, 748, 2
- Dekel A., Birnboim Y., 2006, MNRAS, 368, 2
- Dekel A. et al., 2009, Nature, 457, 451
- Dekel A., Sari R., Ceverino D., 2009, ApJ, 703, 785
- Dekel A., Silk J., 1986, ApJ, 303, 39
- Di Matteo T., Springel V., Hernquist L., 2005, Nature, 433, 604
- Djorgovski S., Davis M., 1987, ApJ, 313, 59
- D'Onofrio M. et al., 2008, ApJ, 685, 875
- Dressler A., 1980, ApJ, 236, 351
- Dressler A., Lynden-Bell D., Burstein D., Davies R. L., Faber S. M., Terlevich R., Wegner G., 1987, ApJ, 313, 42
- Driver S. P. et al., 2011, MNRAS, 413, 971
- Dutton A. A. et al., 2011a, MNRAS, 417, 1621
- Dutton A. A. et al., 2011b, MNRAS, 416, 322
- Dutton A. A., Maccio' A. V., Mendel J. T., Simard L., 2012, MNRAS submitted (arXiv:1204.2825)
- Dutton A. A., Mendel J. T., Simard L., 2012, MNRAS, 422, L33
- Dutton A. A. et al., 2013, MNRAS, 428, 3183
- Elmegreen D. M., Elmegreen B. G., Ravindranath S., Coe D. A., 2007, ApJ, 658, 763
- Emsellem E. et al., 2007, MNRAS, 379, 401
- Emsellem E. et al., 2004, MNRAS, 352, 721
- Emsellem E., Monnet G., Bacon R., 1994, A&A, 285, 723
- Faber S. M. et al., 1997, AJ, 114, 1771
- Faber S. M., Dressler A., Davies R. L., Burstein D., Lynden-Bell D., 1987, in Nearly Normal Galaxies. From the Planck Time to the Present, Faber S. M., ed., pp. 175–183
- Faber S. M., Jackson R. E., 1976, ApJ, 204, 668
- Falcón-Barroso J., Sánchez-Blázquez P., Vazdekis A., Ricciardelli E., Cardiel N., Cenarro A. J., Gorgas J., Peletier R. F., 2011, A&A, 532, A95
- Ferrarese L., 2012, in Paper presented at the conference on Science from the Next Generation Imaging and Spectroscopic Surveys held 15-18 October, 2012 at ESO Garching, Germany. Online at <http://www.eso.org/sci/meetings/2012/surveys2012/program.html>
- Ferrarese L. et al., 2012, ApJS, 200, 4
- Ferrarese L. et al., 2006a, ApJ, 644, L21
- Ferrarese L. et al., 2006b, ApJS, 164, 334
- Ferrarese L., Merritt D., 2000, ApJ, 539, L9
- Ferreras I., La Barbera F., de la Rosa I. G., Vazdekis A., de Carvalho R. R., Falcón-Barroso J., Ricciardelli E., 2013, MNRAS, 429, L15
- Ferreras I., Saha P., Burles S., 2008, MNRAS, 383, 857
- Ferreras I., Saha P., Leier D., Courbin F., Falco E. E., 2010, MNRAS, 409, L30
- Forbes D. A., Lasky P., Graham A. W., Spitler L., 2008, MNRAS, 389, 1924
- Forbes D. A., Spitler L. R., Graham A. W., Foster C., Hau G. K. T., Benson A., 2011, MNRAS, 413, 2665
- Förster Schreiber N. M. et al., 2009, ApJ, 706, 1364
- Förster Schreiber N. M. et al., 2006, ApJ, 645, 1062
- Franx M., van Dokkum P. G., Schreiber N. M. F., Wuyts S., Labbé I., Toft S., 2008, ApJ, 688, 770
- Freeman K. C., 1970, ApJ, 160, 811
- Gallazzi A., Bell E. F., 2009, ApJS, 185, 253
- Gallazzi A., Charlot S., Brinchmann J., White S. D. M., 2006, MNRAS, 370, 1106
- Gargiulo A. et al., 2009, MNRAS, 397, 75
- Gavazzi G., Donati A., Cucciati O., Sabatini S., Boselli A., Davies J., Zibetti S., 2005, A&A, 430, 411
- Gavazzi R., Treu T., Rhodes J. D., Koopmans L. V. E., Bolton A. S., Burles S., Massey R. J., Moustakas L. A., 2007, ApJ, 667, 176
- Gebhardt K. et al., 2000, ApJ, 539, L13
- Geha M., Blanton M. R., Yan R., Tinker J. L., 2012, ApJ, 757, 85
- Gelman A., Carlin J., Stern H., Rubin D., 2004, Bayesian data analysis. CRC press
- Genel S. et al., 2012, ApJ, 745, 11
- Genzel R. et al., 2008, ApJ, 687, 59
- Genzel R. et al., 2011, ApJ, 733, 101
- Genzel R. et al., 2006, Nature, 442, 786
- Gerhard O., Kronawitter A., Saglia R. P., Bender R., 2001, AJ, 121, 1936
- Gerhard O. E., Binney J. J., 1996, MNRAS, 279, 993
- Gnedin O. Y., Ceverino D., Gnedin N. Y., Klypin A. A., Kravtsov A. V., Levine R., Nagai D., Yepes G., 2011, ApJ submitted (arXiv:1108.5736)
- Gnedin O. Y., Kravtsov A. V., Klypin A. A., Nagai D., 2004, ApJ, 616, 16
- Governato F. et al., 2010, Nature, 463, 203
- Graham A. W., 2012, ApJ, 746, 113
- Graham A. W., 2013, in Planets, stars and stellar systems, Oswald T. D., Keel W. C., eds., Vol. 6, (Berlin: Springer), pp. 91–139 (arXiv:1108.0997)
- Graham A. W., Guzmán R., 2003, AJ, 125, 2936
- Graham A. W., Worley C. C., 2008, MNRAS, 388, 1708
- Granato G. L., De Zotti G., Silva L., Bressan A., Danese L., 2004, ApJ, 600, 580
- Graves G. J., Faber S. M., 2010, ApJ, 717, 803
- Graves G. J., Faber S. M., Schiavon R. P., 2009, ApJ, 698, 1590
- Grillo C., Gobat R., 2010, MNRAS, 402, L67
- Grillo C., Gobat R., Lombardi M., Rosati P., 2009, A&A, 501, 461
- Gunawardhana M. L. P. et al., 2011, MNRAS, 415, 1647
- Heavens A., Panter B., Jimenez R., Dunlop J., 2004, Nature, 428, 625
- Held E. V., de Zeeuw T., Mould J., Picard A., 1992, AJ, 103, 851
- Hernquist L., 1990, ApJ, 356, 359
- Hernquist L., Spergel D. N., Heyl J. S., 1993, ApJ, 416, 415
- Hilz M., Naab T., Ostriker J. P., 2013, MNRAS, 429, 2924

- Hilz M., Naab T., Ostriker J. P., Thomas J., Burkert A., Jesseit R., 2012, *MNRAS*, 425, 3119
- Hopkins P. F., Bundy K., Hernquist L., Wuyts S., Cox T. J., 2010, *MNRAS*, 401, 1099
- Hopkins P. F., Bundy K., Murray N., Quataert E., Lauer T. R., Ma C.-P., 2009a, *MNRAS*, 398, 898
- Hopkins P. F., Hernquist L., Cox T. J., Di Matteo T., Robertson B., Springel V., 2006, *ApJS*, 163, 1
- Hopkins P. F., Hernquist L., Cox T. J., Keres D., Wuyts S., 2009b, *ApJ*, 691, 1424
- Hopkins P. F., Kereš D., Murray N., Quataert E., Hernquist L., 2012, *MNRAS*, 427, 968
- Hoversten E. A., Glazebrook K., 2008, *ApJ*, 675, 163
- Hubble E. P., 1936, *Realm of the Nebulae*. Yale Univ. Press, New Haven
- Huchra J., Gorenstein M., Kent S., Shapiro I., Smith G., Horine E., Perley R., 1985, *AJ*, 90, 691
- Huchra J. P. et al., 2012, *ApJS*, 199, 26
- Hyde J. B., Bernardi M., 2009a, *MNRAS*, 394, 1978
- Hyde J. B., Bernardi M., 2009b, *MNRAS*, 396, 1171
- Jaffe W., 1983, *MNRAS*, 202, 995
- Jahnke K., Macciò A. V., 2011, *ApJ*, 734, 92
- Janz J., Lisker T., 2008, *ApJ*, 689, L25
- Janz J., Lisker T., 2009, *ApJ*, 696, L102
- Jiang F., van Dokkum P., Bezanson R., Franx M., 2012, *ApJ*, 749, L10
- Kassin S. A., de Jong R. S., Weiner B. J., 2006, *ApJ*, 643, 804
- Kauffmann G. et al., 2003a, *MNRAS*, 341, 33
- Kauffmann G. et al., 2003b, *MNRAS*, 341, 54
- Kelly B. C., 2007, *ApJ*, 665, 1489
- Kelson D. D., van Dokkum P. G., Franx M., Illingworth G. D., Fabricant D., 1997, *ApJ*, 478, L13
- Kennicutt, Jr. R. C., 1998, in *Astronomical Society of the Pacific Conference Series*, Vol. 142, *The Stellar Initial Mass Function* (38th Herstmonceux Conference), Howell G. G., D., ed., p. 1
- Kereš D., Katz N., Weinberg D. H., Davé R., 2005, *MNRAS*, 363, 2
- Khochfar S. et al., 2011, *MNRAS*, 417, 845 (Paper VIII)
- Khochfar S., Ostriker J. P., 2008, *ApJ*, 680, 54
- Khochfar S., Silk J., 2006, *MNRAS*, 370, 902
- Kirby E. M., Jerjen H., Ryder S. D., Driver S. P., 2008, *AJ*, 136, 1866
- Koopmans L. V. E., Treu T., Bolton A. S., Burles S., Moustakas L. A., 2006, *ApJ*, 649, 599
- Kormendy J., 1977, *ApJ*, 218, 333
- Kormendy J., 1985, *ApJ*, 295, 73
- Kormendy J., Bender R., 2012, *ApJS*, 198, 2
- Kormendy J., Fisher D. B., Cornell M. E., Bender R., 2009, *ApJS*, 182, 216
- Kormendy J., Gebhardt K., 2001, in *American Institute of Physics Conference Series*, Vol. 586, *20th Texas Symposium on relativistic astrophysics*, Wheeler J. C., Martel H., eds., pp. 363–+
- Kormendy J., Kennicutt, Jr. R. C., 2004, *ARA&A*, 42, 603
- Krajnović D. et al., 2013, *MNRAS* in press (arXiv:1210.8167), (Paper XVII)
- Krajnović D. et al., 2013, *MNRAS* submitted, (Paper XXIV)
- Kroupa P., 2001, *MNRAS*, 322, 231
- Kroupa P., 2002, *Science*, 295, 82
- Kroupa P., Weidner C., 2003, *ApJ*, 598, 1076
- Kroupa P., Weidner C., Pflamm-Altenburg J., Thies I., Dabringhausen J., Marks M., Maschberger T., 2013, in *Planets, stars and stellar systems*, Oswalt T. D., Gilmore G., eds., Vol. 5, (Berlin: Springer), pp. 115–242 (arXiv:1112.3340)
- Kuntschner H. et al., 2006, *MNRAS*, 369, 497
- Kuntschner H., Emsellem E., et al., 2010, *MNRAS*, 408, 97
- Lablanche P.-Y. et al., 2012, *MNRAS*, 424, 1495 (Paper XII)
- Lauer T. R. et al., 2007a, *ApJ*, 664, 226
- Lauer T. R. et al., 2007b, *ApJ*, 662, 808
- Law D. R., Shapley A. E., Steidel C. C., Reddy N. A., Christensen C. R., Erb D. K., 2012, *Nature*, 487, 338
- Longhetti M., Saracco P., 2009, *MNRAS*, 394, 774
- Magorrian J. et al., 1998, *AJ*, 115, 2285
- Mancini C. et al., 2010, *MNRAS*, 401, 933
- Maraston C., 2005, *MNRAS*, 362, 799
- Maraston C., Daddi E., Renzini A., Cimatti A., Dickinson M., Papovich C., Pasquali A., Pirzkal N., 2006, *ApJ*, 652, 85
- Martig M., Bournaud F., Teyssier R., Dekel A., 2009, *ApJ*, 707, 250
- Matković A., Guzmán R., 2005, *MNRAS*, 362, 289
- McConnell N. J., Ma C.-P., Gebhardt K., Wright S. A., Murphy J. D., Lauer T. R., Graham J. R., Richstone D. O., 2011, *Nature*, 480, 215
- Mei S. et al., 2007, *ApJ*, 655, 144
- Meurer G. R. et al., 2009, *ApJ*, 695, 765
- Mihos J. C., Hernquist L., 1994, *ApJ*, 431, L9
- Misgeld I., Hilker M., 2011, *MNRAS*, 414, 3699
- Misgeld I., Hilker M., Mieske S., 2009, *A&A*, 496, 683
- Misgeld I., Mieske S., Hilker M., 2008, *A&A*, 486, 697
- Moster B. P., Somerville R. S., Maulbetsch C., van den Bosch F. C., Macciò A. V., Naab T., Oser L., 2010, *ApJ*, 710, 903
- Naab T., Johansson P. H., Ostriker J. P., 2009, *ApJ*, 699, L178
- Napolitano N. R., Romanowsky A. J., Tortora C., 2010, *MNRAS*, 405, 2351
- Navarro J. F., Frenk C. S., White S. D. M., 1996, *ApJ*, 462, 563
- Newman A. B., Ellis R. S., Bundy K., Treu T., 2012, *ApJ*, 746, 162
- Nipoti C., Londrillo P., Ciotti L., 2003, *MNRAS*, 342, 501
- Nipoti C., Treu T., Auger M. W., Bolton A. S., 2009, *ApJ*, 706, L86
- Onodera M. et al., 2012, *ApJ*, 755, 26
- Oser L., Naab T., Ostriker J. P., Johansson P. H., 2012, *ApJ*, 744, 63
- Oser L., Ostriker J. P., Naab T., Johansson P. H., Burkert A., 2010, *ApJ*, 725, 2312
- Padmanabhan N. et al., 2004, *New Astr.*, 9, 329
- Paturel G., Petit C., Prugniel P., Theureau G., Rousseau J., Brouty M., Dubois P., Cambrésy L., 2003, *A&A*, 412, 45
- Peng C. Y., 2007, *ApJ*, 671, 1098
- Press W. H., Teukolsky S. A., Vetterling W. T., Flannery B. P., 1992, *Numerical recipes in FORTRAN. The art of scientific computing*. Cambridge: University Press, —c1992, 2nd ed.
- Prugniel P., Simien F., 1996, *A&A*, 309, 749
- Renzini A., 2005, in *Astrophysics and Space Science Library*, Vol. 327, *The Initial Mass Function 50 Years Later*, Corbelli E., Palla F., Zinnecker H., eds., p. 221
- Renzini A., Ciotti L., 1993, *ApJ*, 416, L49+
- Robertson B., Cox T. J., Hernquist L., Franx M., Hopkins P. F., Martini P., Springel V., 2006, *ApJ*, 641, 21
- Rybicki G. B., 1987, in *IAU Symposium*, Vol. 127, *Structure and Dynamics of Elliptical Galaxies*, de Zeeuw P. T., ed., p. 397
- Salpeter E. E., 1955, *ApJ*, 121, 161
- Sánchez-Blázquez P. et al., 2006, *MNRAS*, 371, 703
- Sandage A., 1961, *The Hubble Atlas*. Carnegie Institution, Washington
- Sarzi M. et al., 2013, *MNRAS* in press (arXiv:1301.2589), (Paper XIX)
- Sarzi M. et al., 2010, *MNRAS*, 402, 2187
- Schechter P., 1976, *ApJ*, 203, 297
- Schiavon R. P., Barbuy B., Bruzual A. G., 2000, *ApJ*, 532, 453
- Schulz A. E., Mandelbaum R., Padmanabhan N., 2010, *MNRAS*, 408, 1463
- Scott N. et al., 2009, *MNRAS*, 398, 1835
- Scott N. et al., 2013, *MNRAS* in press (arXiv:1211.4615), (Paper XXI)
- Sersic J. L., 1968, *Atlas de galaxias australes*. Cordoba, Argentina: Observatorio Astronomico, 1968
- Shankar F., Bernardi M., 2009, *MNRAS*, 396, L76
- Shen S. et al., 2003, *MNRAS*, 343, 978
- Silk J., Rees M. J., 1998, *A&A*, 331, L1
- Silverman B. W., 1986, *Density estimation for statistics and data analysis*, Vol. 26. Chapman & Hall/CRC, Boca Raton
- Skrutskie M. F. et al., 2006, *AJ*, 131, 1163
- Smith R. J., Lucey J. R., Carter D., 2012, *MNRAS*, 426, 2994
- Sonnenfeld A., Treu T., Gavazzi R., Marshall P. J., Auger M. W., Suyu S. H., Koopmans L. V. E., Bolton A. S., 2012, *ApJ*, 752, 163
- Spiniello C., Trager S. C., Koopmans L. V. E., Chen Y. P., 2012, *ApJ*, 753, L32
- Spinrad H., Taylor B. J., 1971, *ApJS*, 22, 445

- Spitzer, Jr. L., Baade W., 1951, *ApJ*, 113, 413
- Suyu S. H. et al., 2012, *ApJ*, 750, 10
- Tacconi L. J. et al., 2010, *Nature*, 463, 781
- Thomas D., Maraston C., Bender R., Mendes de Oliveira C., 2005, *ApJ*, 621, 673
- Thomas J., Saglia R. P., Bender R., Thomas D., Gebhardt K., Magorrian J., Corsini E. M., Wegner G., 2009, *ApJ*, 691, 770
- Thomas J. et al., 2011, *MNRAS*, 415, 545
- Tonry J. L., Dressler A., Blakeslee J. P., Ajhar E. A., Fletcher A. B., Lupino G. A., Metzger M. R., Moore C. B., 2001, *ApJ*, 546, 681
- Tortora C., Napolitano N. R., Romanowsky A. J., Capaccioli M., Covone G., 2009, *MNRAS*, 396, 1132
- Tortora C., Napolitano N. R., Romanowsky A. J., Jetzer P., 2010, *ApJ*, 721, L1
- Tortora C., Romanowsky A. J., Napolitano N. R., 2013, *ApJ*, 765, 8
- Tremblay B., Merritt D., 1996, *AJ*, 111, 2243
- Treu T., Auger M. W., Koopmans L. V. E., Gavazzi R., Marshall P. J., Bolton A. S., 2010, *ApJ*, 709, 1195
- Treu T., Dutton A. A., Auger M. W., Marshall P. J., Bolton A. S., Brewer B. J., Koo D. C., Koopmans L. V. E., 2011, *MNRAS*, 417, 1601
- Treu T. et al., 2005, *ApJ*, 633, 174
- Trujillo I., Conselice C. J., Bundy K., Cooper M. C., Eisenhardt P., Ellis R. S., 2007, *MNRAS*, 382, 109
- Trujillo I., Ferreras I., de La Rosa I. G., 2011, *MNRAS*, 938
- Trujillo I. et al., 2006, *MNRAS*, 373, L36
- Valentinuzzi T. et al., 2010, *ApJ*, 712, 226
- van de Ven G., Falcón-Barroso J., McDermid R. M., Cappellari M., Miller B. W., de Zeeuw P. T., 2010, *ApJ*, 719, 1481
- van den Bergh S., 1976, *ApJ*, 206, 883
- van den Bosch F. C., 1997, *MNRAS*, 287, 543
- van der Marel R. P., van Dokkum P. G., 2007, *ApJ*, 668, 756
- van der Wel A., Bell E. F., van den Bosch F. C., Gallazzi A., Rix H.-W., 2009a, *ApJ*, 698, 1232
- van der Wel A., Rix H.-W., Holden B. P., Bell E. F., Robaina A. R., 2009b, *ApJ*, 706, L120
- van der Wel A. et al., 2011, *ApJ*, 730, 38
- van Dokkum P., 2011, *Nature*, 473, 160
- van Dokkum P. G., 2008, *ApJ*, 674, 29
- van Dokkum P. G., Conroy C., 2010, *Nature*, 468, 940
- van Dokkum P. G., Conroy C., 2011, *ApJ*, 735, L13
- van Dokkum P. G., Conroy C., 2012, *ApJ*, 760, 70
- van Dokkum P. G., Franx M., 1996, *MNRAS*, 281, 985
- van Dokkum P. G., Franx M., Kelson D. D., Illingworth G. D., 1998, *ApJ*, 504, L17
- van Dokkum P. G. et al., 2008, *ApJ*, 677, L5
- van Dokkum P. G. et al., 2010, *ApJ*, 709, 1018
- Vazdekis A., Ricciardelli E., Cenarro A. J., Rivero-González J. G., Díaz-García L. A., Falcón-Barroso J., 2012, *MNRAS*, 424, 157
- Wake D. A., van Dokkum P. G., Franx M., 2012, *ApJ*, 751, L44
- Wegner G. A., Corsini E. M., Thomas J., Saglia R. P., Bender R., Pu S. B., 2012, *AJ*, 144, 78
- Weijmans A.-M. et al., 2013, *MNRAS* submitted, (Paper XXIII)
- Wilkins S. M., Hopkins A. M., Trentham N., Tojeiro R., 2008, *MNRAS*, 391, 363
- Williams M. J., Bureau M., Cappellari M., 2009, *MNRAS*, 400, 1665
- Williams M. J., Bureau M., Cappellari M., 2010, *MNRAS*, 409, 1330
- Williams R. J., Quadri R. F., Franx M., van Dokkum P., Toft S., Kriek M., Labbé I., 2010, *ApJ*, 713, 738
- Wolf J., Martinez G. D., Bullock J. S., Kaplinghat M., Geha M., Muñoz R. R., Simon J. D., Avedo F. F., 2010, *MNRAS*, 406, 1220
- Worthey G., 1994, *ApJS*, 95, 107
- Wuyts S. et al., 2011, *ApJ*, 742, 96
- Wuyts S., Franx M., Cox T. J., Hernquist L., Hopkins P. F., Robertson B. E., van Dokkum P. G., 2009, *ApJ*, 696, 348
- York D. G., Adelman J., Anderson, Jr. J. E., et al., 2000, *AJ*, 120, 1579
- Young C. K., Currie M. J., 1994, *MNRAS*, 268, L11
- Young L. M. et al., 2011, *MNRAS*, 414, 940 (Paper IV)
- Zaritsky D., Gonzalez A. H., Zabludoff A. I., 2006, *ApJ*, 638, 725
- Zepf S. E., Silk J., 1996, *ApJ*, 466, 114
- Zhu G., Blanton M. R., Moustakas J., 2010, *ApJ*, 722, 491

Table 1. Central velocity dispersion, dark matter fraction and M/L for the ATLAS^{3D} sample of 260 early-type galaxies.

Galaxy	$\log \sigma(R_e/8)$ (km s^{-1})	$f_{\text{DM}}(r = R_e)$	$\log(M/L)_{\text{stars}}$ ($M_{\odot}/L_{\odot,r}$)	$\log(M/L)_{\text{Salp}}$ ($M_{\odot}/L_{\odot,r}$)
(1)	(2)	(3)	(4)	(5)
IC0560	1.920	0.12	0.429	0.381
IC0598	2.001	0.24	0.268	0.364
IC0676	1.825	0.20	0.481	0.261
IC0719	1.934	0.11	0.833	0.518
IC0782	1.818	0.57	0.318	0.510
IC1024	1.772	0.15	0.570	0.194
IC3631	1.733	0.00	-0.153	0.109
NGC0448	1.988	0.14	0.396	0.673
NGC0474	2.223	0.00	0.620	0.724
NGC0502	2.107	0.00	0.466	0.725
NGC0509	1.792	0.07	0.495	0.388
NGC0516	1.743	0.16	0.421	0.573
NGC0524	2.386	0.43	0.625	0.840
NGC0525	1.954	0.00	0.710	0.706
NGC0661	2.279	0.08	0.929	0.754
NGC0680	2.316	0.08	0.713	0.739
NGC0770	2.045	0.19	0.362	0.696
NGC0821	2.300	0.18	0.734	0.773
NGC0936	2.292	0.38	0.504	0.824
NGC1023	2.316	0.00	0.531	0.827
NGC1121	2.235	0.13	0.706	0.786
NGC1222	1.840	0.76	0.219	-0.014
NGC1248	1.965	0.00	0.326	0.488
NGC1266	1.897	0.00	0.598	0.364
NGC1289	2.133	0.14	0.577	0.578
NGC1665	1.999	0.12	0.450	0.567
NGC2481	2.271	0.06	0.642	0.706
NGC2549	2.174	0.17	0.666	0.710
NGC2577	2.320	0.06	0.843	0.790
NGC2592	2.384	0.01	0.882	0.812
NGC2594	2.263	0.07	0.659	0.729
NGC2679	1.983	0.17	0.423	0.576
NGC2685	1.972	0.18	0.327	0.620
NGC2695	2.342	0.11	0.672	0.795
NGC2698	2.355	0.00	0.749	0.802
NGC2699	2.181	0.07	0.538	0.724
NGC2764	1.954	0.00	0.649	0.154
NGC2768	2.306	0.00	0.933	0.772
NGC2778	2.215	0.04	0.842	0.768
NGC2824	2.127	0.45	0.262	0.343
NGC2852	2.306	0.00	0.834	0.784
NGC2859	2.252	0.06	0.546	0.732
NGC2880	2.150	0.25	0.580	0.675
NGC2950	2.230	0.11	0.539	0.661
NGC2962	2.244	0.00	0.813	0.782
NGC2974	2.375	0.07	0.950	0.787
NGC3032	1.983	0.42	0.164	-0.070
NGC3073	1.755	0.44	0.105	-0.145
NGC3098	2.038	0.18	0.517	0.612
NGC3156	1.838	0.23	0.245	0.132
NGC3182	2.060	0.48	0.336	0.606
NGC3193	2.299	0.08	0.532	0.794
NGC3226	2.250	0.00	0.873	0.784
NGC3230	2.303	0.04	0.754	0.795
NGC3245	2.322	0.11	0.588	0.727
NGC3248	1.990	0.15	0.420	0.527

Table 1 (cont'd)

Galaxy	$\log \sigma(R_e/8)$ (km s^{-1})	$f_{\text{DM}}(r = R_e)$	$\log(M/L)_{\text{stars}}$ ($M_{\odot}/L_{\odot r}$)	$\log(M/L)_{\text{Salp}}$ ($M_{\odot}/L_{\odot r}$)
(1)	(2)	(3)	(4)	(5)
NGC3301	2.096	0.21	0.297	0.448
NGC3377	2.163	0.06	0.553	0.731
NGC3379	2.329	0.33	0.600	0.838
NGC3384	2.208	0.22	0.404	0.724
NGC3400	1.878	0.18	0.522	0.687
NGC3412	2.022	0.47	0.211	0.648
NGC3414	2.365	0.14	0.714	0.826
NGC3457	1.889	0.21	0.173	0.517
NGC3458	2.253	0.05	0.627	0.767
NGC3489	2.028	0.24	0.113	0.394
NGC3499	1.845	0.25	0.189	0.449
NGC3522	2.011	0.24	0.582	0.648
NGC3530	2.049	0.23	0.493	0.695
NGC3595	2.206	0.00	0.602	0.749
NGC3599	1.874	0.11	0.234	0.467
NGC3605	1.966	0.00	0.459	0.668
NGC3607	2.360	0.07	0.661	0.799
NGC3608	2.286	0.09	0.669	0.829
NGC3610	2.244	0.12	0.409	0.566
NGC3613	2.327	0.00	0.772	0.761
NGC3619	2.221	0.28	0.615	0.693
NGC3626	2.134	0.23	0.332	0.261
NGC3630	2.241	0.13	0.550	0.745
NGC3640	2.267	0.17	0.534	0.722
NGC3641	2.241	0.34	0.821	0.773
NGC3648	2.277	0.21	0.694	0.807
NGC3658	2.216	0.00	0.572	0.721
NGC3665	2.352	0.14	0.748	0.764
NGC3674	2.346	0.04	0.811	0.809
NGC3694	1.892	0.60	0.082	0.437
NGC3757	2.170	0.20	0.610	0.705
NGC3796	1.955	0.34	0.243	0.387
NGC3838	2.144	0.00	0.589	0.694
NGC3941	2.144	0.00	0.400	0.693
NGC3945	2.262	0.22	0.549	0.760
NGC3998	2.445	0.15	0.920	0.818
NGC4026	2.279	0.08	0.633	0.747
NGC4036	2.276	0.00	0.697	0.783
NGC4078	2.242	0.07	0.823	0.755
NGC4111	2.201	0.00	0.651	0.616
NGC4119	1.771	0.20	0.423	0.306
NGC4143	2.336	0.16	0.646	0.834
NGC4150	1.945	0.20	0.336	0.358
NGC4168	2.241	0.25	0.775	0.738
NGC4179	2.264	0.08	0.724	0.771
NGC4191	2.127	0.22	0.533	0.685
NGC4203	2.202	0.00	0.536	0.827
NGC4215	2.147	0.04	0.589	0.652
NGC4233	2.337	0.22	0.724	0.812
NGC4249	1.885	0.78	-0.071	0.672
NGC4251	2.137	0.08	0.360	0.684
NGC4255	2.255	0.04	0.785	0.743
NGC4259	2.060	0.51	0.212	0.728
NGC4261	2.469	0.28	0.820	0.859
NGC4262	2.290	0.02	0.751	0.782
NGC4264	2.021	0.31	0.464	0.668

Table 1 (cont'd)

Galaxy	$\log \sigma(R_e/8)$ (km s^{-1})	$f_{\text{DM}}(r = R_e)$	$\log(M/L)_{\text{stars}}$ ($M_{\odot}/L_{\odot r}$)	$\log(M/L)_{\text{Salp}}$ ($M_{\odot}/L_{\odot r}$)
(1)	(2)	(3)	(4)	(5)
NGC4267	2.193	0.17	0.548	0.831
NGC4268	2.194	0.20	0.744	0.672
NGC4270	2.153	0.00	0.542	0.669
NGC4278	2.395	0.26	0.734	0.846
NGC4281	2.409	0.00	0.956	0.788
NGC4283	2.078	0.00	0.568	0.770
NGC4324	1.969	0.32	0.310	0.620
NGC4339	2.093	0.00	0.684	0.728
NGC4340	2.030	0.34	0.484	0.700
NGC4342	2.403	0.01	1.018	0.822
NGC4346	2.162	0.22	0.517	0.718
NGC4350	2.300	0.08	0.717	0.826
NGC4365	2.408	0.18	0.688	0.862
NGC4371	2.142	0.27	0.596	0.789
NGC4374	2.460	0.15	0.761	0.846
NGC4377	2.164	0.53	0.284	0.733
NGC4379	2.069	0.00	0.600	0.735
NGC4382	2.264	0.50	0.438	0.578
NGC4387	2.024	0.00	0.574	0.759
NGC4406	2.336	0.17	0.717	0.816
NGC4417	2.183	0.13	0.569	0.771
NGC4425	1.921	0.14	0.529	0.638
NGC4429	2.292	0.00	0.787	0.824
NGC4434	2.116	0.00	0.416	0.753
NGC4435	2.214	0.13	0.518	0.734
NGC4442	2.288	0.00	0.672	0.838
NGC4452	1.771	0.00	0.706	0.679
NGC4458	2.032	0.15	0.460	0.740
NGC4459	2.252	0.20	0.572	0.721
NGC4461	2.173	0.06	0.621	0.792
NGC4472	2.460	0.00	0.746	0.856
NGC4473	2.287	0.24	0.574	0.780
NGC4474	1.964	0.05	0.482	0.660
NGC4476	1.854	0.20	0.324	0.382
NGC4477	2.234	0.47	0.602	0.788
NGC4478	2.160	0.00	0.715	0.775
NGC4483	1.966	0.30	0.472	0.757
NGC4486	2.497	0.13	0.815	0.883
NGC4486A	2.156	0.00	0.656	0.705
NGC4489	1.812	0.77	-0.030	0.556
NGC4494	2.188	0.20	0.511	0.699
NGC4503	2.170	0.00	0.733	0.815
NGC4521	2.247	0.16	0.774	0.778
NGC4526	2.371	0.00	0.748	0.775
NGC4528	2.044	0.01	0.580	0.641
NGC4546	2.345	0.00	0.737	0.810
NGC4550	1.974	0.21	0.537	0.656
NGC4551	1.998	0.00	0.697	0.769
NGC4552	2.418	0.21	0.743	0.861
NGC4564	2.241	0.04	0.669	0.792
NGC4570	2.281	0.15	0.574	0.820
NGC4578	2.043	0.38	0.514	0.739
NGC4596	2.179	0.18	0.638	0.776
NGC4608	2.139	0.00	0.633	0.766
NGC4612	1.934	0.56	0.168	0.448
NGC4621	2.350	0.03	0.759	0.846

Table 1 (cont'd)

Galaxy	$\log \sigma(R_e/8)$ (km s^{-1})	$f_{\text{DM}}(r = R_e)$	$\log(M/L)_{\text{stars}}$ ($M_{\odot}/L_{\odot r}$)	$\log(M/L)_{\text{Salp}}$ ($M_{\odot}/L_{\odot r}$)
(1)	(2)	(3)	(4)	(5)
NGC4623	1.835	0.50	0.309	0.723
NGC4624	2.156	0.25	0.582	0.794
NGC4636	2.300	0.39	0.789	0.844
NGC4638	2.080	0.05	0.433	0.703
NGC4643	2.184	0.14	0.635	0.755
NGC4649	2.498	0.29	0.774	0.871
NGC4660	2.338	0.10	0.601	0.817
NGC4684	1.837	0.10	0.304	0.571
NGC4690	2.052	0.00	0.590	0.542
NGC4694	1.792	0.00	0.150	-0.042
NGC4697	2.257	0.00	0.703	0.800
NGC4710	2.021	0.00	0.645	0.587
NGC4733	1.750	0.00	0.338	0.518
NGC4753	2.263	0.25	0.519	0.615
NGC4754	2.254	0.21	0.596	0.815
NGC4762	2.129	0.23	0.417	0.760
NGC4803	2.016	0.55	0.336	0.670
NGC5103	2.061	0.22	0.376	0.695
NGC5173	2.005	0.33	0.261	0.507
NGC5198	2.304	0.00	0.794	0.809
NGC5273	1.870	0.55	0.247	0.502
NGC5308	2.376	0.09	0.744	0.790
NGC5322	2.395	0.16	0.634	0.700
NGC5342	2.254	0.21	0.676	0.773
NGC5353	2.462	0.11	0.769	0.855
NGC5355	1.918	0.47	0.227	0.279
NGC5358	1.886	0.27	0.477	0.623
NGC5379	1.840	0.73	0.307	0.468
NGC5422	2.242	0.21	0.670	0.758
NGC5473	2.332	0.49	0.511	0.783
NGC5475	2.041	0.33	0.411	0.544
NGC5481	2.174	0.00	0.776	0.740
NGC5485	2.253	0.09	0.806	0.777
NGC5493	2.208	0.20	0.305	0.514
NGC5500	1.915	0.13	0.649	0.650
NGC5507	2.320	0.17	0.708	0.786
NGC5557	2.406	0.01	0.659	0.774
NGC5574	1.885	0.04	0.378	0.368
NGC5576	2.269	0.00	0.454	0.705
NGC5582	2.180	0.37	0.599	0.752
NGC5611	2.120	0.06	0.651	0.645
NGC5631	2.207	0.10	0.580	0.627
NGC5638	2.202	0.06	0.647	0.791
NGC5687	2.280	0.12	0.864	0.768
NGC5770	1.988	0.12	0.328	0.576
NGC5813	2.354	0.47	0.699	0.816
NGC5831	2.220	0.14	0.610	0.743
NGC5838	2.441	0.00	0.901	0.834
NGC5839	2.208	0.01	0.716	0.802
NGC5845	2.431	0.00	0.695	0.810
NGC5846	2.365	0.14	0.844	0.853
NGC5854	2.061	0.12	0.356	0.472
NGC5864	2.079	0.00	0.570	0.636
NGC5866	2.205	0.00	0.663	0.640
NGC5869	2.260	0.24	0.750	0.787
NGC6010	2.213	0.17	0.685	0.719

Table 1 (cont'd)

Galaxy	$\log \sigma(R_e/8)$ (km s^{-1})	$f_{\text{DM}}(r = R_e)$	$\log(M/L)_{\text{stars}}$ ($M_{\odot}/L_{\odot r}$)	$\log(M/L)_{\text{Salp}}$ ($M_{\odot}/L_{\odot r}$)
(1)	(2)	(3)	(4)	(5)
NGC6014	1.953	0.48	0.429	0.518
NGC6017	2.041	0.00	0.447	0.526
NGC6149	2.031	0.31	0.425	0.611
NGC6278	2.338	0.16	0.667	0.812
NGC6547	2.285	0.12	0.740	0.629
NGC6548	2.240	0.20	0.768	0.698
NGC6703	2.260	0.00	0.775	0.736
NGC6798	2.137	0.16	0.564	0.684
NGC7280	2.061	0.36	0.406	0.444
NGC7332	2.143	0.07	0.288	0.528
NGC7454	2.049	0.13	0.673	0.585
NGC7457	1.874	0.45	0.238	0.455
NGC7465	1.971	0.05	0.333	0.348
NGC7693	1.816	0.00	0.586	0.357
NGC7710	2.019	0.00	0.537	0.616
PGC016060	1.992	0.00	0.651	0.509
PGC028887	2.132	0.35	0.643	0.730
PGC029321	1.762	0.90	-0.541	0.362
PGC035754	2.067	0.00	0.577	0.634
PGC042549	2.004	0.14	0.422	0.506
PGC044433	2.123	0.10	0.688	0.724
PGC050395	1.869	0.74	-0.061	0.581
PGC051753	1.892	0.15	0.584	0.621
PGC054452	1.818	0.28	0.411	0.534
PGC056772	1.916	0.00	0.620	0.379
PGC058114	1.902	0.474
PGC061468	1.823	0.90	-0.280	0.415
PGC071531	1.969	0.479
PGC170172	2.020	0.16	0.111	0.650
UGC03960	1.946	0.04	0.729	0.606
UGC04551	2.265	0.04	0.658	0.737
UGC05408	1.919	0.00	0.194	0.331
UGC06062	2.156	0.25	0.629	0.760
UGC06176	2.045	0.50	0.331	0.328
UGC08876	2.152	0.00	0.760	0.740
UGC09519	1.992	0.10	0.513	0.345

Note. — Column (1): The Name is the principal designation from LEDA (Paturel et al. 2003), which is used as standard designation for our project. Column (2): Central stellar velocity dispersion (1σ random error of 5% or 0.021 dex) within a circular aperture of radius $R = R_e/8$, where R_e is the value given in table 1 of Paper XV. This $\sigma(R_e/8)$ is measured by co-adding all SAURON spectra contained within the aperture. The velocity dispersion is measured on that single spectrum using pPXF and adopting a Gaussian line-of-sight velocity distribution (keyword MOMENTS=2). Column (3): fraction of dark matter contained within an iso-surface of volume $V = 4\pi R_e^3/3$ (a sphere of radius $r = R_e$ for spherical galaxies). Column (4): Mass-to-light ratio (1σ random error of 6% or 0.027 dex) of the stellar component (approximately within a sphere of radius $r \sim R_e$) in the SDSS r -band, for the best-fitting JAM model with NFW halo (model B in Paper XV) for the assumed distance and extinction of table 3 of Paper I. A description of the quality of the model fits is given in table 1 of Paper XV. The $(M/L)_{\text{stars}}$ and f_{DM} of models with qual < 2 should be used with caution. Column (5): Mass-to-light ratio of the stellar population (1σ random error of 6% or 0.027 dex) within $1R_e$, in the SDSS r -band, assuming a Salpeter IMF slope from 0.1 to $100 M_{\odot}$ and using the models of Vazdekis et al. (2012). This value is mass weighted over the multiple populations fitted to the SAURON spectra with pPXF. Table 1 is also available from our project website <http://purl.org/atlas3d>.