

Planck 2013 results. XXII. Constraints on inflation

Planck Collaboration: P. A. R. Ade⁸⁶, N. Aghanim⁵⁹, C. Armitage-Caplan⁹², M. Arnaud⁷², M. Ashdown^{69,6}, F. Atrio-Barandela¹⁹, J. Aumont⁵⁹, C. Baccigalupi⁸⁵, A. J. Banday^{95,10}, R. B. Barreiro⁶⁶, J. G. Bartlett^{1,67}, N. Bartolo³³, E. Battaner⁹⁶, K. Benabed^{60,94}, A. Benoît⁵⁷, A. Benoit-Lévy^{26,60,94}, J.-P. Bernard¹⁰, M. Bersanelli^{36,50}, P. Bielewicz^{95,10,85}, J. Bobin⁷², J. J. Bock^{67,11}, A. Bonaldi⁶⁸, J. R. Bond⁹, J. Borrill^{14,89}, F. R. Bouchet^{60,94}, M. Bridges^{69,6,63}, M. Bucher^{1*}, C. Burigana^{49,34}, R. C. Butler⁴⁹, E. Calabrese⁹², J.-F. Cardoso^{73,1,60}, A. Catalano^{74,71}, A. Challinor^{63,69,12}, A. Chamballu^{72,16,59}, L.-Y. Chiang⁶², H. C. Chiang^{28,7}, P. R. Christensen^{81,39}, S. Church⁹¹, D. L. Clements⁵⁵, S. Colombi^{60,94}, L. P. L. Colombo^{25,67}, F. Couchot⁷⁰, A. Coulais⁷¹, B. P. Crill^{67,82}, A. Curto^{6,66}, F. Cuttaia⁴⁹, L. Danese⁸⁵, R. D. Davies⁶⁸, R. J. Davis⁶⁸, P. de Bernardis³⁵, A. de Rosa⁴⁹, G. de Zotti^{46,85}, J. Delabrouille¹, J.-M. Delouis^{60,94}, F.-X. Désert⁵³, C. Dickinson⁶⁸, J. M. Diego⁶⁶, H. Dole^{59,58}, S. Donzelli⁵⁰, O. Doré^{67,11}, M. Douspis⁵⁹, J. Dunkley⁹², X. Dupac⁴², G. Efstathiou⁶³, T. A. Enßlin⁷⁸, H. K. Eriksen⁶⁴, F. Finelli^{49,51*}, O. Forni^{95,10}, M. Frailis⁴⁸, E. Franceschi⁴⁹, S. Galeotta⁴⁸, K. Ganga¹, C. Gauthier^{1,77}, M. Giard^{95,10}, G. Giardini⁴³, Y. Giraud-Héraud¹, J. González-Nuevo^{66,85}, K. M. Górski^{67,98}, S. Gratton^{69,63}, A. Gregorio^{37,48}, A. Gruppuso⁴⁹, J. Hamann⁹³, F. K. Hansen⁶⁴, D. Hanson^{79,67,9}, D. Harrison^{63,69}, S. Henrot-Versillé⁷⁰, C. Hernández-Monteagudo^{13,78}, D. Herranz⁶⁶, S. R. Hildebrandt¹¹, E. Hivon^{60,94}, M. Hobson⁶, W. A. Holmes⁶⁷, A. Hornstrup¹⁷, W. Hovest⁷⁸, K. M. Huffenberger⁹⁷, T. R. Jaffe^{95,10}, A. H. Jaffe⁵⁵, W. C. Jones²⁸, M. Juvela²⁷, E. Keihänen²⁷, R. Kesitalo^{23,14}, T. S. Kisner⁷⁶, R. Kneissl^{41,8}, J. Knoche⁷⁸, L. Knox³⁰, M. Kunz^{18,59,3}, H. Kurki-Suonio^{27,45}, G. Lagache⁵⁹, A. Lähteenmäki^{2,45}, J.-M. Lamarre⁷¹, A. Lasenby^{6,69}, R. J. Laureijs⁴³, C. R. Lawrence⁶⁷, S. Leach⁸⁵, J. P. Leahy⁶⁸, R. Leonardi⁴², J. Lesgourgues^{93,84}, M. Liguori³³, P. B. Lilje⁶⁴, M. Linden-Vørnle¹⁷, M. López-Cañiego⁶⁶, P. M. Lubin³¹, J. F. Macías-Pérez⁷⁴, B. Maffei⁶⁸, D. Maino^{36,50}, N. Mandolesi^{49,5,34}, M. Maris⁴⁸, D. J. Marshall⁷², P. G. Martin⁹, E. Martínez-González⁶⁶, S. Masi³⁵, S. Matarrese³³, F. Matthai⁷⁸, P. Mazzotta³⁸, P. R. Meinhold³¹, A. Melchiorri^{35,52}, L. Mendes⁴², A. Mennella^{36,50}, M. Migliaccio^{63,69}, S. Mitra^{54,67}, M.-A. Miville-Deschênes^{59,9}, A. Moneti⁶⁰, L. Montier^{95,10}, G. Morgante⁴⁹, D. Mortlock⁵⁵, A. Moss⁸⁷, D. Munshi⁸⁶, P. Naselsky^{81,39}, F. Nati³⁵, P. Natoli^{34,4,49}, C. B. Netterfield²¹, H. U. Nørgaard-Nielsen¹⁷, F. Noviello⁶⁸, D. Novikov⁵⁵, I. Novikov⁸¹, I. J. O'Dwyer⁶⁷, S. Osborne⁹¹, C. A. Oxborrow¹⁷, F. Paci⁸⁵, L. Pagano^{35,52}, F. Pajot⁵⁹, R. Paladini⁵⁶, S. Pandolfi³⁸, D. Paoletti^{49,51}, B. Partridge⁴⁴, F. Pasian⁴⁸, G. Patanchon¹, H. V. Peiris²⁶, O. Perdereau⁷⁰, L. Perotto⁷⁴, F. Perrotta⁸⁵, F. Piacentini³⁵, M. Piat¹, E. Pierpaoli²⁵, D. Pietrobon⁶⁷, S. Plaszczynski⁷⁰, E. Pointecouteau^{95,10}, G. Polenta^{4,47}, N. Ponthieu^{59,53}, L. Popa⁶¹, T. Poutanen^{45,27,2}, G. W. Pratt⁷², G. Prézeau^{11,67}, S. Prunet^{60,94}, J.-L. Puget⁵⁹, J. P. Rachen^{22,78}, R. Rebolo^{65,15,40}, M. Reinecke⁷⁸, M. Remazeilles^{59,1}, C. Renault⁷⁴, S. Ricciardi⁴⁹, T. Riller⁷⁸, I. Ristorcelli^{95,10}, G. Rocha^{67,11}, C. Rosset¹, G. Roudier^{1,71,67}, M. Rowan-Robinson⁵⁵, J. A. Rubiño-Martín^{65,40}, B. Rusholme⁵⁶, M. Sandri⁴⁹, D. Santos⁷⁴, M. Savelainen^{27,45}, G. Savini⁸³, D. Scott²⁴, M. D. Seiffert^{67,11}, E. P. S. Shellard¹², L. D. Spencer⁸⁶, J.-L. Starck⁷², V. Stolyarov^{6,69,90}, R. Stompor¹, R. Sudiwala⁸⁶, R. Sunyaev^{78,88}, F. Sureau⁷², D. Sutton^{63,69}, A.-S. Suur-Uski^{27,45}, J.-F. Sygnet⁶⁰, J. A. Tauber⁴³, D. Tavagnacco^{48,37}, L. Terenzi⁴⁹, L. Toffolatti^{20,66}, M. Tomasi⁵⁰, M. Tristram⁷⁰, M. Tucci^{18,70}, J. Tuovinen⁸⁰, L. Valenziano⁴⁹, J. Valiviita^{45,27,64}, B. Van Tent⁷⁵, J. Varis⁸⁰, P. Vielva⁶⁶, F. Villa⁴⁹, N. Vittorio³⁸, L. A. Wade⁶⁷, B. D. Wandelt^{60,94,32}, M. White²⁹, A. Wilkinson⁶⁸, D. Yvon¹⁶, A. Zacchei⁴⁸, and A. Zonca³¹

(Affiliations can be found after the references)

Preprint online version: March 22, 2013

ABSTRACT

We analyse the implications of the *Planck* data for cosmic inflation. The *Planck* nominal mission temperature anisotropy measurements, combined with the *WMAP* large-angle polarization, constrain the scalar spectral index to $n_s = 0.9603 \pm 0.0073$, ruling out exact scale invariance at over 5σ . *Planck* establishes an upper bound on the tensor-to-scalar ratio at $r < 0.11$ (95% CL). The *Planck* data shrink the space of allowed standard inflationary models, preferring potentials with $V'' < 0$. Exponential potential models, the simplest hybrid inflationary models, and monomial potential models of degree $n \geq 2$ do not provide a good fit to the data. *Planck* does not find statistically significant running of the scalar spectral index, obtaining $dn_s/d\ln k = -0.0134 \pm 0.0090$. We verify these conclusions through a numerical analysis, which makes no slow-roll approximation, and carry out a Bayesian parameter estimation and model-selection analysis for a number of inflationary models including monomial, natural, and hilltop potentials. For each model, we present the *Planck* constraints on the parameters of the potential and explore several possibilities for the post-inflationary entropy generation epoch, thus obtaining nontrivial data-driven constraints. We also present a direct reconstruction of the observable range of the inflaton potential. Unless a quartic term is allowed in the potential, we find results consistent with second-order slow-roll predictions. We investigate whether the primordial power spectrum contains any features. A penalized likelihood approach suggests a feature near the highest wavenumbers probed by *Planck* at an estimated significance of $\sim 3\sigma$ after correction for the “look elsewhere” effect. We find that models with a parameterized oscillatory feature improve the fit by $\Delta\chi^2_{\text{eff}} \approx 10$; however, Bayesian evidence does not prefer these models. We constrain several single-field inflation models with generalized Lagrangians by combining power spectrum data with *Planck* bounds on f_{NL} . *Planck* constrains with unprecedented accuracy the amplitude and possible correlation of non-decaying isocurvature fluctuations. The fractional primordial contributions of CDM isocurvature modes of the types expected in the curvaton and axion scenarios have upper bounds of 0.25% and 3.9% (95% CL), respectively. In models with arbitrarily correlated CDM or neutrino isocurvature modes, an anticorrelated isocurvature component can improve the χ^2_{eff} by ~ 4 due to a moderate tension between $l < 40$ and higher multipoles. Nonetheless, the data are consistent with adiabatic initial conditions.

Key words. Cosmology: theory – early Universe – inflation

1. Introduction

This paper, one of a set associated with the 2013 release of data from the *Planck*¹ mission (Planck Collaboration I, 2013), describes the implications of the *Planck* measurement of cosmic microwave background (CMB) anisotropies for cosmic inflation. In this first release only the *Planck* temperature data resulting from the ‘nominal’ mission is used, which includes 2.6 surveys of the sky. The interpretation of the CMB polarization as seen by *Planck* will be in a later series of publications. This paper exploits the data presented in Planck Collaboration XV (2013); Planck Collaboration XVII (2013). Other closely related papers are the estimates of cosmological parameters in Planck Collaboration XVI (2013) and investigations of non-Gaussianity in Planck Collaboration XXIV (2013).

In the early 1980s inflationary cosmology, which postulates an epoch of nearly exponential expansion, was proposed in order to resolve a number of puzzles of standard Big Bang cosmology such as the entropy, flatness, and monopole problems (Starobinsky, 1980; Guth, 1981; Albrecht & Steinhardt, 1982; Linde, 1982, 1983). During inflation, cosmological fluctuations resulting from quantum fluctuations are generated and can be calculated using the semiclassical theory of quantum fields in curved spacetime (Mukhanov & Chibisov, 1981, 1982; Hawking, 1982; Guth & Pi, 1982; Starobinsky, 1982; Bardeen et al., 1983; Mukhanov, 1985).

Cosmological observations pre-*Planck* are consistent with the simplest models of inflation within the slow-roll paradigm. Recent observations of the CMB anisotropies (Story et al., 2012; Bennett et al., 2012; Hinshaw et al., 2012a; Hou et al., 2012; Das et al., 2013) and of large-scale structure (Beutler et al., 2011; Padmanabhan et al., 2012; Anderson et al., 2012) indicate that our Universe is very close to spatially flat and has primordial density fluctuations that are nearly Gaussian and adiabatic and are described by a nearly scale-invariant power spectrum. Pre-*Planck* CMB observations also indicate that the amplitude of gravitational waves, with a nearly scale-invariant spectrum generated in the early Universe (Grishchuk, 1975; Starobinsky, 1979; Rubakov et al., 1982; Fabbri & Pollock, 1983), is small.

Most of the results in this paper are based on the two-point statistics of the CMB as measured by *Planck*, exploiting the data presented in Planck Collaboration XV (2013); Planck Collaboration XVI (2013) and Planck Collaboration XVII (2013). The *Planck* results testing the Gaussianity of the primordial CMB component are described in the companion papers Planck Collaboration XXIII (2013); Planck Collaboration XXIV (2013); Planck Collaboration XXV (2013). *Planck* finds values for the non-Gaussian f_{NL} parameter connected to the CMB bispectrum that are consistent with the Gaussian hypothesis (Planck Collaboration XXIV, 2013), and this result has important implications for inflation. The simplest slow-roll inflationary models predict a level of bispectral non-Gaussianity f_{NL} of the order of the slow-roll parameters and therefore too small to be detected by *Planck*.

The paper is organized as follows. Section 2 reviews inflationary theory emphasizing in particular those aspects used

later in the paper. In Section 3 the statistical methodology and the *Planck* and other astrophysical likelihoods are described. Section 4 studies constraints on slow-roll inflation and studies their robustness under generalizations of the minimal assumptions of our baseline cosmological model. In Section 5 Bayesian model comparison of several inflationary models taking into account the uncertainty from the end of inflation to the beginning of the radiation dominated era is carried out. Section 6 reconstructs the inflationary potential over the range corresponding to the scales observable in the CMB. In Section 7 a penalized likelihood reconstruction of the primordial perturbation spectrum is performed. Section 8 reports on a parametric search for oscillations and features in the primordial scalar power spectrum. Section 9 examines constraints on non-canonical single-field models of inflation including the f_{NL} measurements from Planck Collaboration XXIV (2013). In Section 10 constraints on isocurvature modes are established, thus testing the hypothesis that initial conditions were solely adiabatic. We summarize our conclusions in Section 11. Definitions of the most relevant symbols used in this paper can be found in Tables 1 and 2.

2. Lightning review of inflation

Before describing cosmic inflation, which was developed in the early 1980s, it is useful to review the state of theory prior to its introduction. Lifshitz & Khalatnikov (1963) first wrote down the equations for the evolution of linearized perturbations about a homogeneous and isotropic Friedmann-Lemaître-Robertson-Walker (FLRW) spacetime within the framework of general relativity. The general framework adopted was based on two assumptions:

- (i) the cosmological perturbations can be described by a single component fluid, at least at very early times;
- (ii) the initial cosmological perturbations were statistically homogeneous and isotropic, and also Gaussian.

These are the simplest—but by no means unique—assumptions for defining a stochastic process for the initial conditions. Assumption (i), where only a single *adiabatic* mode is excited, is just the simplest possibility; in Sect. 10 we shall describe isocurvature perturbations, where other available modes are excited, and report on the constraints established by *Planck*. Assumption (ii) is *a priori* more dubious given the understanding at the time. An appeal can be made to the fact that any physics at weak coupling could explain (ii), but at the time these assumptions were somewhat *ad hoc*.

Even with the strong assumptions (i) and (ii), comparisons with observations cannot be made without further restrictions on the functional form of the primordial power spectrum of large-scale curvature inhomogeneities \mathcal{R} , $\mathcal{P}_{\mathcal{R}}(k) \propto k^{n_s-1}$, where n_s is the (scalar) spectral index. The notion of a *scale-invariant* (i.e., $n_s = 1$) primordial power spectrum was introduced by Harrison (1970); Zeldovich (1972); and Peebles & Yu (1970) to address this problem. These authors showed that the scale-invariant power law was consistent with the crude constraints on large- and small-scale perturbations available at the time; however, other than its mathematical simplicity, no compelling theoretical explanation for this ansatz was put forth. An important current question, addressed in Sect. 4, is whether $n_s = 1$ (i.e., exact scale invariance) is consistent with the data, or whether there is convincing evidence for small deviations from exact scale invariance. Although the inflationary potential can be tuned to obtain $n_s = 1$, inflationary models generically predict deviations from $n_s = 1$, usually on the red side (i.e., $n_s < 1$).

* Corresponding authors: Martin Bucher bucher@apc.univ-paris7.fr, Fabio Finelli finelli@iasfbo.inaf.it

¹ *Planck* (<http://www.esa.int/Planck>) is a project of the European Space Agency (ESA) with instruments provided by two scientific consortia funded by ESA member states (in particular the lead countries France and Italy), with contributions from NASA (USA) and telescope reflectors provided by a collaboration between ESA and a scientific consortium led and funded by Denmark.

Table 1. Cosmological parameter definitions

Parameter	Definition
Ω_b	Baryon fraction today (compared to critical density)
Ω_c	Cold dark matter fraction today (compared to critical density)
θ_{MC}	Approximation to the angular size of sound horizon at last scattering
τ	Thomson scattering optical depth of reionized intergalactic medium
N_{eff}	Number of massive and massless neutrinos
Σm_ν	Sum of neutrino masses
Y_P	Fraction of baryonic mass in primordial helium
Ω_K	Spatial curvature parameter
w_{de}	Dark energy equation of state parameter (i.e., p/ρ) (assumed constant)
\mathcal{R}	Curvature perturbation
\mathcal{I}	Isocurvature perturbation
$\mathcal{P}_X = k^3 X_k ^2 / 2\pi^2$.	Power spectrum of X
A_X	X power spectrum amplitude (at $k_* = 0.05 \text{ Mpc}^{-1}$)
n_s	Scalar spectrum spectral index (at $k_* = 0.05 \text{ Mpc}^{-1}$)
$dn_s/d \ln k$	Running of scalar spectral index (at $k_* = 0.05 \text{ Mpc}^{-1}$)
$d^2 n_s / d \ln k^2$	Running of running of scalar spectral index (at $k_* = 0.05 \text{ Mpc}^{-1}$)
r	Tensor-to-scalar power ratio (at $k_* = 0.05 \text{ Mpc}^{-1}$)
n_t	Tensor spectrum spectral index (at $k_* = 0.05 \text{ Mpc}^{-1}$)
$dn_t / d \ln k$	Running of tensor spectral index (at $k_* = 0.05 \text{ Mpc}^{-1}$)

2.1. Cosmic inflation

Inflation was developed in a series of papers by [Starobinsky \(1980\)](#); [Guth \(1981\)](#); [Albrecht & Steinhardt \(1982\)](#); and [Linde \(1982, 1983\)](#). (See also [Sato \(1981\)](#) and [Brout et al. \(1978\)](#) for interesting precursors.) By generating an equation of state with a negative pressure (i.e., $w = p/\rho \approx -1$) before the radiation epoch, inflation solves a number of cosmological conundrums (the monopole, horizon, smoothness, and entropy problems), which had plagued all cosmological models extrapolating a matter-radiation equation of state all the way back to the singularity. Such an equation of state with a large negative pressure and the resulting nearly-exponential expansion are obtained from a scalar field, the *inflaton*, with a canonical kinetic term, slowly rolling in the framework of Einstein gravity.

The homogeneous evolution of the inflaton field, ϕ , is governed by the equation of motion

$$\ddot{\phi}(t) + 3H(t)\dot{\phi}(t) + V_\phi = 0, \quad (1)$$

and the Friedmann equation

$$H^2 = \frac{1}{3M_{\text{pl}}^2} \left(\frac{1}{2} \dot{\phi}^2 + V(\phi) \right). \quad (2)$$

Here $H = \dot{a}/a$ is the Hubble parameter, the subscript ϕ denotes the derivative with respect to ϕ , $M_{\text{pl}} = (8\pi G)^{-1/2}$ is the reduced Planck mass, and V is the potential. The evolution during the stage of quasi-exponential expansion, when the scalar field rolls slowly down the potential, can be approximated by neglecting the second time derivative in Eq. 1 and the kinetic energy term in Eq. 2, so that

$$3H(t)\dot{\phi}(t) \approx -V_\phi, \quad (3)$$

$$H^2 \approx \frac{V(\phi)}{3M_{\text{pl}}^2}. \quad (4)$$

Necessary conditions for the *slow-roll* described above are $\epsilon_V \ll 1$ and $|\eta_V| \ll 1$, where the slow-roll parameters ϵ_V and η_V are

defined as

$$\epsilon_V = \frac{M_{\text{pl}}^2 V_\phi^2}{2V^2}, \quad (5)$$

$$\eta_V = \frac{M_{\text{pl}}^2 V_{\phi\phi}}{V}. \quad (6)$$

An analogous hierarchy of parameters, called the Hubble flow function (HFF) slow-roll parameters, measures instead the deviation from an exact exponential expansion. This hierarchy is defined as $\epsilon_1 = -\dot{H}/H^2$, $\epsilon_{i+1} \equiv \dot{\epsilon}_i/(H\epsilon_i)$, with $i \geq 1$. By using Eq. 3, we have that $\epsilon_1 \approx \epsilon_V$, $\epsilon_2 \approx -2\eta_V + 4\epsilon_V$.

2.2. Quantum generation of fluctuations

Without quantum fluctuations, inflationary theory would fail. Classically, any initial spatial curvature or gradients in the scalar field, as well any inhomogeneities in other fields, would rapidly decay away during the quasi-exponential expansion. The resulting Universe would be too homogeneous and isotropic compared with observations. Quantum fluctuations must exist in order to satisfy the uncertainty relations that follow from the canonical commutation relations of quantum field theory. The quantum fluctuations in the inflaton, and in the transverse and traceless parts of the metric, are amplified by the nearly exponential expansion, yielding the scalar and tensor primordial power spectra, respectively.

Many essentially equivalent approaches to quantizing the linearized cosmological fluctuations can be found in the original literature (see, e.g., [Mukhanov & Chibisov, 1981](#); [Hawking, 1982](#); [Guth & Pi, 1982](#); [Starobinsky, 1982](#); [Bardeen et al., 1983](#)). A simple formalism, which we shall follow here, was introduced by [Mukhanov \(1988\)](#); [Mukhanov et al. \(1992\)](#) and [Sasaki \(1986\)](#). In this approach a gauge-invariant inflaton fluctuation Q is constructed and canonically quantized. This gauge-invariant variable Q is the inflaton fluctuation $\delta\phi(t, x)$ in the uniform curvature gauge. The mode function of the inflaton fluctuations $\delta\phi(t, x)$ obeys the evolution equation

$$(a\delta\phi_k)'' + \left(k^2 - \frac{z''}{z} \right) (a\delta\phi_k) = 0, \quad (7)$$

Table 2. Conventions and definitions for inflation physics

Parameter	Definition
ϕ	Inflaton
$V(\phi)$	Inflaton potential
a	Scale factor
t	Cosmic (proper) time
δX	Fluctuation of X
$\dot{X} = dX/dt$	Derivative with respect to proper time
$X' = dX/d\eta$	Derivative with respect to conformal time
$X_\phi = \partial X/\partial \phi$	Partial derivative with respect to ϕ
M_{pl}	Reduced Planck mass ($= 2.435 \times 10^{18}$ GeV)
\mathcal{Q}	Scalar perturbation variable
$h^{+, \times}$	Gravitational wave amplitude of (+, \times)-polarization component
X_*	X evaluated at Hubble exit during inflation of mode with wavenumber k_*
X_e	X evaluated at end of inflation
$\epsilon_V = M_{\text{pl}}^2 V_\phi^2 / 2V^2$	First slow-roll parameter for $V(\phi)$
$\eta_V = M_{\text{pl}}^2 V_{\phi\phi} / V^2$	Second slow-roll parameter for $V(\phi)$
$\xi_V^2 = M_{\text{pl}}^4 V_\phi V_{\phi\phi\phi} / V^2$	Third slow-roll parameter for $V(\phi)$
$\varpi_V^3 = M_{\text{pl}}^6 V_\phi^2 V_{\phi\phi\phi\phi} / V^3$	Fourth slow-roll parameter for $V(\phi)$
$\epsilon_1 = -\dot{H}/H^2$	First Hubble hierarchy parameter
$\epsilon_{n+1} = \dot{\epsilon}_n / H \epsilon_n$	$(n+1)$ st Hubble hierarchy parameter (where $n \geq 1$)
$N(t) = \int_t^{t_e} dt H$	Number of e -folds to end of inflation
$\delta\sigma$	Curvature field perturbation
δs	Isocurvature field perturbation

where $z = a\dot{\phi}/H$. The gauge-invariant field fluctuation is directly related to the comoving curvature perturbation²

$$\mathcal{R} = -H \frac{\delta\phi}{\dot{\phi}}. \quad (8)$$

Analogously, gravitational waves are described by the two polarization states (+, \times) of the transverse and traceless parts of the metric fluctuations and are amplified by the expansion of the Universe as well (Grishchuk, 1975). The evolution equation for their mode function is

$$(ah_k^{+, \times})'' + \left(k^2 - \frac{a''}{a}\right)(ah_k^{+, \times}) = 0. \quad (9)$$

Early discussions of the generation of gravitational waves during inflation include Starobinsky (1979); Rubakov et al. (1982); Fabbri & Pollock (1983); Abbott & Wise (1984).

Because the primordial perturbations are small, of order 10^{-5} , the linearized equations 7 and 9 provide an accurate description for the generation and subsequent evolution of the cosmological perturbations during inflation. In this paper we use two approaches for solving for the cosmological perturbations. Firstly, we use an approximate treatment based on the *slow-roll* approximation described below. Secondly, we use an almost exact approach based on numerical integration of the ordinary differential equations 7 and 9 for each value of the co-moving wavenumber k . For fixed k the evolution may be divided into three epochs: (i) sub-Hubble evolution, (ii) Hubble crossing evolution, and (iii) super-Hubble evolution. During (i) the wavelength is much smaller than the Hubble length, and the mode oscillates as it would in a non-expanding Universe (i.e., Minkowski space). Therefore we can proceed with quantization as we would

in Minkowski space. We quantize by singling out the positive frequency solution, as in the Bunch-Davies vacuum (Bunch & Davies, 1978). This epoch is the oscillating regime in the WKB approximation. In (iii), by contrast, there are two solutions, a growing and a decaying mode, and the evolution becomes independent of k . We care only about the growing mode. On scales much larger than the Hubble radius (i.e., $k \ll aH$), both curvature and tensor fluctuations admit solutions constant in time.³ All the interesting, or nontrivial, evolution takes place between (i) and (iii)—that is, during (ii), a few e -folds before and after Hubble crossing, and this is the interval where the numerical integration is most useful since the asymptotic expansions are not valid in this transition region. Two numerical codes are used in this paper, ModeCode (Adams et al., 2001; Peiris et al., 2003; Mortonson et al., 2009; Easther & Peiris, 2012), and the inflation module of Lesgourgues & Valkenburg (2007) as implemented in CLASS (Lesgourgues, 2011; Blas et al., 2011).⁴

It is convenient to expand the power spectra of curvature and tensor perturbations as

$$\mathcal{P}_{\mathcal{R}}(k) = A_s \left(\frac{k}{k_*}\right)^{n_s - 1 + \frac{1}{2} \frac{dn_s}{dn} \ln k \ln(k/k_*) + \frac{1}{6} \frac{d^2 n_s}{d^2 n} \ln^2 k \ln(k/k_*)^2 + \dots} \quad (10)$$

$$\mathcal{P}_t(k) = A_t \left(\frac{k}{k_*}\right)^{n_t + \frac{1}{2} \frac{dn_t}{dn} \ln k \ln(k/k_*) + \dots}, \quad (11)$$

where $A_s (A_t)$ is the scalar (tensor) amplitude and $n_s (n_t)$, $dn_s/d \ln k$ ($dn_t/d \ln k$) and $d^2 n_s/d \ln^2 k$ are the scalar (tensor) spectral index, the running of the scalar (tensor) spectral index, and the running of the running of the scalar spectral index, respectively.

² Another important quantity is the curvature perturbation on uniform density hypersurfaces ζ (in the Newtonian gauge, $\zeta = -\psi - H\delta\rho/\rho$, where ψ is the generalized gravitational potential) which is related to the perturbed spatial curvature $^{(3)}R = -4\nabla^2 \zeta/a^2$. On large scales $\zeta \approx R$.

³ On large scales, the curvature fluctuation is constant in time when non-adiabatic pressure terms are negligible. This condition is typically violated in multi-field inflationary models.

⁴ <http://zuserver2.star.ucl.ac.uk/~hiranya/ModeCode/>, <http://class-code.net>

The parameters of the scalar and tensor power spectra may be calculated approximately in the framework of the slow-roll approximation by evaluating the following equations at the value of the inflation field ϕ_* where the mode $k_* = a_* H_*$ crosses the Hubble radius for the first time. (For a nice review of the slow-roll approximation, see for example [Liddle & Lyth \(1993\)](#)). The number of e -folds before the end of inflation, N_* , at which the pivot scale k_* exits from the Hubble radius, is

$$N_* = \int_{t_*}^{t_e} dt H \approx \frac{1}{M_{\text{pl}}^2} \int_{\phi_*}^{\phi_e} d\phi \frac{V}{V_\phi}, \quad (12)$$

where the equality holds in the slow-roll approximation, and subscript ‘e’ refers to the end of inflation.

The coefficients of Eqs. 10 and 11 at their respective leading orders in the slow-roll parameters are given by

$$A_s \approx \frac{V}{24\pi^2 M_{\text{pl}}^4 \epsilon_V} \quad (13)$$

$$A_t \approx \frac{2V}{3\pi^2 M_{\text{pl}}^4} \quad (14)$$

$$n_s - 1 \approx 2\eta_V - 6\epsilon_V \quad (15)$$

$$n_t \approx -2\epsilon_V \quad (16)$$

$$dn_s/d \ln k \approx -16\epsilon_V \eta_V + 24\epsilon_V^2 + 2\xi_V^2 \quad (17)$$

$$dn_t/d \ln k \approx -4\epsilon_V \eta_V + 8\epsilon_V^2 \quad (18)$$

$$d^2 n_s/d \ln k^2 \approx -192\epsilon_V^3 + 192\epsilon_V^2 \eta_V - 32\epsilon_V \eta_V^2 - 24\epsilon_V \xi_V^2 + 2\eta_V \xi_V^2 + 2\varpi_V^3, \quad (19)$$

where the *slow-roll parameters* ϵ_V and η_V are defined in Eqs. 5 and 6, and the higher order parameters are defined as follows

$$\xi_V^2 = \frac{M_{\text{pl}}^4 V_\phi V_{\phi\phi\phi}}{V^2}, \quad (20)$$

and

$$\varpi_V^3 = \frac{M_{\text{pl}}^6 V_\phi^2 V_{\phi\phi\phi\phi}}{V^3}. \quad (21)$$

In single field inflation with a standard kinetic term, as discussed here, the tensor spectrum shape is not independent from the other parameters. The slow-roll paradigm implies a tensor-to-scalar ratio, at the pivot scale, of

$$r = \frac{\mathcal{P}_t(k_*)}{\mathcal{P}_s(k_*)} \approx 16\epsilon \approx -8n_t, \quad (22)$$

referred to as the consistency relation. This consistency relation is also useful to understand how r is connected to the evolution of the inflaton:

$$\frac{\Delta\phi}{M_{\text{pl}}} \approx \frac{1}{\sqrt{8}} \int_0^N dN \sqrt{r}. \quad (23)$$

The above relation, called the Lyth bound ([Lyth, 1997](#)), implies that an inflaton variation of the order of the Planck mass is needed to produce $r \gtrsim 0.01$. Such a threshold is useful to classify large and small field inflationary models with respect to the Lyth bound.

2.3. Ending inflation and the epoch of entropy generation

The greatest uncertainty in calculating the perturbation spectrum predicted from a particular inflationary potential arises in establishing the correspondence between the comoving wavenumber today, and the inflaton energy density when the mode of that wavenumber crossed the Hubble radius during inflation ([Kinney & Riotto, 2006](#)). This correspondence depends both on the inflationary model and on the cosmological evolution from the end of inflation to the present.

After the slow-roll stage, $\dot{\phi}$ becomes as important as the *cosmological damping* term $3H\dot{\phi}$. Inflation ends gradually as the inflaton picks up kinetic energy so that w is no longer slightly above -1 , but rather far from that value. We may arbitrarily deem that inflation ends when $w = -1/3$ (the value dividing the cases of an expanding and a contracting comoving Hubble radius), or, equivalently, at $\epsilon_V \approx 1$, after which the epoch of entropy generation starts. Because of couplings to other fields, the energy initially in the form of scalar field vacuum energy is transferred to the other fields by perturbative decay (reheating), possibly preceded by a non-perturbative stage (preheating). There is considerable uncertainty about the mechanisms of entropy generation, or thermalization, which subsequently lead to a standard $w = 1/3$ equation of state for radiation.

On the other hand, if we want to identify some k_* today with the value of the inflaton field at the time this scale left the horizon, Eq. 12 needs to be matched to an expression that quantifies how much k_* has shrunk relative to the size of the horizon between the end of inflation and the time that mode re-enters the horizon. This quantity depends both on the inflationary potential and the details of the entropy generation process, and is given by

$$N_* \approx 71.21 - \ln\left(\frac{k_*}{a_0 H_0}\right) + \frac{1}{4} \ln\left(\frac{V_{\text{hor}}}{M_{\text{pl}}^4}\right) + \frac{1}{4} \ln\left(\frac{V_{\text{hor}}}{\rho_{\text{end}}}\right) + \frac{1 - 3w_{\text{int}}}{12(1 + w_{\text{int}})} \ln\left(\frac{\rho_{\text{th}}}{\rho_{\text{end}}}\right), \quad (24)$$

where ρ_{end} is the energy density at the end of inflation, ρ_{th} is an energy scale by which the Universe has thermalized, $a_0 H_0$ is the present horizon scale, V_{hor} is the potential energy when the present horizon scale left the horizon during inflation, and w_{int} characterizes the effective equation of state between the end of inflation and the energy scale specified by ρ_{th} . In predicting the primordial power spectra at observable scales for a specific inflaton potential, this uncertainty in the reheating history of the Universe becomes relevant and can be taken into account by allowing N_* to vary over a range of values. Note that w_{int} is not intended to provide a detailed model for entropy generation, but rather to parameterize the uncertainty regarding the expansion rate of the Universe during this intermediate era. Nevertheless, constraints on w_{int} provide observational limits on the uncertain physics during this period.

The first two terms of Eq. 24 are model independent, with the second term being roughly 5 for $k_* = 0.05 \text{ Mpc}^{-1}$. If thermalization occurs rapidly, or if the reheating stage is close to radiation-like, the magnitude of the last term in Eq. 24 is $\lesssim 1$. For most reasonable inflation models, the fourth term is $\mathcal{O}(1)$ and the third term ~ -10 , motivating the commonly assumed range $50 < N_* < 60$. Nonetheless, more extreme values on both ends are in principle possible ([Liddle & Leach, 2003](#)). In the figures of Sect. 4 we will mark the range $50 < N_* < 60$ to guide the reader's eye.

2.4. Perturbations from cosmic inflation at higher order

To calculate the quantum fluctuations generated during cosmic inflation, a linearized quantum field theory in a time-dependent background can be used. The leading order is the two-point correlation function

$$\langle \mathcal{R}(\mathbf{k}_1) \mathcal{R}(\mathbf{k}_2) \rangle = (2\pi)^3 \frac{2\pi^2}{k^3} \mathcal{P}_{\mathcal{R}}(k) \delta^3(\mathbf{k}_1 + \mathbf{k}_2), \quad (25)$$

but the inflaton self-interactions and the nonlinearity of Einstein gravity give small higher-order corrections, of which the next-to-leading order is the three point function

$$\langle \mathcal{R}(\mathbf{k}_1) \mathcal{R}(\mathbf{k}_2) \mathcal{R}(\mathbf{k}_3) \rangle = (2\pi)^3 B_{\mathcal{R}}(k_1, k_2, k_3) \delta^3(\mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3), \quad (26)$$

which is in general non-zero.

For single-field inflation with a standard kinetic term in a smooth potential (with initial fluctuations in the Bunch-Davies vacuum), the non-Gaussian contribution to the curvature perturbation during inflation is $\mathcal{O}(\epsilon_V, \eta_V)$ (Acquaviva et al., 2003; Maldacena, 2003), i.e., at an undetectable level smaller than other general relativistic contributions, such as the cross-correlation between the integrated Sachs-Wolfe effect and weak gravitational lensing of the CMB. For a general scalar field Lagrangian, the non-Gaussian contribution can be large enough to be accessible to *Planck* with $f_{\text{NL}} \sim c_s^{-2}$ (Chen et al., 2007), where c_s is the sound speed of inflaton fluctuations (see Sect. 9). Other higher order kinetic and spatial derivative terms contribute to larger non-Gaussianities. For a review of non-Gaussianity generated during inflation, see, for example, Bartolo et al. (2004a); Chen (2010), and the companion paper Planck Collaboration XXIV (2013).

2.5. Multi-field models of cosmic inflation

Inflation as described thus far assumes a single scalar field that drives and terminates the quasi-exponential expansion, and also generates the large-scale curvature perturbations. When there is more than one field with an effective mass smaller than H , isocurvature perturbations are also generated during inflation, by the same mechanism of amplification due to the stretching of the spacetime geometry (Axenides et al., 1983; Linde, 1985). Models with an N component inflaton ϕ_i can be best analysed in terms of an average field σ moving along the classical trajectory ($\dot{\sigma}^2 \equiv \sum_{i=1}^N \dot{\phi}_i^2$), whose fluctuations are related to the curvature perturbation according to

$$\mathcal{R} = -H \frac{\sum_{i=1}^N \dot{\phi}_i Q_i}{\dot{\sigma}^2}, \quad (27)$$

where Q_i is the gauge-invariant field fluctuation associated with ϕ_i . The $N - 1$ normal directions are connected to $N - 1$ isocurvature perturbations δs_{ij} according to

$$\delta s_{ij} = \frac{\dot{\phi}_i Q_j - \dot{\phi}_j Q_i}{\dot{\sigma}}. \quad (28)$$

If the trajectory of the average field is curved in field space, then during inflation both curvature and isocurvature fluctuations are generated with non-vanishing correlations.

Isocurvature perturbations can be converted into curvature perturbations on large scales, but the opposite does not hold (Mollerach, 1990). If such isocurvature perturbations are not totally converted into curvature perturbations and persist after nucleosynthesis, these can lead to observable effects in CMB

anisotropies and in structure formation. In Sect. 10, we present the *Planck* constraints on a combination of curvature and isocurvature initial conditions, and the implications for important two-field scenarios, such as the curvaton (Lyth & Wands, 2002) and axion (Lyth, 1990) models.

Isocurvature perturbations may lead to a higher level of non-Gaussianity compared to a single inflaton with a standard kinetic term (Groot Nibbelink & van Tent, 2000). There is no reason to expect the inflaton to be a single-component field: the scalar sector of the Standard Model and its extensions contain numerous scalar fields.

3. Methodology

3.1. Cosmological model and parameters

The parameters of the models to be estimated in this paper fall into three categories: (i) parameters describing the initial perturbations, i.e., characterizing the particular inflationary scenario in question; (ii) parameters determining cosmological evolution at late times ($z \lesssim 10^4$); and (iii) parameters that quantify our uncertainty about the instrument and foreground contributions to the angular power spectrum. These will be described in Section 3.2.1.

Unless specified otherwise, we assume that the late-time cosmology is the standard flat six parameter Λ CDM model, whose energy content consists of photons, baryons, cold dark matter, neutrinos (assuming $N_{\text{eff}} = 3.046$ effective species, one of which is taken to be massive with a mass of $m_\nu = 0.06$ eV), and a cosmological constant. The primordial helium fraction, Y_p , is set as a function of $\Omega_b h^2$ and N_{eff} according to the Big Bang nucleosynthesis consistency condition (Ichikawa & Takahashi, 2006; Hamann et al., 2008b), and we fix the CMB mean temperature to $T_0 = 2.7255$ K (Fixsen, 2009). Reionization is assumed to occur rapidly at a redshift z_{re} , with optical depth τ . This model can be characterized by four free cosmological parameters: $\Omega_b h^2, \Omega_c h^2, \theta_{\text{MC}}, \tau$, defined in Table 1, in addition to the parameters describing the initial perturbations.

3.2. Data

The CMB data we use consists of the *Planck* CMB temperature likelihood supplemented by the *WMAP* large-scale polarization likelihood (henceforth *Planck+WP*), as described in Sect. 3.2.1. The large scale angular E -mode polarization spectrum is important for constraining reionization. This breaks the degeneracy in the temperature data between the primordial power spectrum amplitude and the optical depth to reionization. In some cases, we also include the *Planck* lensing likelihood, measurements of the CMB at higher resolution from the Atacama Cosmology Telescope (ACT) and the South Pole Telescope (SPT), and measurements of the baryon acoustic oscillation (BAO) scale. This follows the same approach as Planck Collaboration XVI (2013).

3.2.1. *Planck* CMB temperature data

The *Planck* CMB likelihood is based on a hybrid approach, which combines a Gaussian likelihood approximation based on temperature pseudo cross-spectra at high multipoles (Hamimeche & Lewis, 2008), with a pixel-based temperature and polarization likelihood at low multipoles. We summarize the likelihood here, and for a detailed description refer the reader to Planck Collaboration XV (2013).

The small-scale *Planck* temperature likelihood is based on pseudo cross-spectra between pairs of maps at 100, 143, and 217 GHz, masked to retain 49%, 31%, and 31% of the sky, respectively. This results in angular auto- and cross-correlation power spectra covering multipole ranges of $50 \leq \ell \leq 1200$ at 100 GHz, $50 \leq \ell \leq 2000$ at 143 GHz, and $500 \leq \ell \leq 2500$ at 217 GHz and for the 143×217 GHz cross-spectrum. In addition to instrumental uncertainties, mitigated here by using only cross-spectra among different detectors, small scale foreground and CMB secondary anisotropies need to be accounted for. The foreground model used in the *Planck* high- ℓ likelihood is described in detail in Planck Collaboration XV (2013); Planck Collaboration XVI (2013), and includes contributions to the cross-frequency power spectra from unresolved radio point sources, the cosmic infrared background (CIB), and the thermal and kinetic Sunyaev-Zeldovich effects. There are eleven adjustable *nuisance* parameters: $(A_{100}^{\text{PS}}, A_{143}^{\text{PS}}, A_{217}^{\text{PS}}, r_{143 \times 217}^{\text{PS}}, A_{143}^{\text{CIB}}, A_{217}^{\text{CIB}}, r_{143 \times 217}^{\text{CIB}}, \gamma^{\text{CIB}}, A_{143}^{\text{SZ}}, A^{\text{kSZ}}, \xi^{\text{SZ-CIB}})$. In addition, the calibration parameters for the 100 and 217 GHz channels, c_{100} and c_{217} , relative to the 143 GHz channel, and the dominant beam uncertainty eigenmode amplitude B_1^1 are left free in the analysis, with other beam uncertainties marginalized analytically, so the *Planck* high- ℓ likelihood includes 14 nuisance parameters.

The low- ℓ *Planck* likelihood combines the *Planck* temperature data with the large scale 9-year *WMAP* polarization data for this release. Using the procedure introduced in Page et al. (2007), the temperature and polarization likelihood can be separated assuming negligible noise in the temperature map. The temperature likelihood is based on a Gibbs approach (Eriksen et al., 2007), mapping out the distribution of the $\ell < 50$ CMB temperature multipoles from a foreground-cleaned combination of the 30 – 353 GHz maps. The polarization likelihood uses a pixel-based approach using the *WMAP* 9-year polarization maps at 33, 41, and 61 GHz, and includes the temperature-polarization cross-correlation (Page et al., 2007). Its angular range is $\ell \leq 23$ for *TE*, *EE*, and *BB*.

3.2.2. *Planck* lensing data

The primary CMB anisotropies are distorted by the gravitational potential induced by intervening matter. Such lensing, which tends to broaden and smooth out the acoustic oscillations, is taken into account as a correction to the observed temperature power spectrum. The lensing power spectrum can also be recovered by measuring higher-order correlation functions.

For some of our analysis we include the *Planck* lensing likelihood, derived in Planck Collaboration XVII (2013), which measures the non-Gaussian trispectrum of the CMB, which is proportional to the power spectrum of the lensing potential. As described in Planck Collaboration XVII (2013), this potential is reconstructed using quadratic estimators (Okamoto & Hu, 2003), and its power spectrum is used to estimate the lensing deflection power spectrum. The spectrum is estimated from the 143 and 217 GHz maps, using multipoles in the range $40 < \ell < 400$. The theoretical predictions for the lensing potential power spectrum are calculated at linear order.

3.2.3. ACT and SPT temperature data

In some of our analysis we include data from ACT and SPT. ACT measures the power spectrum at 148 and 218 GHz, and the cross-spectrum (Das et al., 2013), and covers angular scales

$500 < \ell < 10000$ at 148 GHz and $1500 < \ell < 10000$ at 218 GHz. We use these data in the range $\ell > 1000$ in combination with *Planck*. SPT measures the power spectrum for angular scales $2000 < \ell < 10000$ at 95, 150, and 220 GHz (Reichardt et al., 2012). The spectrum at larger scales is also measured at 150 GHz in Story et al. (2012) but we do not include these in our analysis. To model the foregrounds for ACT and SPT we follow a similar approach to the likelihood described in Dunkley et al. (2013), extending the model used for the *Planck* high- ℓ likelihood. Additional nuisance parameters are included to model the Poisson source levels in each experiment, the residual Galactic dust, and inter-frequency calibration parameters. They are described in more detail in Planck Collaboration XVI (2013).

3.2.4. BAO data

The BAO angular scale acts as a *standard ruler* and allows us to map out the expansion history of the Universe after last scattering. The BAO scale, extracted from galaxy redshift surveys, provides a constraint on the late-time geometry and breaks degeneracies with other cosmological parameters. Galaxy surveys constrain the ratio $D_V(\bar{z})/r_s$, where $D_V(\bar{z})$ is the spherically averaged distance scale to the effective survey redshift \bar{z} and r_s is the sound horizon (Mehta et al., 2012).

In this analysis we consider a combination of the measurements by the 6dFGRS (Beutler et al. (2011), $\bar{z} = 0.106$), SDSS-II (Padmanabhan et al. 2012, $\bar{z} = 0.35$), and BOSS CMASS (Anderson et al. 2012, $\bar{z} = 0.57$) surveys, assuming no correlation between the three data points. This likelihood is described further in Planck Collaboration XVI (2013).

3.3. Parameter estimation

Given a model \mathcal{M} with free parameters $\hat{x} \equiv \{x_1, \dots, x_X\}$ and a likelihood function of the data $\mathcal{L}(\text{data}|\hat{x})$, the (posterior) probability density \mathcal{P} as a function of the parameters can be expressed as

$$\mathcal{P}(\hat{x}|\text{data}, \mathcal{M}) \propto \mathcal{L}(\text{data}|\hat{x}) \cdot P(\hat{x}|\mathcal{M}), \quad (29)$$

where $P(\hat{x}|\mathcal{M})$ represents the data-independent prior probability density. Unless specified otherwise, we choose wide top-hat prior distributions for all cosmological parameters.

We construct the posterior parameter probabilities using the Markov Chain Monte Carlo (MCMC) sampler as implemented in the CosmoMC (Lewis & Bridle, 2002) or MontePython (Audren et al., 2012) packages. In some cases, when the calculation of the Bayesian evidence (see below) is desired, or when the likelihood function deviates strongly from a multivariate Gaussian, we use the nested sampling algorithm provided by the MultiNest add-on module (Feroz & Hobson, 2008; Feroz et al., 2009) instead of the Metropolis-Hastings algorithm.

Joint two-dimensional and one-dimensional posterior distributions are obtained by marginalization. Numerical values and constraints on parameters are quoted in terms of the mean and 68% central Bayesian interval of the respective one-dimensional marginalized posterior distribution.

3.4. Model selection

Two approaches to model selection are commonly used in statistics. The first approach examines the logarithm of the likelihood ratio, or *effective* χ^2 ,

$$\Delta\chi_{\text{eff}}^2 \equiv 2 [\ln \mathcal{L}_{\text{max}}(\mathcal{M}_1) - \ln \mathcal{L}_{\text{max}}(\mathcal{M}_2)], \quad (30)$$

between models \mathcal{M}_1 and \mathcal{M}_2 , corrected for the fact that models with more parameters provide a better fit due to fitting away noise, even when the more complicated model is not correct. Various information criteria have been proposed based on this idea (Akaike, 1974; Schwarz, 1978), see also Liddle (2007). These quantities have the advantage of being independent of prior choice and fairly easy to calculate. The second approach is a Bayesian approach (Cox, 1946; Jeffreys, 1998; Jaynes & Bretthorst, 2003), based on evaluating ratios of the *model averaged likelihood*, or Bayesian evidence, defined by

$$\mathcal{E}_i = \int d\mathbf{x}^X P(\hat{\mathbf{x}}|\mathcal{M}_i) \mathcal{L}(\text{data}|\hat{\mathbf{x}}). \quad (31)$$

Evidence ratios are naturally interpreted as *betting odds* between models.⁵ Nested sampling algorithms allow rapid numerical evaluation of \mathcal{E} . In this paper we will consider both the effective χ^2 and the Bayesian evidence.

4. Constraints on slow-roll inflationary models

In this section we describe constraints on the slow-roll inflationary paradigm, using *Planck*+WP data in combination with the likelihoods described in Sects. 3.2.2–3.2.4. First we concentrate on characterizing the primordial power spectrum using *Planck* and other data. We start by showing that the empirical pre-inflationary Harrison-Zeldovich (HZ) spectrum with $n_s = 1$ does not fit the *Planck* measurements. We further examine whether generalizing the cosmological model, for example by allowing the number of neutrino species to vary, allowing the Helium fraction to vary, or admitting a non-standard reionization scenario could reconcile the data with $n_s = 1$, and conclude that $n_s \neq 1$ is robust. Next we consider evidence for a running of n_s and show it constrained to be small, although we find a preference for negative running at modest statistical significance.

We then investigate the *Planck* constraints on slow roll inflation, allowing a tilt for the spectral index and the presence of tensor modes, and discuss the implications for the simplest standard inflationary models. In this section the question is studied using the slow roll approximation, but later sections move beyond the slow roll approximation. We show that compared to previous experiments, *Planck* significantly narrows the space of possible inflationary models. Finally we comment on the implications for inflation of the *Planck* constraints on possible deviations from spatial flatness.

4.1. Ruling out exact scale invariance

The simplest ansatz for characterizing the statistical properties of the primordial cosmological perturbations is the so-called HZ model proposed by Harrison (1970); Zeldovich (1972); and Peebles & Yu (1970). These authors pointed out that a power spectrum with exact scale invariance for the Newtonian gravitation potential fitted the data available at the time, but without giving any theoretical justification for this form of the spectrum. Under exact scale invariance, which would constitute a sort of unexplained new symmetry, the primordial perturbations in the

Newtonian gravitational potential look exactly the same whether they are magnified or demagnified. In this simple model, vector and tensor perturbations are absent and the spectrum of curvature perturbations is characterized by a single parameter, the amplitude A_s . Inflation, on the other hand, generically breaks this rescaling symmetry. Although under inflation, scale invariance still holds approximately, inflation must end: therefore, as different scales are being imprinted, the physical conditions must also evolve. (However, see Starobinsky (2005) for the canonical inflationary model designed to mimic $n_s = 1$, albeit with $r \neq 0$).

Although a detection of a violation of scale invariance would not definitively prove that inflation is responsible for the generation of the primordial perturbations, ruling out the HZ model would confirm the expectation of small deviations from scale invariance, almost always on the red side, which are generic to all inflationary models without fine tuning. We examine in detail the viability of the HZ model using statistics to compare it to the more general model where the spectral index is allowed to vary, as motivated by slow-roll inflation.

When the cosmological model with $n_s = 1$ is compared with a model in which n_s is allowed to vary, we find that allowing n_s to deviate from one decreases the best-fit effective χ^2 by 27.9 with respect to the HZ model. Thus the significance of the finding that $n_s \neq 1$ is in excess of 5σ . The parameters and maximum likelihood of this comparison are reported in Table 3.

One might wonder whether $n_s = 1$ could be reconciled with the data by relaxing some of the assumptions of the underlying cosmological model. Of particular interest is exploring those parameters known to be almost degenerate with the spectral index, such as the effective number of neutrino species N_{eff} and the primordial helium fraction Y_p , which both alter the damping tail of the temperature spectrum (Trotta & Hansen, 2004; Hou et al., 2011), somewhat mimicking a spectral tilt. Assuming a Harrison-Zeldovich spectrum and allowing N_{eff} or Y_p to float, and thus deviate from their standard values, gives almost as good a fit to *Planck*+WP data as the Λ CDM model with a varying spectral index, with $\Delta\chi_{\text{eff}}^2 = 2.8$ and 2.2, respectively. However, as shown in Table 3, the HZ, HZ+ N_{eff} , and HZ+ Y_p models require significantly higher baryon densities and reionization optical depths compared to Λ CDM. In the HZ+ Y_p model, one obtains a helium fraction of $Y_p = 0.3194 \pm 0.013$. This value is incompatible both with direct astrophysical measurements of this quantity (Aver et al., 2012) and with standard Big Bang nucleosynthesis (Hamann et al., 2008b). The HZ+ N_{eff} model, on the other hand, would imply the presence of $\Delta N_{\text{eff}} \approx 1$ new effective neutrino species beyond the three known species. When BAO measurements are included in the likelihood, $\Delta\chi_{\text{eff}}^2$ increases to 39.2 (HZ), 4.6 (HZ+ Y_p), and 8.0 (HZ+ N_{eff}), respectively, for the three models. The significance of this detection is also discussed in Planck Collaboration XVI (2013).

4.2. Constraining inflationary models using the slow roll approximation

We now consider all inflationary models that can be described by the primordial power spectrum parameters consisting of the scalar amplitude, A_s , the spectral index, n_s , and the tensor-to-scalar ratio r , all defined at the pivot scale k_* . We assume that the spectral index is independent of the wavenumber k . Negligible running of the spectral index is expected if the slow roll condition is satisfied and higher order corrections in the slow roll approximations can be neglected. In the next subsection we relax this assumption.

⁵ Note that since the average is performed over the entire support of the prior probability density, the evidence depends strongly on the probability range for the adjustable parameters. Whereas in parameter inference, the exact extent of the prior ranges often becomes irrelevant as long as they are “wide enough” (i.e., containing the bulk of the high-likelihood region in parameter space), the value of the evidence will generally depend on precisely how wide the prior range was chosen.

	HZ	HZ + Y_p	HZ + N_{eff}	ΛCDM
$10^5 \Omega_b h^2$	2296 ± 24	2296 ± 23	2285 ± 23	2205 ± 28
$10^4 \Omega_c h^2$	1088 ± 13	1158 ± 20	1298 ± 43	1199 ± 27
$100 \theta_{\text{MC}}$	1.04292 ± 0.00054	1.04439 ± 0.00063	1.04052 ± 0.00067	1.04131 ± 0.00063
τ	$0.125^{+0.016}_{-0.014}$	$0.109^{+0.013}_{-0.014}$	$0.105^{+0.014}_{-0.013}$	$0.089^{+0.012}_{-0.014}$
$\ln(10^{10} A_s)$	$3.133^{+0.032}_{-0.028}$	$3.137^{+0.027}_{-0.028}$	$3.143^{+0.027}_{-0.026}$	$3.089^{+0.024}_{-0.027}$
n_s	—	—	—	0.9603 ± 0.0073
N_{eff}	—	—	3.98 ± 0.19	—
Y_p	—	0.3194 ± 0.013	—	—
$-2\Delta \ln(\mathcal{L}_{\text{max}})$	27.9	2.2	2.8	0

Table 3. Constraints on cosmological parameters and best-fit $-2\Delta \ln(\mathcal{L})$ with respect to the standard ΛCDM model, using *Planck*+WP data, testing the significance of the deviation from the HZ model.

Sampling the power spectrum parameters A_s , n_s , and r is not the only method for constraining slow roll inflation. Another possibility is to sample the HFF in the analytic expressions for the scalar and tensor power spectra (Stewart & Lyth, 1993; Gong & Stewart, 2001; Leach et al., 2002). In the Appendix, we perform a comparison of slow-roll inflationary predictions by sampling the HFF with *Planck* data, and show that the results obtained in this way agree with those derived by sampling the power spectrum parameters. This confirms similar studies with previous data (Hamann et al., 2008c; Finelli et al., 2010).

The spectral index estimated from *Planck*+WP data is

$$n_s = 0.9603 \pm 0.0073. \quad (32)$$

This tight bound on n_s is crucial for constraining inflation. The *Planck* constraint on r depends slightly on the pivot scale; we adopt $k_* = 0.002 \text{ Mpc}^{-1}$ to quote our result, with $r_{0.002} < 0.12$ at 95% CL. This bound improves on the most recent results, including the WMAP 9-year constraint of $r < 0.38$ (Hinshaw et al., 2012a), the WMAP7 + ACT limit of $r < 0.28$ (Sievers et al., 2013), and the WMAP7 + SPT limit of $r < 0.18$ (Story et al., 2012). The new bound from *Planck* is consistent with the limit from temperature anisotropies alone (Knox & Turner, 1994). When a possible tensor component is included, the spectral index from *Planck*+WP is not significantly changed, with $n_s = 0.9624 \pm 0.0075$.

The *Planck* constraint on r corresponds to an upper bound on the energy scale of inflation

$$V_* = \frac{3\pi^2 A_s}{2} r M_{\text{pl}}^4 = (1.94 \times 10^{16} \text{ GeV})^4 \frac{r_*}{0.12}, \quad (33)$$

at 95% CL. This is equivalent to an upper bound on the Hubble parameter during inflation of $H_*/M_{\text{pl}} < 3.7 \times 10^{-5}$. In terms of slow-roll parameters, *Planck*+WP constraints imply $\epsilon_V < 0.008$ at 95% CL, and $\eta_V = -0.010^{+0.005}_{-0.011}$.

The *Planck* results on n_s and r are robust to the addition of external data sets (see Table 4). When the high- ℓ CMB ACT+SPT data are added, we obtain $n_s = 0.9600 \pm 0.0072$ and $r_{0.002} < 0.11$ at 95% CL. Including the *Planck* lensing likelihood gives $n_s = 0.9653 \pm 0.0069$ and $r_{0.002} < 0.13$, and adding BAO data gives $n_s = 0.9643 \pm 0.0059$ and $r_{0.002} < 0.12$. These bounds are robust to the small changes in the polarization likelihood at low multipoles. To test this robustness, instead of using the WMAP polarization likelihood, we impose a Gaussian prior $\tau = 0.07 \pm 0.013$ to take into account small shifts due to uncertainties in residual foreground contamination or instrument systematics in the evaluation of τ , as performed in Appendix B of Planck Collaboration XVI (2013). We find at most a reduction of 8% for the upper bound on r .

It is useful to plot the inflationary potentials in the n_s – r plane using the first two slow-roll parameters evaluated at the pivot scale $k_* = 0.002 \text{ Mpc}^{-1}$ (Dodelson et al., 1997). Given our ignorance of the details of the epoch of entropy generation, we assume that the number of e -folds N_* to the end of inflation lies in the interval [50, 60]. This uncertainty is plotted for those potentials predicting an exit from inflation without changing the potential.

Fig. 1 shows the *Planck* constraints in the n_s – r plane and indicates the predictions of a number of representative inflationary potentials. The sensitivity of *Planck* data to high multipoles removes the degeneracy between n_s and r found using WMAP data. *Planck* data favour models with a concave potential. As shown in Fig. 1, most of the joint 95% allowed region lies below the convex potential limit, and concave models with a red tilt in the range [0.945–0.98] are allowed by *Planck* at 95% CL. In the following we consider the status of several illustrative and commonly discussed inflationary potentials in light of the *Planck* observations.

Power law potential and chaotic inflation

The simplest class of inflationary models is characterized by a single monomial potential of the form

$$V(\phi) = \lambda M_{\text{pl}}^4 \left(\frac{\phi}{M_{\text{pl}}} \right)^n. \quad (34)$$

This class of potentials includes the simplest chaotic models, in which inflation starts from large values for the inflaton, $\phi > M_{\text{pl}}$. Inflation ends by violation of the slow-roll regime, and we assume this occurs at $\epsilon_V = 1$. According to Eqs. 5, 6, and 15, this class of potentials predicts to lowest order in slow-roll parameters $n_s - 1 \approx -n(n+2)M_{\text{pl}}^2/\phi_*^2$, $r \approx 8n^2 M_{\text{pl}}^2/\phi_*^2$, $\phi_*^2 \approx nM_{\text{pl}}^2(4N_* + n)/2$. The $\lambda\phi^4$ model lies well outside of the joint 99.7% CL region in the n_s – r plane. This result confirms previous findings from e.g., Hinshaw et al. (2012a) in which this model is well outside the 95% CL for the WMAP 9-year data and is further excluded by CMB data at smaller scales.

The model with a quadratic potential, $n = 2$ (Linde, 1983), often considered the simplest example for inflation, now lies outside the joint 95% CL for the *Planck*+WP+high- ℓ data for $N_* \lesssim 60$ e -folds, as shown in Fig. 1.

A linear potential with $n = 1$ (McAllister et al., 2010), motivated by axion monodromy, has $\eta_V = 0$ and lies within the 95% CL region. Inflation with $n = 2/3$ (Silverstein & Westphal, 2008), however, also motivated by axion monodromy, now lies on the boundary of the joint 95% CL region. More permissive

Model	Parameter	<i>Planck</i> +WP	<i>Planck</i> +WP+lensing	<i>Planck</i> + WP+high- ℓ	<i>Planck</i> +WP+BAO
Λ CDM + tensor	n_s	0.9624 ± 0.0075	0.9653 ± 0.0069	0.9600 ± 0.0071	0.9643 ± 0.0059
	$r_{0.002}$	< 0.12	< 0.13	< 0.11	< 0.12
	$-2\Delta \ln \mathcal{L}_{\max}$	0	0	0	-0.31

Table 4. Constraints on the primordial perturbation parameters in the Λ CDM+ r model from *Planck* combined with other data sets. The constraints are given at the pivot scale $k_* = 0.002 \text{ Mpc}^{-1}$.




Fig. 1. Marginalized joint 68% and 95% CL regions for n_s and $r_{0.002}$ from *Planck* in combination with other data sets compared to the theoretical predictions of selected inflationary models.

reheating priors allowing $N_* < 50$ could reconcile this model with the *Planck* data.

Exponential potential and power law inflation

Inflation with an exponential potential

$$V(\phi) = \Lambda^4 \exp\left(-\lambda \frac{\phi}{M_{\text{pl}}}\right) \quad (35)$$

is called power law inflation (Lucchin & Matarrese, 1985), because the exact solution for the scale factor is given by $a(t) \propto t^{2/\lambda^2}$. This model is incomplete, since inflation would not end without an additional mechanism to stop it. Assuming such a mechanism exists and leaves predictions for cosmological perturbations unmodified, this class of models predicts $r = -8(n_s - 1)$ and is now outside the joint 99.7% CL contour.

Inverse power law potential

Intermediate models (Barrow, 1990; Muslimov, 1990) with inverse power law potentials

$$V(\phi) = \Lambda^4 \left(\frac{\phi}{M_{\text{pl}}}\right)^{-\beta} \quad (36)$$

lead to inflation with $a(t) \propto \exp(At^f)$, with $A > 0$ and $0 < f < 1$, where $f = 4/(4 + \beta)$ and $\beta > 0$. In intermediate inflation there is no natural end to inflation, but if the exit mechanism leaves the inflationary predictions on cosmological perturbations unmodified, this class of models predicts $r \approx -8\beta(n_s - 1)/(\beta - 2)$ (Barrow & Liddle, 1993). It is disfavoured, being outside the joint 95% CL contour for any β .

Hill-top models

In another interesting class of potentials, the inflaton rolls away from an unstable equilibrium as in the first new inflationary models (Albrecht & Steinhardt, 1982; Linde, 1982). We consider

$$V(\phi) \approx \Lambda^4 \left(1 - \frac{\phi^p}{\mu^p} + \dots\right), \quad (37)$$

where the ellipsis indicates higher order terms negligible during inflation, but needed to ensure the positiveness of the potential later on. An exponent of $p = 2$ is allowed only as a large field inflationary model and predicts $n_s - 1 \approx -4M_{\text{pl}}^2/\mu^2 + 3r/8$ and $r \approx 32\phi_*^2 M_{\text{pl}}^2/\mu^4$. This potential leads to predictions in agreement with *Planck*+WP+BAO joint 95% CL contours for super-Planckian values of μ , i.e., $\mu \gtrsim 9 M_{\text{pl}}$.

Models with $p \geq 3$ predict $n_s - 1 \approx -(2/N)(p - 1)/(p - 2)$ when $r \sim 0$. The hill-top potential with $p = 3$ lies outside the

joint 95% CL region for *Planck*+WP+BAO data; the case with $p = 4$ is also in tension with *Planck*+WP+BAO, but allowed within the joint 95% CL region for $N_* \gtrsim 50$. For larger values of r these models provide a better fit to the *Planck*+WP+BAO data.

A simple symmetry breaking potential

The symmetry-breaking potential (Olive, 1990)

$$V(\phi) = \Lambda^4 \left(1 - \frac{\phi^2}{\mu^2} \right)^2, \quad (38)$$

can be considered as a self-consistent completion of the hill-top model with $p = 2$ (although it has a different limiting large-field branch for non-zero r). This potential leads to predictions in agreement with *Planck* + WP + BAO joint 95% CL contours for super-Planckian value of μ , i.e. $\mu \gtrsim 13 M_{\text{pl}}$.

Natural inflation

Another interesting class of potentials is *natural* inflation (Freese et al., 1990; Adams et al., 1993), initially motivated by its origin in symmetry-breaking in an attempt to naturally give rise to the extremely flat potentials required for inflationary cosmology. In *natural inflation* the effective one-dimensional potential takes the form

$$V(\phi) = \Lambda^4 \left[1 + \cos \left(\frac{\phi}{f} \right) \right], \quad (39)$$

where f is a scale which determines the slope of the potential. Depending on the value of f , the model falls into the large field ($f \gtrsim 1.5 M_{\text{pl}}$) or small field ($f \lesssim 1.5 M_{\text{pl}}$) classification scheme. Therefore, $n_s \approx 1 - M_{\text{pl}}^2/f^2$ holds for small f and $n_s \approx 1 - 2/N$, $r \approx 8/N$ holds for large f , approximating the $m^2\phi^2$ potential in the latter case (with $N_* \approx (2f^2/M_{\text{pl}}^2) \ln[\sin(\phi_e/f)/\sin(\phi_*/f)]$). This model agrees with *Planck*+WP data for $f \gtrsim 5 M_{\text{pl}}$.

Hybrid inflation

In hybrid inflationary models a second field, χ , coupled to the inflaton, undergoes symmetry breaking. The simplest example of this class is

$$V(\phi, \chi) = \Lambda^4 \left(1 - \frac{\chi^2}{\mu^2} \right)^2 + U(\phi) + \frac{g^2}{2} \phi^2 \chi^2. \quad (40)$$

For most of their parameter space, these models can be considered effectively as single field models for the inflaton ϕ . The second field χ is close to the origin during the slow-roll regime for ϕ , and inflation ends either by breakdown of slow roll for the inflaton at $\epsilon_\phi \approx M_{\text{pl}}^2 (dU/d\phi)^2 / (\Lambda^4 + U(\phi))^2 \approx 1$ or by the waterfall transition of χ . The simplest models with

$$U(\phi) = \frac{m^2}{2} \phi^2 \quad (41)$$

are disfavoured for most of the parameter space (Cort s & Liddle, 2009). Models with $m^2\phi^2/2 \sim \Lambda^4$ are disfavoured due to a high tensor-to-scalar ratio, and models with $U(\phi) \ll \Lambda^4$ predict a spectral index $n_s > 1$, also disfavoured by the *Planck* data.

We discuss hybrid inflationary models predicting $n_s < 1$ separately. As an example, the spontaneously broken SUSY model (Dvali et al., 1994)

$$U(\phi) = \alpha_h \Lambda^4 \ln \left(\frac{\phi}{\mu} \right), \quad (42)$$

predicts $n_s - 1 \approx -(1 + 3\alpha_h/2)/N_*$ and $r \approx 8\alpha_h/N_*$. For $\alpha_h \ll 1$ and $N_* \approx 50$, $n_s \approx 0.98$ is disfavoured by *Planck*+WP+BAO data at more than 95% CL. However, more permissive reheating priors allowing $N_* < 50$ or a non-negligible α_h give models that are consistent with the *Planck* data.

R^2 inflation

Inflationary models can also be accommodated within extended theories of gravity. These theories can be analysed either in the original (Jordan) frame or in the conformally-related Einstein frame with a Klein-Gordon scalar field. Due to the invariance of curvature and tensor perturbation power spectra with respect to this conformal transformation, we can use the same methodology described earlier.

The first inflationary model proposed was of this type and based on higher order gravitational terms in the action (Starobinsky, 1980)

$$S = \int d^4x \sqrt{-g} \frac{M_{\text{pl}}^2}{2} \left(R + \frac{R^2}{6M^2} \right), \quad (43)$$

with the motivation to include semi-classical quantum effects. The predictions for R^2 inflation were first studied in Mukhanov & Chibisov (1981); Starobinsky (1983) and can be summarized as: $n_s - 1 \approx -8(4N_* + 9)/(4N_* + 3)$ and $r \approx 192/(4N_* + 3)^2$. Since r is suppressed by another $1/N_*$ with respect to the scalar tilt, this model predicts a tiny amount of gravitational waves. This model predicts $n_s = 0.963$ for $N_* = 55$ and is fully consistent with the *Planck* constraints.

Non-minimally coupled inflaton

A non-minimal coupling of the inflaton to gravity with the action

$$S = \int d^4x \sqrt{-g} \left[\frac{M_{\text{pl}}^2 + \xi \phi^2}{2} R - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{\lambda}{4} (\phi^2 - \phi_0^2)^2 \right], \quad (44)$$

leads to several interesting consequences such as a lowering of the tensor-to-scalar ratio.

The case of a massless self-interacting inflaton ($\phi_0 = 0$) agrees with the *Planck*+WP data for $\xi \neq 0$. Within the range $50 < N_* < 60$, this model is within the *Planck*+WP joint 95% CL region for $\xi > 0.0019$, improving previous bounds (Okada et al., 2010).

The amplitude of scalar perturbations is proportional to λ/ξ^2 for $\xi \gg 1$, and therefore the problem of tiny values for the inflaton self-coupling λ can be alleviated (Salopek et al., 1989; Fakir & Unruh, 1990). The regime $\phi_0 \ll M_{\text{pl}}$ is allowed and ϕ could be the Standard Model Higgs as proposed in Bezrukov & Shaposhnikov (2009). The Higgs case with $\xi \gg 1$ has the same predictions as the R^2 model in terms of n_s and r as a function of N_* . The reheating mechanism in the Higgs case can be more efficient than in R^2 case and therefore predicts a slightly larger n_s . This model is fully consistent with *Planck* constraints.

The case with $\xi < 0$ and $|\xi|\phi_0^2/M_{\text{pl}}^2 \sim 1$ was also recently emphasized in Linde et al. (2011). With the symmetry breaking potential in Eq. 44, the large field case with inflaton $\phi > \phi_0$ is disfavoured by *Planck* data, whereas the small field case $\phi < \phi_0$ is in agreement with the data.

4.3. Running spectral index

We have shown that the single parameter Harrison-Zeldovich spectrum does not fit the data, and that at least the first two terms, A_s and n_s , in the expansion of the primordial power spectrum in powers of $\ln(k)$ given in Eq. 10 are needed. Here we consider whether the data require the next term known as the *running of the spectral index* (Kosowsky & Turner, 1995), defined as the derivative of the spectral index with respect to $\ln k$, $dn_s/d \ln k$ for scalar or tensor fluctuations. In slow-roll single-field inflation, the running is second order in the Hubble slow-roll parameters, for scalar and for tensor perturbations (see Eqs. 17 and 18, respectively) (Leach et al., 2002), and thus is typically suppressed with respect to, e.g., $n_s - 1$ and n_t , which are first order. If the slow-roll approximation holds and the inflaton has reached its attractor solution, $dn_s/d \ln k$ and $dn_t/d \ln k$ are related to the potential slow-roll parameters as in Eqs. 17 and 18. Given the tight constraints on the first two slow-roll parameters ϵ_V and η_V (ϵ_1 and ϵ_2) from the present data, typical values of the running to which *Planck* is sensitive (Pahud et al., 2007) would generically be dominated by the contribution from the third derivative of the potential, encoded in $|\xi_V^2|$ (or ϵ_3).

While it is easy to see that the running is invariant under a change in pivot scale, the same does not hold true for the spectral index and the amplitude of the primordial power spectrum. It is convenient to choose k_* such that $dn_s/d \ln k$ and n_s are decorrelated (Cort  s et al., 2007). This approach minimizes the inferred variance of n_s and facilitates comparison with constraints on n_s in the power law models. Note however that the decorrelation pivot scale k_*^{dec} depends both on the model and data set considered in the analysis.

We consider a model parameterizing the power spectrum using $A_s(k_*)$, $n_s(k_*)$, and $dn_s/d \ln k$, where $k_* = 0.05 \text{ Mpc}^{-1}$. The joint constraints on n_s and $dn_s/d \ln k$ at the decorrelation scale of $k_*^{\text{dec}} = 0.04 \text{ Mpc}^{-1}$ are shown in Fig. 4. The *Planck*+WP constraints on the running do not change significantly when complementary data sets such as *Planck* lensing, CMB high- ℓ , and BAO data are included. We find

$$dn_s/d \ln k = -0.013 \pm 0.009 \quad (68\% \text{ CL, Planck+WP}), \quad (45)$$

which is negative at the 1.5σ level. This reduces the the uncertainty compared to previous CMB results. Error bars are reduced by 60% compared to the *WMAP* 9-year results (Hinshaw et al., 2012a), and by 20-30% compared to *WMAP* supplemented by SPT and ACT data (Hou et al., 2012; Sievers et al., 2013). *Planck* finds a smaller scalar running than SPT+WMAP7 (Hou et al., 2012), and larger than ACT+WMAP7 (Sievers et al., 2013). The best-fit likelihood improves by only $\Delta\chi_{\text{eff}}^2 \approx 1.5$ (3 when high- ℓ data are included) with respect to the minimal case in which n_s is scale independent, indicating that the deviation from scale independence is not very significant. The constraint for the spectral index in this case is 0.9630 ± 0.0065 at 68% CL at the decorrelation pivot scale $k_* = 0.038 \text{ Mpc}^{-1}$. This result implies that the third derivative of the potential is small, i.e., $|\xi_V^2| \sim 0.007$, but compatible with zero at 95% CL, for inflation at low energy (i.e., with $\epsilon_V \approx 0$).

We also test the possibility that the running depends on the wavelength so that $d^2n_s/d \ln k^2$ is nonzero. With *Planck*+WP data, we find $d^2n_s/d \ln k^2 = 0.020^{+0.016}_{-0.015}$. This result is stable with respect to the addition of complementary data sets. When $d^2n_s/d \ln k^2$ is allowed in the fit, we find a value for the running $dn_s/d \ln k$ consistent with zero.

Finally we allow a non-zero primordial gravitational wave spectrum together with the running. The tensor spectral in-




Fig. 2. Marginalized joint 68% and 95% CL for $(dn_s/d \ln k, n_s)$ using *Planck*+WP+BAO, either marginalizing over r or fixing $r = 0$ at $k_* = 0.038 \text{ Mpc}^{-1}$. The purple strip shows the prediction for single monomial chaotic inflationary models with $50 < N_* < 60$ for comparison.

dex and its running are set by the slow-roll consistency relations to second order, with $n_t = -r(2 - r/8 - n_s)/8$ and $dn_t/d \ln k = r(r/8 + n_s - 1)/8$. *Planck* measures the running to be $dn_s/d \ln k = -0.016 \pm 0.010$ when tensors are included (see Table 5 and Fig. 4). The constraints on the tensor-to-scalar ratio are relaxed compared to the case with no running, due to an anti-correlation between r and $dn_s/d \ln k$, as shown in Fig. 4 for *Planck*+WP+BAO.

Varying both tensors and running, *Planck*+WP implications for slow-roll parameters are $\epsilon_V < 0.015$ at 95% CL, $\eta_V = -0.014^{+0.015}_{-0.011}$, $|\xi_V^2| = 0.009 \pm 0.006$.

In summary, the *Planck* data prefer a negative running for the scalar spectral index of order $dn_s/d \ln k \approx -0.015$ at the 1.5σ significance level, alone and in combination with other astrophysical data sets. Weak statistical evidence for non-zero negative values of $dn_s/d \ln k$ has been claimed in several previous investigations with the *WMAP* data and smaller scale CMB data (e.g., Spergel et al., 2003; Peiris et al., 2003; Dunkley et al., 2011; Hinshaw et al., 2012a; Hou et al., 2012).

If primordial, negative values for $dn_s/d \ln k$ of order 10^{-2} , would be interesting for the physics of inflation. The running of the scalar spectral index is a key prediction for inflationary models. It is strictly zero for power law inflation, whose fit to *Planck* was shown to be quite poor in the previous section. Chaotic monomial models with $V(\phi) \propto \phi^n$ predict $dn_s/d \ln k \approx -8(n+2)/(4N+n)^2 \approx (n_s-1)^2$, and the same order of magnitude (10^{-3}) is quite typical for many slow-roll inflationary models, such as natural inflation or hilltop inflation, to name a few. It was pointed out that a large negative running of $dn_s/d \ln k \lesssim -10^{-2}$ would make it difficult to support the $N_* \approx 50$ e -foldings required from inflation (Easther & Peiris, 2006), but this holds only without nonzero derivatives higher than the third order in the inflationary potential. Designing inflationary models that predict a negative running of $O(10^{-2})$ with an acceptable n_s and number of e -folds is not impossible, as the case with modulated oscillations in the inflationary potential demonstrates (Kobayashi & Takahashi, 2011). This occurs, for instance, in the axion monodromy model when the instanton contribution is taken into ac-

Model	Parameter	<i>Planck</i> +WP	<i>Planck</i> +WP+lensing	<i>Planck</i> +WP+high- ℓ	<i>Planck</i> +WP+BAO
Λ CDM + $dn_s/d \ln k$	n_s	0.9561 ± 0.0080	0.9615 ± 0.0072	0.9548 ± 0.0073	0.9596 ± 0.0063
	$dn_s/d \ln k$	-0.0134 ± 0.0090	-0.0094 ± 0.0085	-0.0149 ± 0.0085	-0.0130 ± 0.0090
	$-2\Delta \ln \mathcal{L}_{\max}$	-1.50	-0.77	-2.95	-1.45
Λ CDM + $dn_s/d \ln k$ + $d^2 n_s/d \ln k^2$	n_s	$0.9514^{+0.087}_{-0.090}$	$0.9573^{+0.077}_{-0.079}$	$0.9476^{+0.086}_{-0.088}$	$0.9568^{+0.068}_{-0.063}$
	$dn_s/d \ln k$	$0.001^{+0.016}_{-0.014}$	$0.006^{+0.015}_{-0.014}$	$0.001^{+0.013}_{-0.014}$	$0.000^{+0.016}_{-0.013}$
	$d^2 n_s/d \ln k^2$	$0.020^{+0.016}_{-0.015}$	$0.019^{+0.014}_{-0.014}$	$0.022^{+0.014}_{-0.013}$	$0.017^{+0.016}_{-0.014}$
	$-2\Delta \ln \mathcal{L}_{\max}$	-2.65	-2.14	-5.42	-2.40
Λ CDM + r + $dn_s/d \ln k$	n_s	0.9583 ± 0.0081	0.9633 ± 0.0072	0.9570 ± 0.0075	0.9607 ± 0.0063
	r	< 0.25	< 0.26	< 0.23	< 0.25
	$dn_s/d \ln k$	0.021 ± 0.012	0.017 ± 0.012	$-0.022^{+0.011}_{-0.010}$	$-0.021^{+0.012}_{-0.010}$
	$-2\Delta \ln \mathcal{L}_{\max}$	-1.53	-0.26	-3.25	-1.5

Table 5. Constraints on the primordial perturbation parameters for Λ CDM+ $dn_s/d \ln k$, Λ CDM+ $dn_s/d \ln k$ + r and Λ CDM+ $dn_s/d \ln k$ + $d^2 n_s/d \ln k^2$ models from *Planck* combined with other data sets. Constraints on the spectral index and its dependence on the wavelength are given at the pivot scale of $k_* = 0.05 \text{ Mpc}^{-1}$.




Fig. 3. Marginalized joint 68% and 95% CL regions for $(d^2 n_s/d \ln k^2, dn_s/d \ln k)$ using *Planck*+WP+BAO.




Fig. 4. Marginalized joint 68% and 95% CL regions for (r, n_s) , using *Planck*+WP+BAO with and without a running spectral index.

count (McAllister et al., 2010) giving the potential

$$V(\phi) = \mu^3 \phi + \Lambda^4 \cos\left(\frac{\phi}{f}\right). \quad (46)$$

4.4. Open inflation

Most models of inflation predict a nearly flat spatial geometry with small deviations from perfect spatial flatness of order $|\Omega_K| \sim 10^{-5}$. Curvature fluctuations may be regarded as local fluctuations in the spatial curvature, and even in models of inflation where the perturbations are calculated about a spatially flat background, the spatial curvature on the largest scales accessible to observation now are subject to fluctuations from perfect spatial flatness (i.e., $\Omega_K = 0$). This prediction for this fluctuation is calculated by simply extrapolating the power law spectrum to the largest scale accessible today, so that Ω_K as probed by the CMB roughly represents the local curvature fluctuation averaged over our (causal) horizon volume. Although it has sometimes been claimed that spatial flatness is a firm prediction of inflation, it

was realized early on that spatial flatness is not an inexorable consequence of inflation, and large amounts of spatial curvature (i.e., large compared to the above prediction) can be introduced in a precise way while retaining all the advantages of inflation (Gott, 1982; Gott & Statler, 1984) through bubble nucleation by false vacuum decay (Coleman & De Luccia, 1980). This proposal gained credence when it was shown how to calculate the perturbations in this model around and beyond the curvature scale (Bucher et al., 1995; Bucher & Turok, 1995; Yamamoto et al., 1995; Tanaka & Sasaki, 1994). See also (Ratra & Peebles, 1995, 1994; Lyth & Stewart, 1990). For more refined later calculations see for example Garriga et al. (1998, 1999); Gratton & Turok (1999) and references therein. For predictions of the tensor perturbations see for example Bucher & Cohn (1997); Sasaki et al. (1997); Hertog & Turok (2000).

An interesting proposal using *singular* instantons and not requiring a false vacuum may be found in Hawking & Turok (1998), and for calculations of the resulting perturbation spectra see (Hertog & Turok, 2000; Gratton et al., 2000). Models of this sort have been studied more recently in the context of the string




Fig. 5. Joint posterior for tensors and running of the scalar spectral index marginalizing over other parameters. The contours are set at 68% and 95%. The red contours apply for *Planck*+WP+high- ℓ data. The colour of the scattered points indicates the distribution of Ω_K in this case. The blue contours apply when BAO data is also included. (Ω_K is then found to be well-constrained close to zero.) Solid black lines guide the eye to the *no-tensors, no-running* solution.

landscape (see for example [Vilenkin \(2007\)](#) for a nice review). Although some proposals for universes with positive curvature within the framework of inflation have been put forth ([Gratton et al., 2002](#)), it is much harder to obtain a closed universe with a spatial geometry of positive spatial curvature (i.e., $\Omega_K < 0$) ([Linde, 2003](#)).

Theoretically, it is of interest to measure Ω_K to an accuracy of approximately 10^{-4} or slightly better to test the extrapolation of simple flat inflation for this observable. A statistically significant positive value would suggest that open inflation, perhaps in the context of the landscape, was at play. A statistically significant negative value could pose difficulties for the inflationary paradigm. For a recent discussion of these questions, see for example [Freivogel et al. \(2006\)](#), [Kleban & Schillo \(2012\)](#) and [Guth & Nomura \(2012\)](#).

In order to see how much spatial curvature is allowed, we consider a rather general model including the parameters r , n_s , and $dn_s/d \ln k$ as well as Ω_K . We find that $\Omega_K = -0.058^{+0.046}_{-0.026}$ with *Planck*+WP, and $\Omega_K = -0.0004 \pm 0.00036$ with *Planck*+WP+BAO. More details can be found by consulting the parameter tables available online.⁶ Figure 5 shows r and $dn_s/d \ln k$ for this family of models. We conclude that any possible spatial curvature is small in magnitude even within this general model and that the spatial curvature scale is constrained to lie far beyond the horizon today. Open models predict a tensor spectrum enhanced at small wavenumber $k \lesssim 1$, where $k = 1$ corresponds to the curvature scale, but our constraint on Ω_K and cosmic variance imply that this aspect is likely unobservable.

4.5. Relaxing the assumption of the late-time cosmological concordance model

The joint constraints on n_s and r shown in Fig. 1 are one of the central results of this paper. However, they are derived assuming the standard Λ CDM cosmology at late times (i.e., $z \lesssim 10^4$). It is therefore natural to ask how robust our conclusions are to changes of the late time cosmological model. We discuss two classes of models: firstly, changes to the Λ CDM energy content, and secondly, a more general reionization model. These extensions can lead to degeneracies of the additional parameters with n_s or r .

4.5.1. Extensions to the energy content

We consider the Λ CDM+ r + N_{eff} , Λ CDM+ r + Y_p , Λ CDM+ r + $\sum m_\nu$ and Λ CDM+ r + w extensions of the standard model. This selection is motivated by the impact on the CMB damping tail of the first two and the effect on the Sachs-Wolfe plateau at low multipoles for the latter two. The resulting contours are shown in Fig. 6. While the lower limit on n_s is stable under all extensions considered here, the models that affect the high- ℓ part of the spectrum permit significantly bluer spectral tilts, and accordingly also leads to a weaker bound on the tensor-to-scalar ratio (Λ CDM+ r + N_{eff} gives $n_s = 0.9764 \pm 0.0106$ and $r_{0.002} < 0.15$ at 95% CL, while for Λ CDM+ r + Y_p we obtain $n_s = 0.9810 \pm 0.0111$ and $r_{0.002} < 0.18$ at 95% CL). The models affecting the large-scale part of the power spectrum, on the other hand, do not lead to a notable degradation of constraints on either n_s or r ($n_s = 0.9648 \pm 0.0061$ and $r_{0.002} < 0.13$ at 95% CL for Λ CDM+ r + $\sum m_\nu$, and $n_s = 0.9601 \pm 0.0070$ and $r_{0.002} < 0.11$ at 95% CL for Λ CDM+ r + w).

4.5.2. General reionization scenario

In the standard rapid reionization scenario typically used in CMB analysis, the Universe is assumed to be completely transparent after recombination, but the ionization fraction increases from zero to one over a duration $\Delta z \approx 1$ at a certain redshift z_{reion} , which is the only unknown parameter of the reionization model. This model is obviously simplistic, but for CMB analysis it works quite well because the CMB has little sensitivity to the details of how the ionization fraction changes from 0 to 1. In this section we study to what extent allowing more general reionization scenarios may alter some of the conclusions concerning the constraints on n_s and r as well as τ . As discussed in [Mortonson & Hu \(2008a\)](#) and [Mortonson & Hu \(2008b\)](#), CMB anisotropies constrain reionization almost entirely by using the shape of the large-scale EE power spectrum, and the power is redistributed from larger to smaller scales for reionization processes which take place during a non-negligible redshift interval, since they start at an earlier epoch.

We use the method developed by [Mortonson & Hu \(2008a\)](#) to describe and constrain the reionization history. A complete principal component basis serves to describe the effect of reionization on the large-scale E -mode polarization power spectrum. Following [Mortonson & Hu \(2008a\)](#) we bin the ionization history $x_e(z_i)$ using 95 equal width bins with $\Delta z = 0.25$ ranging from $z_{\text{min}} = 6$ to $z_{\text{max}} = 30$. For the redshifts $z < z_{\text{min}}$ we assume values for x_e which take into account first (and possibly second) helium ionization and complete hydrogen ionization ($x_e = 1.16$ for $z < 3$ and $x_e = 1.08$ for $z < 6$). For $z > 30$ we fix $x_e = 2 \times 10^{-4}$ as the value of x_e expected before reionization (and after primordial recombination). Any reionization

⁶ Available at: http://www.sciops.esa.int/index.php?project=planck&page=Planck_Legacy_Archive




Fig. 6. Marginalized joint 68% and 95% CL regions for *Planck*+WP+BAO data for Λ CDM+ r + N_{eff} and Λ CDM+ r + Y_{He} (top); and Λ CDM+ r + $\sum m_\nu$ and Λ CDM+ r + w (bottom). Shown for comparison are the corresponding contours for the Λ CDM+ r model.

history can be parametrized as a free function of redshift by decomposing the ionization fraction as $x_e(z) = x_e^f(z) + \sum_\mu m_\mu S_\mu(z)$, where the principal components, $S_\mu(z)$, are the eigenfunctions of the Fisher matrix computed by taking the derivatives of the EE polarization power spectrum with respect to $x_e(z)$. Following Mortonson & Hu (2008a) we consider here the first five eigenfunctions, $S_\mu(z)$ with $\mu = 1, \dots, 5$, which will be varied with the other cosmological parameters.

In Fig. 7 we plot the 68% and 95% CL regions for n_s and r . The constraint on the tensor-to-scalar ratio is not significantly affected by this additional marginalization, increasing to $r_{0.002} < 0.13$ at 95% CL. The scalar spectral index is increased to $n_s = 0.9650 \pm 0.0080$, compared with $n_s = 0.9603 \pm 0.0073$ obtained with the rapid reionization scheme. This is the same trend as noted in Pandolfi et al. (2010) using *WMAP* data, but the effect is less significant due to the improved measurement of the spectrum by *Planck*. The larger freedom in the reionization history increases the posterior on the derived optical depth, which is still partially degenerate with the scalar spectral index of primordial perturbations. The $n_s = 1$ model is still excluded at high significance; however, we find $\Delta\chi^2_{\text{eff}} = 12.5$ compared to the Λ CDM model.




Fig. 7. Marginalized joint 68% and 95% CL regions for *Planck*+WP data for Λ CDM+ r for instantaneous and general reionization.

5. Inflationary model comparison

In Sect. 4.2 several representative families of parameterized models for the inflationary potential were analysed within the slow-roll approximation in the neighbourhood of the pivot scale k_* . Approximate constraints were applied to reject models for which there is no plausible scenario for entropy generation. In this section we revisit some of the parametric models defined in Sect. 4.2. Here, however, the modes for the first order perturbations, as described in Sect. 2.2, are integrated numerically. Thus there is no slow-roll approximation, and the issue of the existence of a plausible scenario for entropy generation is examined more carefully. We perform a statistical model comparison between the competing parameterized potentials, both within the framework of Bayesian model comparison and in terms of the relative likelihoods of the best-fit models from each parameterized family.

As noted in Sect. 2.3, there is considerable uncertainty surrounding what occurs during the epoch of entropy generation, particularly with respect to the energy scale at which entropy generation ends, and the average equation of state between that epoch and the end of inflation. For this reason, we explore a number of scenarios for this intermediate era during which entropy generation takes place. The models compared in this section include inflation with power law potentials, defined in Eq. 34, with several representative values for n ; natural inflation, in Eq. 39; and hilltop inflation, with $p = 4$ and $\lambda = 4\Lambda^4/\mu^4$ in Eq. 37.

The free parameters in these inflationary potentials may vary over several orders of magnitude corresponding to unknown scales in high energy particle physics. Consequently a logarithmic prior is a sensible choice for these parameters. However, there is no theoretical guidance on how to truncate these model priors. We therefore adopt broad priors initially, and then truncate them as follows in order to compare all the models on an equal footing.

The strongest constraint on the inflationary parameter space comes from the amplitude of the primordial power spectrum. This is a free parameter in most models, and successful structure formation in a universe dominated by cold dark matter has long been known to require primordial fluctuations with $\mathcal{R} \approx 10^{-5}$,

Model	Priors
$n = 2/3$	$-13 < \log_{10}(\lambda) < -7$
$n = 1$	$-13 < \log_{10}(\lambda) < -7$
$n = 2$	$-13.5 < \log_{10}(m^2) < -8$
$n = 4$	$-16 < \log_{10}(\lambda) < -10$
Natural	$-5 < \log_{10}(\Lambda) < 0$ $0 < \log_{10}(f) < 2.5$
Hilltop	$-8 < \log_{10}(\Lambda) < -1$
$p = 4$	$-17 < \log_{10}(\lambda) < -10$
Λ CDM	$0.9 < n_s < 1.02$ $3.0 < \log_{10}[10^{10} A_s] < 3.2$

Matching	Prior
N_*	$20 \leq N \leq 90$

Table 6. Model priors. Dimensionful quantities are expressed in units with M_{pl} set to unity. The matching parameter N_* , computed at $k_* = 0.05 \text{ Mpc}^{-1}$, allows us to marginalize over the uncertainty in connecting the inflationary era to astrophysical scales. Additional cuts are made in these parameter ranges to select out physically relevant subspaces which are compatible with basic structure-formation requirements and different entropy-generation scenarios, as described in the text, resulting in non-rectangular model priors in some cases. We also marginalize over the concordance cosmological parameters and foreground parameters of the *Planck* likelihood, as described in the text.

or $A_s \approx 10^{-10}$ (see e.g., [Zeldovich, 1972](#); [Linde, 1990](#)). We can therefore immediately reject models for which A_s is far from this value, so regions of parameter space which do not yield $10^{-11} \leq A_s \leq 10^{-7}$ are *a priori* excluded. This range is generous relative to estimates of A_s prior to *Planck* (e.g., [Komatsu et al., 2011](#)), but the results do not depend strongly on the range chosen. This effectively truncates the logarithmic priors on the model parameters, leaving a parameter subspace compatible with basic structure formation requirements.

For the single parameter models this requirement defines the range of λ in Eq. 34. However, for generic multi-parameter models, an A_s -based cut may select a nontrivial region of parameter space, as happens for the two cases considered here (see e.g., Fig. 1 in [Easther & Peiris \(2012\)](#)). Without the A_s -based cut in the prior, the parameter volume for both natural and hilltop inflation would be rectangular, and the corresponding Bayesian evidence values computed for these models would be lowered.

As discussed in Sect. 2.3, specifying an inflationary potential does not enable us to predict the late-time CMB angular power spectra. The subsequent expansion history and details of the epoch of entropy generation are required to relate the value of the inflaton field at horizon crossing to comoving wavenumbers in today’s Universe, via Eq. 24. Physically, the fundamental parameter that sets the observable perturbation spectrum is the value of the field ϕ_* at which the pivot mode leaves the horizon. It can be rescaled by a shift $\phi_* \rightarrow \phi_* + \phi_0$, and the range over which ϕ changes during inflation varies greatly between models. Consequently we treat the remaining number of e -folds, N_* , after the pivot scale leaves the horizon as a free parameter with a wide uniform prior, since this quantity has a consistent interpretation across models. The pivot scale used to compute N_* is $k_* = 0.05 \text{ Mpc}^{-1}$. However, given our ignorance concerning the epoch of entropy generation, a multitude of entropy generation

scenarios for each inflationary potential can occur. Some possibilities are as follows, with parameters referring to Eq. 24.

1. *Instantaneous entropy generation.*
At the end of inflation, all the energy in the inflaton field is instantaneously converted into radiation.
2. *Restrictive entropy generation scenario (narrow range for w_{int}).*
 $\rho_{\text{th}}^{1/4} = 10^9 \text{ GeV}$, and $w_{\text{int}} \in [-1/3, 1/3]$.
3. *Permissive entropy generation scenario (wide range for w_{int}).*
 $\rho_{\text{th}}^{1/4} = 10^3 \text{ GeV}$, and $w_{\text{int}} \in [-1/3, 1]$.

The equations of state with w_{int} in the range $[1/3, 1]$ appear less plausible, but models with these values have been put forward ([Pallis, 2006](#)), so this cannot be completely excluded. Moreover the w_{int} parameterization captures a variety of scenarios in which the post-inflationary Universe is thermalized, but not radiation-dominated, including phases of coherent oscillations ([Martin & Ringeval, 2010](#); [Easther et al., 2011](#)), resonance ([Traschen & Brandenberger, 1990](#); [Kofman et al., 1997](#); [Allahverdi et al., 2010](#)), kination ([Spokoyny, 1993](#); [Chung et al., 2007](#)), secondary or thermal inflation ([Lyth & Stewart, 1996](#)), moduli domination ([Banks et al., 1994](#); [de Carlos et al., 1993](#)), primordial black hole domination ([Anantua et al., 2009](#)), or a frustrated cosmic string network ([Burgess et al., 2005](#)), all of which lead to an expansion rate differing from that of a radiation-dominated universe.

At the other extreme, the decision to exclude $w_{\text{int}} < -1/3$, as done here, is not completely justifiable. We cannot, for example, rule out a first-order phase transition at a lower energy scale that would drive w_{int} below $-1/3$, but here we neglect this possibility. Our analysis does not preclude a secondary period of inflation, but does require that the *average* expansion during the post-inflationary regime parameterized by w_{int} should not be inflationary. This caveat should be kept in mind.

For some of the parameterized models, tighter constraints can, in principle, be placed on w_{int} . It has been argued (see e.g., [Liddle & Leach, 2003](#)) that for the $\lambda\phi^4$ potential, the uncertainties concerning entropy generation contribute almost no uncertainty in the determination of ϕ_* . This is because according to the virial theorem, a field sloshing about $\phi = 0$ in a quartic potential has the same average w , namely $w = 1/3$, as the radiation equation of state. More generally, for a potential of the form ϕ^n around the minimum, $w_{\text{vir}} = \langle w \rangle = (n - 2)/(n + 2)$ ([Turner, 1983](#)); therefore, one may argue that w_{int} should be restricted to the interval whose endpoints are $1/3$ and w_{vir} . This approach was taken by [Martin & Ringeval \(2010\)](#) in obtaining Bayesian constraints on the reheating temperature for monomial potentials from the CMB. However this scenario requires a carefully tuned potential which has approximately a ϕ^n shape both at large field values, and near the origin, far below the inflationary scale. Typically, potentials for which $V(\phi) \sim \phi^n$ at large field values can have very different shapes near the origin. Thus, following [Easther & Peiris \(2012\)](#), in this paper we choose to explore a broader range of w_{int} for these models (including the cases above as subsets), in order to obtain data-driven constraints on w_{int} .

In this paper we focus on the three representative scenarios itemized above, referred to hereafter as: (1), (2), and (3). Our algorithm draws a value of N_* and then, given the value of ρ_{th} , computes w_{int} , which is a derived parameter—models for which w_{int} lies outside the specified range of each scenario under consideration are excluded.

Our full set of priors for the inflationary physics is given in Table 6. Dimensionful quantities are expressed in units with reduced Planck mass M_{pl} set to unity.

Model	Instantaneous entropy generation		Restrictive entropy generation		Permissive entropy generation	
	$\ln[\mathcal{E}/\mathcal{E}_0]$	$\Delta\chi^2_{\text{eff}}$	$\ln[\mathcal{E}/\mathcal{E}_0]$	$\Delta\chi^2_{\text{eff}}$	$\ln[\mathcal{E}/\mathcal{E}_0]$	$\Delta\chi^2_{\text{eff}}$
$n = 4$	-14.9	25.9	-18.8	27.2	-13.2	17.4
$n = 2$	-4.7	5.4	-7.3	6.3	-6.2	5.0
$n = 1$	-4.1	3.3	-5.4	2.8	-4.9	2.1
$n = 2/3$	-4.7	5.1	-5.2	3.1	-5.2	2.3
Natural	-6.6	5.2	-8.9	5.5	-8.2	5.0
Hilltop	-7.1	6.1	-9.1	7.1	-6.6	2.4
ΛCDM	-4940.7	9808.4

Table 7. Inflationary model comparison results. For each model and set of assumptions concerning entropy generation [(1), (2), (3)], the natural logarithm of the Bayesian evidence ratio as well as $\Delta\chi^2_{\text{eff}}$ for the best-fit model in each category are indicated, relative to the ΛCDM concordance model (denoted by subscript “0”); $\ln \mathcal{E}_0$ and $-2 \ln \mathcal{L}_0$ for the latter are also given.

Due to the non-trivial likelihood surfaces and the large dimensionality of the parameter spaces explored in this section, we use ModeCode coupled with MultiNest v3.0⁷ to map out the parameter space. In addition to the standard nested sampling (NS) algorithm, MultiNest v3.0 enables nested importance sampling (NIS), resulting in substantial speed gains⁸ and significantly enhancing the accuracy of the Bayesian evidence computation over NS alone for the same computational setup.

5.1. Results

Table 7 presents model comparison results for the ensemble of parameterized potential families described above. We report the Bayesian evidence (*model-averaged likelihood*) ratio, which provides a self-consistent framework for calculating the *betting odds* between models (see Sect. 3.4). The uncertainty in these log evidence values is ≈ 0.2 . We also report the $\Delta\chi^2_{\text{eff}}$ values computed from the $2 \ln \mathcal{L}_{\text{max}}$ values found by the sampler.

The monomial models have a single-parameter potential, and the natural and hilltop inflation models have two parameters each. All reheating scenarios except case (1) contribute one additional parameter to the inflationary sector. The evidence ratios and $\Delta\chi^2_{\text{eff}}$ values are presented with respect to the ΛCDM cosmological model.

None of the inflationary models tested here fit the data as well as the ΛCDM model. This mostly reflects that there is no evidence in the data for r different from zero. Furthermore, the priors listed in Table 6 for the ΛCDM primordial sector are purely phenomenological, roughly corresponding to ranges somewhat broader than WMAP constraints. Narrowing them around the best-fit model arbitrarily increases the evidence in its favour. Instead it is instructive to compare the relative evidence for the inflationary models presented.

Table 7 shows that the $\lambda\phi^4$ model is decisively ruled out by Planck, confirming previous analyses by the WMAP team (Peiris et al., 2003; Spergel et al., 2007; Dunkley et al., 2009; Komatsu et al., 2011) based on the model track plotted on the $n_s - r$ plane. Recent model selection analyses (Martin et al., 2011; Easther & Peiris, 2012) with WMAP7 found that the model was already under severe pressure, disfavoured by odds of $\sim 400:1$ against.

With Planck, the odds against this model are *at least* 500,000:1 when compared with ΛCDM , for a very broad set of reheating scenarios. The same conclusion is reflected by the extremely poor $\Delta\chi^2_{\text{eff}}$ values for the model. Given the strength of our results in the flexible setting of the permissive entropy generation scenario, it is possible not just to rule out models where the potential is of the quartic form in the full range from the origin to the inflationary scales, but also a general class where the potential is of the $n = 4$ form in the ϕ -range where the cosmological perturbations are generated, but exhibits a different shape near the origin.

Two other large field models, the quadratic potential and natural inflation, are under some pressure from Planck data, especially when broader entropy generation scenarios are considered. Compared with the ΛCDM model, these models are disfavoured by $\Delta\chi^2_{\text{eff}} \sim 5-6$ depending on the reheating scenario. This reflects the analysis of Sect. 4, where the overlap of the model predictions and the data constraints on the $n_s - r$ plane is seen to be mostly outside the joint 68% CL contour. However, from the Bayesian evidence point of view, it is too early to declare these models incompatible with the data. To make this judgement, it is more conservative to compare these models with the $n = 1$ case, which has the best evidence with respect to ΛCDM , rather than with ΛCDM itself, which provides our reference point for the evidence calculation, but has arbitrary prior ranges. In their simplest forms – instant entropy generation – the $n = 2$ and natural inflation models are only disfavoured by odds of $\sim 1-12:1$ against, which does not rise to a high level of significance⁹.

The models most compatible with the Planck data in the set considered here are the two interesting *axion monodromy* potentials, $n = 1$ (McAllister et al., 2010), and $n = 2/3$ (Silverstein & Westphal, 2008), which are motivated by inflationary model building in the context of string theory. The $p = 4$ hilltop model presents an interesting case. This model was previously found to be compatible with WMAP7 data, performing almost as well as the monodromy potentials (Easther & Peiris, 2012). However it exhibits significant tension with the Planck data, both in terms of evidence ratios and the maximum likelihood. The only exception is the entropy generation scenario (3) which has odds of $> 1000:1$ against compared to ΛCDM , and yet the maximum likelihood is not significantly different to the $n = 1$ case. This indicates that while the extra freedom allowed by the least conservative reheating scenario improves the best fit, this prior is not very predictive of the data. However, the result is counterin-

⁷ Made available ahead of public release to the Planck Collaboration by Farhan Feroz and Mike Hobson.

⁸ We have made extensive tests of NIS vs. NS, and chosen the following settings for the computations presented here: NIS on, constant efficiency mode on, 300 live points, tolerance and efficiency parameters set to 0.5 and 0.02, respectively.

⁹ In comparison, odds of 150:1 are considered highly significant in this context.




Fig. 8. Constraints on n_s vs. r at $k_* = 0.002 \text{ Mpc}^{-1}$ for the inflationary models considered (i.e., power law potentials with $n = 2/3, 1, 2$, and 4 , natural inflation, and hilltop inflation), showing joint 68% and 95% CL. Blue and grey distributions correspond to the restrictive and permissive entropy generation scenarios, respectively. The instant reheating case corresponds to the thin (red) contours in the natural and hilltop panels; for the single parameter models, this case corresponds to the lowest- r extremity of restrictive case. The difference between the *natural inflation* region on Fig. 1 and the natural inflation constraints shown here is due to the strong projection effect described in the text.




Fig. 9. Constraints on n_s vs. w_{int} at $k_* = 0.002 \text{ Mpc}^{-1}$ for the inflationary models considered, as in Fig. 8. The instant reheating case (1) corresponds to $w_{\text{int}} = 1/3$.

tuitive and merits further comment; we will consider it further at the end of this section when parameter estimation results are discussed.

We now turn to parameter constraints. Fig. 8 presents marginalized joint constraints from *Planck* alone on the derived parameters $n_{s,0.002}$ and $r_{0.002}$; Fig. 9 shows the corresponding joint constraints on $n_{s,0.002}$ and w_{int} , again a derived parameter,




Fig. 10. Constraints on $\ln(\lambda)$ vs. w_{int} for the inflationary models considered, as in Figs. 8 and 9.

and Fig. 10 shows joint constraints on w_{int} vs. the potential parameter $\log(\lambda)$ ($\log(\Lambda)$ in the case of natural inflation). It is instructive to consider the three sets of figures together. The restrictive and permissive entropy generation scenarios are shown on all panels; the instant reheating case is shown for the two parameter models, natural inflation and hilltop, in Fig. 8 – for the monomial potentials this case corresponds to the lowest- r extremity of the restrictive case constraints, and to $w_{\text{int}} = 1/3$ in the other two figures.

The quartic potential conflicts with the data because it predicts a high tensor-to-scalar ratio. Hence the model maximizes its likelihood by pushing towards the lower- r , bluer- n_s limits of its parameter space, which corresponds to increasing w_{int} as allowed by the reheating prior. The contours terminate at the lowest- r limit when each reheating case hits its w_{int} prior upper limit (i.e., $w_{\text{int}} = 1/3$ and $w_{\text{int}} = 1$ for restrictive and permissive entropy generation, respectively). For each case, the lower limit on w_{int} , corresponding to the reddest- n_s extremity of the confidence contours, is data-driven. The quadratic potential encounters the same difficulty but at a less extreme level.

The two axion monodromy potentials are compatible with a wide range of entropy generation scenarios. The instantaneous reheating scenario is compatible with the data for both models. For restrictive entropy generation in the $n = 1$ model, we obtain a data-driven upper limit on w_{int} , which just touches the $w_{\text{int}} = 1/3$ case at the 95% CL. At the lower limit, the w_{int} posterior is truncated by the prior, as for the $n = 2/3$ case. For the latter, there is a data-driven upper limit on w_{int} which is controlled by the upper limit on n_s . For permissive entropy generation, the upper and lower limits on w_{int} for both models are data-driven, corresponding to the upper and lower limits on n_s respectively.

The constraints on natural inflation require some interpretation. The relationship between the empirical n_s and r parameters and the potential parameters for natural inflation is discussed in detail by Savage et al. (2006) and Mortonson et al. (2011) along with parameter constraints derived from WMAP3 (Spergel et al., 2007) and WMAP7 (Larson et al., 2011), respectively. In this

model, there is a degeneracy between f and Λ in the limit where these parameters are large so that natural inflation resembles the quadratic model. Our priors are chosen to exclude most of this region. The priors on $\log(f)$ and $\log(\Lambda)$ still allow a region of nearly degenerate models that contribute to a *ridge* seen in the natural inflation panel of Fig. 8. These models closely match the values of n_s and r seen in the quadratic potential constraints. The marginalized constraints on n_s and r depend strongly on the prior on $\log(f)$ due to the projection of a large number of degenerate models into this ridge. Therefore the apparent preference for this region of parameter space over models with lower values of r is largely due to this effect and is not driven by the data. This highly nonlinear mapping between the logarithmic priors on the potential parameters and the power law parameters (which are derived parameters in this analysis) leads to a strong projection effect that accounts for the difference in visual appearance between these contours and the region labeled ‘natural inflation’ in Fig. 1.

Generally, for fixed N_* , decreasing Λ and f reduces both n_s and r . Thus natural inflation models can have lower values of r than the quadratic potential without increasing n_s and N_* . This feature means that the potential parameters for this model are relatively uncorrelated with w_{int} , in contrast with the other models considered here, as illustrated in the Fig. 10. Nevertheless we obtain data-driven bounds on w_{int} in the permissive reheating case as well as a lower bound in the restrictive reheating case. Both bounds overlap with the instantaneous reheating limit.

The left panel of Fig. 11 shows our lower limit on $f \gtrsim 10.0 M_{\text{pl}}$ (95% CL), compared with the WMAP7 limit $f \gtrsim 5.0 M_{\text{pl}}$ (95% CL) reported by Mortonson et al. (2011). Indeed, the Planck limit is in agreement with the Planck prediction presented in that work. There is a hint of an upper limit on f as well, driven by the fact that this corresponds to the quadratic inflation limit, which is in tension with the data. However this is only a 1σ effect.

The $p = 4$ hilltop model has two distinct branches: a *small-field* scenario, where $r \lesssim 0.001$ and $n_s < 0.95$, and a *large-*




Fig. 11. Potential parameters for natural inflation and hilltop inflation, as in Figs. 8, 9, and 10. On the natural inflation panel, instant reheating corresponds to a thin diagonal along the top edge of reheating case (3).

field limit in which $V(\phi) \sim \phi$ (Adshead et al., 2011). Physically, the small-field limit is consistent with the Lyth bound, and we can select it by fixing $\log(\Lambda) < -2.5$ in the prior. We observe that the data select out this small field branch, which requires explanation given that we know that the $n = 1$ model (the limit of the large field branch) is perfectly compatible with the data. In fact, this can be tested by restricting the $\log(\Lambda)$ prior by hand to the large-field branch. The hilltop model constitutes a difficult sampling problem, as is apparent from Fig. 4 in Easther & Peiris (2012). An examination of the nested sampler’s progress in the case of the full hilltop prior reveals the reason for the small-field branch being selected out. In this case, the posterior for the large-field branch is extremely thin compared to the model prior in this regime – much thinner than the posterior for the small-field branch in comparison to its respective prior. Therefore this region occupies very little probability mass, and is dropped in preference to the more predictive small-field branch. This high-likelihood but extremely thin ridge is also responsible for the counterintuitive result reported in the model-selection analysis, where the model was found to have a good $\Delta\chi^2_{\text{eff}}$ with respect to ΛCDM , and yet be highly disfavoured by the Bayesian evidence.

In summary, confirming the results of Sect. 4, this model is in agreement with the data in the limit where it overlaps the linear $n = 1$ model. But since this region of high likelihood occupies a very small fraction of the prior, this model is heavily penalized by the Bayesian evidence for failing to predict the data over most of its prior space.

6. Observable window of inflation

Section 4.2 presented an analysis of several representative inflationary potentials within the framework of the slow-roll approximation, and their compatibility with the *Planck* data. The results are summarized in Fig. 1. In that case the full potential is considered in order to identify a plausible range for the location of ϕ_* on the potential $V(\phi)$. This requires a complete story: i.e., the potential must be specified starting above the point where the largest observable scales first exited the Hubble horizon, and extending to the minimum of the potential.

In this section we explore another approach. We adopt the point of view that we are interested in reconstructing the inflationary potential only over the observable range—that is, the interval of ϕ corresponding to the scales observable today in the CMB. We constrain the potential over the range where these scales exited the horizon during inflation, as well as a few e -folds before and after – the cosmological perturbations are not imprinted instantaneously at the moment of horizon crossing, but rather, gradually over a few e -folds. We expand around ϕ_* , taking the view that a plausible extension of the potential outside this observable range is always possible, so that one has precisely the number of e -folds of inflation needed for ϕ_* to correspond to k_* today.

The argument is that one can always end inflation abruptly by imposing a sharp waterfall feature where needed, or prolong inflation by inserting a sufficiently long plateau into the potential by hand, e.g., for models with a large tilt. A foreseeable objection to this approach is that the extensions of the potential required outside the observable window may render the potential *unnatural*. This possibility should be kept in mind, although *naturalness* is an elusive and uncomfortably subjective concept. The analysis in this section does not rely on the slow-roll approximation. Instead each k mode is integrated exactly by numerical integration, as described in Sect. 2.2, under the assumption of a canonical kinetic term.

Two complementary approaches have been taken in the literature to reconstruct the potential. In the first approach, which we follow in this paper, we expand the potential $V(\phi)$ directly in powers of $(\phi - \phi_*)$. In this case the numerical integration of the slow-roll solution must start sufficiently early so that any initial transient has had a chance to decay, and one is in the attractor solution when the dynamics of the largest observable modes in the Universe today start to have an interesting evolution. This is the approach followed, e.g., by Lesgourgues & Valkenburg (2007); Mortonson et al. (2011) using publicly available codes¹⁰.

A second approach expands $H(\phi)$ as a Taylor series in $(\phi - \phi_*)$. As discussed in Sect. 2.2, this has the advantage that $H(\phi)$ determines both the potential $V(\phi)$ and the solution $\phi(t)$, so the issue of having to start sufficiently early in order to allow the initial transient to decay is avoided. This method was used in, e.g., Kinney (2002), Kinney et al. (2006), Peiris & Easther (2006a), Easther & Peiris (2006), Peiris & Easther (2006b) and Peiris

¹⁰ <http://wwwlapp.in2p3.fr/valkenbu/inflationH/>,
<http://zuserver2.star.ucl.ac.uk/~hiranya/ModeCode/>.




Fig. 12. Posterior distribution for the first three slow-roll potential parameters using *Planck*+WP data. In the $n = 2-4$ cases, the inflaton potential is expanded to the n th order, and the spectrum is obtained by fitting the numerically-computed power spectrum to the data, with no slow-roll approximation, and no assumption about the extrapolation of the potential outside of the observable window.




Fig. 13. Posterior distribution for the Taylor expansion coefficients, V_i , of the inflaton potential. The potential is expanded at n th order, assuming a flat prior on ϵ_V , η_V , ξ_V^2 , ϖ_V^3 . The coefficients V_i are expressed in natural units (where $\sqrt{8\pi}M_{\text{pl}} = 1$). The contours show only half of the allowed regions for potential parameters: the other half is symmetric, with opposite signs for V_1 and V_3 .

& Easther (2008), using analytic and semi-analytic approximations, and in Lesgourgues et al. (2008), Powell & Kinney (2007), Hamann et al. (2008c) and Norena et al. (2012), using a fully numerical approach.

These approaches could lead to results that differ from those in Sect. 4.2. Firstly, if the running of the index is large, the slow-roll approximation taken to second order is not necessarily accurate for all models allowed by the data. The relation between the spectral parameters ($\ln A_s$, n_s , $dn_s/d\ln k$, r) and the underlying inflationary potential $V(\phi)$ is therefore uncertain. Secondly, for




Fig. 14. Observable range of the best-fitting inflaton potentials, when $V(\phi)$ is Taylor expanded at the n th order around the pivot value ϕ_* , in natural units (where $\sqrt{8\pi}M_{\text{pl}} = 1$), assuming a flat prior on ϵ_V , η_V , ξ_V^2 , and ϖ_V^3 , and using *Planck*+WP data. Potentials obtained under the transformation $(\phi - \phi_*) \rightarrow (\phi_* - \phi)$ leave the same observable signature and are also allowed. The sparsity of potentials with a small $V_0 = V(\phi_*)$ comes from the flat prior on ϵ_V rather than on $\ln(V_0)$: in fact V_0 is unbounded from below.

Table 8. Numerical reconstruction of potential parameters, compared to results with the slow-roll approximation, when tensors and running are included (*Planck*+WP 95% CL, with $k_* = 0.05 \text{ Mpc}^{-1}$). The effective χ^2 value is given relative to the model with a quadratic potential.

n	from $V(\phi)$			from slow-roll
	2	3	4	
ϵ_V	< 0.0078	< 0.015	< 0.021	< 0.015
η_V	$-0.011^{+0.018}_{-0.015}$	$-0.016^{+0.028}_{-0.025}$	$0.022^{+0.052}_{-0.047}$	$-0.014^{+0.030}_{-0.022}$
ξ_V^2	–	$0.011^{+0.012}_{-0.011}$	$-0.015^{+0.031}_{-0.032}$	$0.009^{+0.011}_{-0.011}$
ϖ_V^3	–	–	$0.016^{+0.018}_{-0.019}$	–
$\Delta\chi^2_{\text{eff}}$	0	–0.7	–3.7	–0.9

spectra with a large running, there is no guarantee that an inflationary model giving such a spectrum exists. All allowed models have $r \ll 1$, so these models are consistent with $\epsilon_V(k_*) \ll 1$ at the pivot scale. However, towards the edge of the observable range, the potential may become incompatible with $\epsilon_V(k) < 1$ (i.e., with the requirement of inflationary expansion). These possible pitfalls are avoided using the methods in this section, since the data have been fit directly by the candidate $V(\phi)$ or $H(\phi)$, computed numerically without any slow roll assumptions over the entire observable range.

We define a class of models over the observable range based on the expectation that the potential should be smooth. $V(\phi)$ is approximated by a Taylor expansion up to order n , and we explore the cases $n = 2, 3$, and 4. For each $V(\phi)$, we integrate over inflationary fluctuations using the inflation module implemented in CLASS¹¹ (Lesgourgues, 2011; Blas et al., 2011), as described in Lesgourgues & Valkenburg (2007). Potentials are rejected for

¹¹ <http://class-code.net>

Table 9. The scalar amplitude, tilt, running, running of the running, and tensor-to-scalar ratio inferred from a numerical reconstruction of the inflaton potential (*Planck*+WP 95% CL, with $k_* = 0.05 \text{ Mpc}^{-1}$).

n	from $V(\phi)$		
	2	3	4
$\ln[10^{10} A_s]$	$3.087^{+0.050}_{-0.050}$	$3.115^{+0.066}_{-0.063}$	$3.130^{+0.071}_{-0.066}$
n_s	$0.961^{+0.015}_{-0.015}$	$0.958^{+0.017}_{-0.016}$	$0.954^{+0.018}_{-0.018}$
$100 \, dn_s/d \ln k$	$-0.05^{+0.13}_{-0.14}$	$-2.2^{+2.2}_{-2.3}$	$-0.61^{+3.1}_{-3.1}$
$100 \, d^2 n_s/d \ln k^2$	$-0.01^{+0.73}_{-0.75}$	$-0.3^{+1.0}_{-1.2}$	$6.3^{+8.6}_{-7.8}$
r	< 0.12	< 0.22	< 0.35

which the attractor solution cannot be reached when the largest observable scales cross the Hubble radius. The parameters we sample are the potential and its derivatives at the pivot scale, when k_* crosses the Hubble radius during inflation.

To avoid parameter degeneracies, we impose uniform priors on ϵ_V , η_V , ξ_V^2 , and ϖ_V^3 at the pivot field value ϕ_* . The advantage of uniform priors on these parameters is that – to the extent that the slow-roll conditions are satisfied – these coefficients relate linearly to observable quantities such as n_s , r , $dn_s/d \ln k$ and $d^2 n_s/d \ln k^2$. Fig. 12 and Table 8 show the posterior probability for these coefficients, and Fig. 13 shows the posterior probability for the Taylor series coefficients V_i . In Fig. 14, we show the observable range of the best-fitting inflaton potentials (for a sample extracted randomly from the converged Markov chains). The edges of the observable range correspond to Hubble crossing for the minimum and maximum values of k used in the Boltzmann code.

When fitting $V(\phi)$ for each model in parameter space, we compute $(n_s, dn_s/d \ln k, d^2 n_s/d \ln k^2, r)$ at the pivot scale *a posteriori* directly from the numerical primordial spectra. The results are shown in Table 9 and can be compared with those of Table 5 in Sect. 4. We can also use the results of Sect. 4 for the $\Lambda\text{CDM}+r+dn_s/d \ln k$ model to infer the potential parameters (ϵ_V , η_V , ξ_V^2) using the second-order slow-roll expressions, and compare different approaches in the space of potential parameters (see Fig. 12 and the last column in Table 8).

The model with a quadratic potential in the observable window ($n = 2$) leads to bounds on ϵ_V , η_V , n_s , and r very close to the $\Lambda\text{CDM}+r$ case. This is not a surprise since such potentials cannot simultaneously give values of n_s and r compatible with the data, and a large running. A significant $dn_s/d \ln k$ can be generated only in the presence of a large ξ_V^3 (i.e., with a significant V'''). Since quadratic potentials produce little running, they are faithfully described by the slow-roll approximation.

The model with a cubic potential ($n = 3$) has the freedom to generate a large running, $dn_s/d \ln k$. Indeed one can check that the results for the $n = 3$ model are close to those of the $\Lambda\text{CDM}+r+dn_s/d \ln k$ model presented in Sect. 4. The agreement between these two models remains very good, despite the fact that in the presence of a large running, the slow-roll approximation can become inaccurate. The running in a potential with $n = 3$ is not exactly scale invariant; this is not captured by the $\Lambda\text{CDM}+r+dn_s/d \ln k$ parameterization.

The $n = 4$ model has even more freedom, allowing a considerable running of the running $d^2 n_s/d \ln k^2$ (to the extent that inflation holds during the observable e -folds). In that case, the spectrum is better fitted when the two parameters r and $d^2 n_s/d \ln k^2$ are non-zero. In Fig. 14, we see that most $n = 4$

potentials have a long and steep tail for $\phi < \phi_*$, with a kink around $\phi_* - 0.4$ (in natural units). This shape generates a significant running on the largest observable scales, while preserving a smaller running on smaller scales. With such a feature in the scalar primordial spectrum at large scales, combined with a non-zero contribution from tensor fluctuations, the best-fit model for $n = 4$ has a temperature spectrum very close to that of the minimal ΛCDM model for $\ell > 40$, but not for smaller multipoles: the amplitude of the Sachs-Wolfe plateau is smaller. This allows the large-scale data points from *Planck* to be fit slightly better. However, the case $r = dn_s/d \ln k = 0$ still lies at the edge of the 95% CL, and the minimum effective χ^2 of this model is smaller than in the $n = 2$ case by only 3.7.

A comparison of the $n = 3$ and $n = 4$ results clearly shows that the process of expanding the inflation potential at various orders and fitting it to the data does not converge (at least not by $n = 4$). Given the $1 - 2\sigma$ preference of *Planck* data for a non-zero running and running of the running, we find that a model-independent reconstruction of the inflaton potential is not possible under the assumptions of this section. In other words, as long as we assume that $V(\phi)$ can be described during and after observable inflation by a polynomial of order 2 or 3, we can put strong bounds on ϵ_V , η_V . But if we introduce more derivatives to describe the observable part of the potential, and keep complete freedom to extrapolate $V(\phi)$ outside this region, the constraints can be easily evaded.

7. Primordial power spectrum reconstruction

In this section we report on a search for features in the primordial power spectrum. In the basic six parameter model studied in the companion *Planck* paper [Planck Collaboration XVI \(2013\)](#), the primordial power spectrum $\mathcal{P}_{\mathcal{R}}(k)$, which includes only the adiabatic mode, is modeled using the power law $\mathcal{P}_{\mathcal{R}}(k) = A_s (k/k_*)^{n_s-1}$, for which the best fit values are $A_s = 2.20 \times 10^{-9}$ and $n_s = 0.9603$ for a pivot scale $k_* = 0.05 \text{ Mpc}^{-1}$. An extension of this parameterization is also considered allowing for a running of the spectral index ($dn_s/d \ln k = -0.013 \pm 0.009$). But in all cases considered it was assumed that the power spectrum is smooth and without bumps, sharp features, or wiggles. In this section we investigate whether any statistically significant evidence for features is present in the data when these assumptions are relaxed. Allowing an arbitrary function for the input power spectrum is not an option because in this case the recovered primordial power spectrum is dominated by small scale noise. Instead we consider here a penalized likelihood approach where a preference for smooth power spectra is imposed. The companion Sect. 8 of this paper pursues a complementary approach where several parametric models for wiggles and features are explored to see whether a statistically significantly better fit can be obtained.

An extensive literature exists on how to search for features in the power spectrum using a wide range of methods. The following papers and the references therein provide a sampling of the literature on non-parametric reconstruction: Richardson-Lucy deconvolution ([Lucy, 1974](#); [Richardson, 1972](#); [Hamann et al., 2010](#); [Shafieloo & Souradeep, 2004, 2008](#)), deconvolution ([Tocchini-Valentini et al., 2005, 2006](#); [Ichiki & Nagata, 2009](#); [Nagata & Yokoyama, 2008, 2009](#)), smoothing splines ([Verde & Peiris, 2008](#); [Peiris & Verde, 2010](#); [Sealfon et al., 2005](#); [Gauthier & Bucher, 2012](#)), linear interpolation ([Hannestad, 2004](#); [Bridle et al., 2003](#)), and Bayesian model selection ([Bridges et al., 2009](#); [Vázquez et al., 2012](#)). The approach pursued here follows [Tocchini-Valentini et al. \(2006\)](#) and [Gauthier & Bucher \(2012\)](#)




Fig. 15. *Planck* primordial power spectrum feature search results. *Top four panels:* The reconstructed power spectrum at four values for the smoothing parameter λ . The red curves indicate the maximum likelihood configuration for the fractional deviation $f(k)$ of the power spectrum relative to a power law fiducial model (with $A_s = 2.20 \times 10^{-9}$ and $n_s = 0.9603$) for the penalized likelihood. The error bars have a width corresponding to the *minimum reconstructible width* (the minimum width for a Gaussian feature so that the mean square deviation of the expectation value of the reconstruction differs by less than 10%), and a vertical extent showing the 1σ and 2σ limits for the fractional deviation averaged over the box. The grey hashed regions at the far left and right show where the fixing prior (i.e., α) sets $f(k) = 0$. The inner grey regions show where the reconstruction bias is so great that the minimum reconstructible width is undefined. Mock features in this region have reverberations over the entire interval. With $\lambda = 10^3$ and $\lambda = 10^4$, we find statistically significant fluctuations around $k \sim 0.1 \text{ Mpc}^{-1}$. *Lower panels:* The 1σ error bars for three combinations of cosmological parameters at the four values of λ . The maximum likelihood value for the fiducial model is indicated by the dashed line for comparison.

most closely. More technical details and extensive tests validating the method can be found in the latter reference.

Let $\mathcal{P}_0(k) = A_s(k/k_*)^{n_s-1}$ be the best fit power spectrum of the six parameter model. We define a general ansatz for the power spectrum in terms of a fractional variation, $f(k)$, relative to this fiducial model, so that

$$\mathcal{P}_{\mathcal{R}}(k) = \mathcal{P}_0(k)[1 + f(k)]. \quad (47)$$

Any features are then described in terms of $f(k)$.

In this analysis we use the *Planck*+WP likelihood supplemented by the following prior, which is added to $-2 \ln \mathcal{L}$:

$$\begin{aligned} \mathbf{f}^T \mathbf{R}(\lambda, \alpha) \mathbf{f} = & \lambda \int d\kappa \left(\frac{\partial^2 f(\kappa)}{\partial \kappa^2} \right)^2 \\ & + \alpha \int_{-\infty}^{\kappa_{\min}} d\kappa f^2(\kappa) + \alpha \int_{\kappa_{\max}}^{+\infty} d\kappa f^2(\kappa). \end{aligned} \quad (48)$$

where $\kappa = \ln k$. The first term penalizes any deviation from a straight line of the function $f(\kappa)$, and the second term fixes the $f(\kappa)$ to zero where there are effectively no constraints from the data. The value for λ controls the smoothness of the reconstruction, but the precise value of α is less important. It must be large enough to force $f(k)$ towards zero when $\kappa < \kappa_{\min}$ and $\kappa > \kappa_{\max}$ but no so large as to render the matrices ill-conditioned. We use $\alpha = 10^4$.

We represent $f(k)$ using a cubic B-spline on a grid of points in k -space uniformly spaced in $\kappa = \ln k$ with step size $\Delta\kappa = 0.025$ and extended from $\kappa = -12.5$ to $\kappa = -0.3$ giving us a total of 485 knots so that $\mathbf{f} = \{f_i\}_{i=1}^{485}$. The density of grid points is sufficiently large so that artefacts around the scale of the knot spacing are suppressed for the values of λ used here. Given the large number of dimensions, it is not practical to explore the likelihood using MCMC methods. However, for the power spectrum parameters, the predicted C_ℓ s are related by a linear transformation given fixed cosmological parameters, allowing us (for fixed cosmological and nuisance parameters) to find the maximum likelihood solution using the Newton-Raphson method.¹² We define

$$\mathcal{M}(\Theta) = \min_{f_i \in [-1,1]} \left\{ -2 \ln \mathcal{L}(\Theta, \mathbf{f}) + \mathbf{f}^T \mathbf{R}(\lambda, \alpha) \mathbf{f} \right\} \quad (49)$$

where the vector Θ represents the non-power spectrum cosmological parameters and foreground nuisance parameters, and we first minimize over f_i using Newton-Raphson iteration and then in an outer loop minimize over Θ using the downhill simplex algorithm. To achieve this, the *Planck* likelihood code was modified to compute the gradient and Hessian of the likelihood with respect to the C_ℓ s.

The cosmological Boltzmann solver *CAMB* was modified to accept the vector of primordial power spectrum knots \mathbf{f} . By default *CAMB* calculates the C_ℓ s for a subset of ℓ and interpolates to obtain the full multipole power spectrum. Instead we calculate C_ℓ s at each ℓ explicitly.

The boundaries κ_{\min} and κ_{\max} defining where $f(\kappa)$ is allowed to differ from zero are chosen to match the range of ℓ constrained by the high- ℓ likelihood. The likelihood includes C_ℓ s between $\ell = 50$ to $\ell = 2500$, which roughly corresponds to $k \in [0.003, 0.2] \text{ Mpc}^{-1}$. The low- ℓ likelihood covers $\ell = 2$ to $\ell = 49$, which roughly corresponds to $k \in [10^{-4}, 0.003] \text{ Mpc}^{-1}$.

¹² Since the *Planck* likelihood uses a quadratic approximation, the maximum likelihood solution can be found by solving a linear system. Newton-Raphson, however, is needed for other CMB likelihoods which include terms beyond quadratic order.




Fig. 17. CMB multipole spectrum residual for the primordial power spectrum test feature. The test feature (*top*) is set to the anomalously large deviation of the primordial power spectrum reconstruction for $\lambda = 10^3$ in the interval $0.1 \text{ Mpc}^{-1} < k < 0.15 \text{ Mpc}^{-1}$, and is zero elsewhere. *Bottom:* The angular spectrum corresponding to this feature. We observe a large dip at $\ell \approx 1800$.

In this range of ℓ , cosmic variance is large, making feature detection difficult. Calculating the gradient and Hessian of a pixel-based low- ℓ likelihood is computationally time consuming. We therefore use the low- ℓ likelihood only to constrain the cosmological parameters. We choose $k_{\min} = 0.005 \text{ Mpc}^{-1}$ and $k_{\max} = 0.3 \text{ Mpc}^{-1}$. Within this k range, variations in the C_ℓ s due to the f_i are too small to affect the overall likelihood through the low- ℓ likelihood. We observed that the difference in the low- ℓ likelihood between the reconstructed $f(k)$ and $f(k) = 0$ is small ($< 1\%$) compared to the difference in the high- ℓ likelihood.

The cosmological parameters τ and A_s are almost completely degenerate for the temperature anisotropy, except at very low ℓ , so we fix τ to its best fit value for the fiducial model. The likelihood contains additional nuisance parameters that model foreground components and beam shapes, as discussed in [Planck Collaboration XV \(2013\)](#). Many of the nuisance parameters, it can be argued, are unlikely to introduce spurious small scale structure because they represent foreground models with a power law and thus smooth angular power spectrum. However some nuisance parameters, in particular those describing beam uncertainties, could conceivably introduce artefacts into the reconstruction. Unfortunately, converging to the correct maximum likelihood reconstruction with all the beam shape parameters included is prohibitively time consuming. Therefore we fix the nuisance parameters to their fiducial best fit values, leaving a more detailed examination of this issue to future work.

We found that simultaneously allowing extra degrees of freedom for small scale structure and variations in the cosmological parameters changes the best fit fiducial model—that is, A_s and n_s —so that the variations with respect to the fiducial model no longer visibly gave the best straight line fit. Therefore we allow A_s and n_s to vary, so that the fiducial model is indeed the best straight line fit through the plotted data points. Detailed investigation showed that neither the priors, nor low- ℓ , nor high- ℓ data play a significant role in determining the best fit fiducial model. This effect is small and within the error bars for A_s and n_s established assuming the fiducial model. To summarize, we maximize




Fig. 16. CMB multipole spectrum residuals for best fit primordial power spectrum reconstruction at $\lambda = 10^3$. The panels show the C_ℓ spectrum residuals (compared to the best fit power law fiducial model represented by the horizontal straight dashed line) for the four auto- and cross-spectra included in the high- ℓ likelihood. Here, $D_\ell = (\ell(\ell + 1)/(2\pi))C_\ell$. The data points have been binned with $\Delta\ell = 31$ and foregrounds subtracted according to the best fit foreground parameters. The solid black line shows the CMB spectrum residual for the maximum likelihood primordial power spectrum reconstruction with smoothing parameter $\lambda = 10^3$.

the likelihood with respect to the control points f_i and the three cosmological parameters h , $\Omega_c h^2$, and $\Omega_b h^2$. We update the fiducial model (A_s and n_s) at each iteration by finding the best fit power law through the current best fit reconstruction.

Once the maximum likelihood solution has been found, the second derivatives about this solution are readily calculated by extracting the relevant matrices for most of the components, and estimating the remaining components using finite differences. The second derivative matrix is used to estimate the error on the reconstructed f_i and the three cosmological parameters h , $\Omega_c h^2$, and $\Omega_b h^2$. Monte Carlo simulations of a fiducial data set with a simplified CMB likelihood including some of the non-Gaussianities¹³ suggest that this approximation of the error is accurate for $\lambda \gtrsim 10^3$.

Fig. 15 summarizes our results, showing the estimated $f(k)$ in bins, and corresponding 1σ and 2σ errors. Errors in k are also shown to represent the minimum reconstructible width evaluated at the middle of each bin. This is the minimum width that a Gaussian feature must have to be reconstructed with a small enough bias such that the mean reconstruction differs by less than 10% rms. The minimum reconstructible width is closely

related to the correlation length, so that the errors between adjacent bins are weakly correlated and the total number of bins represents roughly the effective number of independent degrees of freedom.

While the plots with a significant roughness penalty—that is, with $\lambda = 10^6$ and $\lambda = 10^5$ —do not show any statistically significant evidence for features standing out from the noise of the reconstruction, for a smaller roughness penalty—that is, for $\lambda = 10^4$ and $\lambda = 10^3$ —a nominally statistically significant feature is clearly visible around $k \approx 0.13 \text{ Mpc}^{-1}$. We do not understand the origin of this feature, which may be primordial or may arise as a foreground or other systematic error in the high- ℓ portion of the likelihood. It should be noted that most of the robustness tests described in the likelihood paper assume smooth power spectra. The maximum excursions are locally at 3.2σ and 3.9σ for $\lambda = 10^4$ and $\lambda = 10^3$, respectively. In each of these two cases we correct for the *look elsewhere* effect by calculating the probability that one of the plotted error bars deviates by the same number of or more standard deviations. This calculation is carried out using the covariance matrix of the plotted error bars. We obtain $p = 1.74\%$ and $p = 0.21\%$, which corresponds to 2.4σ and 3.1σ , respectively. Additional simulations were carried out to validate the method by generating mock data according to the fiducial model and measuring the errors of the reconstruction obtained. These investigations confirm the error model. These

¹³ The simplified likelihood $-2 \ln \mathcal{L}_{\text{toy}} = \sum_{\ell=2}^{\ell_{\text{max}}} (2\ell + 1) \left(\frac{C_\ell^{\text{obs}}}{C_\ell + N_\ell} - \ln \frac{C_\ell^{\text{obs}}}{C_\ell + N_\ell} - 1 \right)$ assumes full-sky coverage and isotropic instrument noise.

tests were carried out both with and without test features. It can be argued that foregrounds are unlikely to explain the observed feature because all the foreground models involve smooth power law templates whereas this feature is localized in multipole number. It is important to assess by means of a more extensive set of simulations whether the statistical significance assigned to this result is accurate.

We investigate which CMB angular multipoles correspond to this apparent feature. Fig. 16 shows the C_ℓ residual from the reconstructed power spectrum with the best fit power law power spectrum subtracted together with the data for each frequency map correlation combinations used in the `CamSpec` likelihood. We observe a smooth dip around $\ell \approx 1800$, which is significant compared to the error bars, in particular for the 217 GHz map. To determine whether this dip is in fact responsible for large deviation in the reconstruction, we take the $\lambda = 10^3$ best fit reconstruction and set $f(k) = 0$ everywhere except for $0.1 \text{ Mpc}^{-1} < k < 0.15 \text{ Mpc}^{-1}$ —the region where the large deviation is located—and calculated the corresponding C_ℓ spectrum. Fig. 17 plots the C_ℓ residuals of this test feature, which show a large dip at around $\ell \approx 1800$, thus demonstrating that the dip in the C_ℓ residual of the data centred at $\ell \approx 1800$ is responsible for the large excursions in the primordial power spectrum reconstructions. The features in the temperature power spectrum, particularly the broad dip at $\ell \approx 1800$, cannot be explained at this level by any of the known systematics that have been propagated through the full data analysis pipeline (Planck Collaboration VI, 2013). Nevertheless, we caution the reader that these (and other small) departures from the best fit Λ CDM spectrum may possibly be due to unknown systematic effects, or inaccurate propagation of known systematics, into the final power spectra. This possibility will be investigated intensively over the coming year using the data from the full mission.

8. Parametric searches for primordial power spectrum features

In this section we continue to investigate deviations of the primordial power spectrum from a smooth, featureless function, in this case by testing a set of theoretically motivated models.

8.1. Models and priors

We consider three models describing features in the primordial power spectrum, adding a global oscillation, a localized oscillation, or a cutoff to the large scale power spectrum.

8.1.1. Wiggles model

Due to the exponential growth of the scale factor during inflation, a periodically recurring event in proper time which affects the amplitude of curvature perturbations would produce features that are periodic in $\ln k$. This occurs, for instance, for non-Bunch-Davies initial conditions (Easther et al., 2001; Danielsson, 2002; Martin & Brandenberger, 2003; Bozza et al., 2003), or, e.g., in the axion monodromy model (Silverstein & Westphal, 2008), as a consequence of instanton-induced corrections to the potential (Flauger et al., 2010). In these scenarios the primordial spectrum has an oscillation superimposed on an underlying smooth spectrum.

Here we consider the following parameterization of the primordial spectrum (referred to as the *wiggles* model):

$$\mathcal{P}_{\mathcal{R}}(k) = \mathcal{P}_0(k) \left\{ 1 + \alpha_w \sin \left[\omega \ln \left(\frac{k}{k_*} \right) + \varphi \right] \right\}, \quad (50)$$

with amplitude α_w , frequency ω , and phase φ to quantify the superimposed oscillations. The underlying smooth spectrum has the standard power law form

$$\mathcal{P}_0(k) = A_s \left(\frac{k}{k_*} \right)^{n_s-1}. \quad (51)$$

The prior ranges for the wiggles model parameters are given in Table 10. The obvious prior for the phase φ is uniform over the interval $(0, 2\pi)$. We choose a uniform prior on α_w (a logarithmic prior on α_w introduces considerable dependence of the resulting marginalized posteriors on the lower limit and does not contain the smooth spectrum as a limiting case). The sensitivity to primordial wiggles is limited at high frequencies by the width of the transfer function (Hamann et al., 2008a) and at low frequencies by the requirement of at least one full oscillation in the observable part of the power spectrum. Since *Planck* data are sensitive to wavenumbers over a range of roughly six orders of magnitude, this condition implies $\omega \gtrsim 0.5$. Here we restrict the analysis to $\omega < 100$ and assume a uniform prior. Larger values of the frequency are theoretically possible, e.g., in axion monodromy models (Flauger et al., 2010), but the amplitude of the oscillations in the C_ℓ s will be suppressed with respect to the primordial one. A comprehensive search for higher frequency oscillations is currently underway.

8.1.2. Step-inflation model

If the slow roll of inflation is briefly interrupted, for instance by a phase transition (Starobinsky, 1998; Hunt & Sarkar, 2004), a burst of resonant particle production (Chung et al., 2000), a sudden turn in field space (Achúcarro et al., 2011) or a step in the inflation potential (Adams et al., 2001), a localized oscillatory feature is superimposed on the scalar primordial power spectrum. We adopt the approximate parameterization for such a feature from a step in the potential, introduced by Adshead et al. (2012), with

$$\mathcal{P}_{\mathcal{R}}(k) = \exp \left[\ln \mathcal{P}_0(k) + \frac{\mathcal{A}_f}{3} \frac{k\eta_f/x_d}{\sinh(k\eta_f/x_d)} W'(k\eta_f) \right], \quad (52)$$

where

$$W'(x) = \left(-3 + \frac{9}{x^2} \right) \cos 2x + \left(15 - \frac{9}{x^2} \right) \frac{\sin 2x}{2x}. \quad (53)$$

As in the wiggles model, we choose a uniform prior on the amplitude parameter \mathcal{A}_f (see Table 10). The parameter η_f determines both the frequency of the feature and its location, which is required to lie in the observable range. The damping envelope of the feature is set by the ratio η_f/x_d . We impose uniform priors on the logarithms of η_f and x_d .

8.1.3. Cutoff model

A number of models were suggested to explain the apparent lack of power in the quadrupole and octupole of the *WMAP* temperature power spectrum. Typically in these models, the onset of a slow-roll phase coincides with the time when the largest

Model	Parameter	Prior range
Wiggles	α_w	[0, 0.2]
	ω	[0.5, 100]
	φ	[0, 2 π]
Step-inflation	\mathcal{A}_f	[0, 0.2]
	$\ln(\eta_f/\text{Mpc})$	[0, 12]
	$\ln x_d$	[-1, 5]
Cutoff	$\ln(k_c/\text{Mpc}^{-1})$	[-12, -4]
	λ_c	[0, 15]

Table 10. Prior ranges imposed for the wiggles, step-inflation, and cutoff model parameters.

observable scales exited the horizon during inflation. This naturally suppresses the primordial power spectrum at large scales (see, e.g., [Sinha & Souradeep, 2006](#)). We consider a phenomenological parameterization of a cutoff proposed in [Contaldi et al. \(2003\)](#), given by

$$\mathcal{P}_{\mathcal{R}}(k) = \mathcal{P}_0(k) \left\{ 1 - \exp \left[- \left(\frac{k}{k_c} \right)^{\lambda_c} \right] \right\}. \quad (54)$$

We apply uniform priors on λ_c , which determines the steepness of the cutoff, and on the logarithm of the cutoff scale k_c .

8.2. Method

To achieve the necessary numerical precision for models with features in the primordial spectra, we modify the standard settings of the CAMB numerical code in order to calculate C_ℓ at each ℓ rather than interpolating and refine the grid in wavenumber for the numerical integration. These changes significantly slow down the computation. In the models considered here, the likelihood function has characteristics that make sampling difficult, such as extended plateaus and multiple isolated maxima, rendering the Metropolis-Hastings algorithm inefficient. We therefore use the nested sampling algorithm implemented in the MultiNest add-on ([Feroz & Hobson, 2008](#); [Feroz et al., 2009](#)) to CosmoMC ([Lewis & Bridle, 2002](#)), which is also able to calculate the Bayesian evidence and likelihood profiles.

The signatures of the feature models under investigation are rather unique and generally cannot be mimicked by other parameters, which only lead to smooth variations of the power spectrum (with an exception of highly tuned very low-frequency oscillations that can change the acoustic peak structure). We thus restrict ourselves to varying only the parameters describing the features and keep all remaining cosmological and nuisance parameters fixed to their Λ CDM best-fit values.¹⁴

8.3. Results

For all three models we find that including these additional features improves the quality of the fit with respect to a pure power law spectrum. For the *Planck*+WP data, we show the best-fit primordial curvature power spectra and temperature angular power

¹⁴ An a-posteriori maximization of the likelihood in a narrow parameter range around the best-fit feature model parameters, including a variation of all remaining cosmological and nuisance parameters, shows that the change in the best-fit χ^2_{eff} is merely $\mathcal{O}(1)$ and hence does not affect our conclusions.




Fig. 18. *Top:* Best fit primordial spectrum of curvature perturbations for the power law (black), wiggles (red), step-inflation (green), and cutoff (blue) models. *Bottom:* Residuals of the temperature angular power spectrum. Note that the scale of the vertical axis changes at $\ell = 50$. *Inset:* Zoom on the region of the first acoustic peak.

spectrum residuals in Fig. 18, and report the best fit parameter values in Table 11. Since in all three cases the likelihood functions do not tend to zero in all directions of the respective parameter spaces, the Bayesian quantities (i.e., posterior distributions and Bayes factors) depend considerably on the choice of prior. For this reason, we also quote two prior-independent quantities, the effective χ^2 (i.e., $-2\Delta \ln \mathcal{L}_{\text{max}} = 2 \ln \mathcal{L}_{\text{max}} - 2 \ln \mathcal{L}_{\text{max}}^{\Lambda\text{CDM}}$) and the profile $-2\Delta \ln \mathcal{L}_{\text{max}}$ as a function of selected model parameters plotted alongside the marginalized posteriors in Fig. 19, which illustrates the unconventional shape of the likelihood functions.

For the wiggles model, oscillations around the first acoustic peak and in the $700 < \ell < 900$ range improve the fit to the data, whereas for the best fit step-inflation model the spectrum between the Sachs-Wolfe plateau and the first acoustic peak is fit better. Quantitatively, the cutoff model improves the fit only modestly, with $\Delta\chi^2_{\text{eff}} \approx 3$, but both the wiggles and step-inflation models lead to a larger improvement, with $\Delta\chi^2_{\text{eff}} \approx 10$, at the cost of three new parameters. Already for pre-*Planck* data, improvements of $\Delta\chi^2_{\text{eff}} \approx 10$ have been reported in related analyses (e.g., [Peiris et al. 2003](#); [Martin & Ringeval 2004](#); [Elgarøy et al. 2003](#);

Covi et al. 2006; Meerburg et al. 2012; Benetti et al. 2013; Peiris et al. 2013). Note that in the step-inflation model, the best-fit does not coincide with the maximum of the marginalized posterior probability, indicating that some degree of fine-tuning is necessary to reach the maximum of the likelihood. The maximum of the marginalized posterior at $\ln(\eta_f/\text{Mpc}) \approx 7.2$ actually reproduces the feature at $\ell \approx 20-40$ found previously in *WMAP* data (Peiris et al., 2003). It is also worth pointing out that the secondary peak at $\ln(\eta_f/\text{Mpc}) \approx 4$ corresponds to a feature at multipoles $\ell \approx 1800$, i.e., where the analysis of Sect. 7 found a feature. However, the model is not able to describe this feature very well, yielding only an improvement of $\Delta\chi^2_{\text{eff}} \approx 3$.

Whether or not these findings can be considered statistically significant or arise simply from over-fitting noisy data is not a trivial question (see, for instance, the discussion in Bennett et al. (2011)). From a frequentist statistics point of view, an answer would require the rather involved procedure of repeating the analysis on a large set of simulations. In designing the test statistic, special care would need to be taken in making sure to take into account the *look elsewhere* effect (i.e., the fact that a particular observed anomaly may be very unlikely, whereas the probability of observing *some* anomaly may be much larger). From a Bayesian statistics point of view, it is the Bayesian evidence that can tell us how probable the extended models are, compared to the baseline power-law primordial power spectrum.

For the models considered here, and for our choice of prior probabilities, the Bayesian evidence in fact favours, albeit weakly, the simple power-law spectrum over the more complex models. The reason is that the Bayesian evidence punishes a lack of predictivity in these models: most of the volume of their parameter space is not compatible with the data; a good match to observations is obtained only in a small region. Nonetheless, the observed features remain interesting, since, if they are real, they will also leave traces in other observables. Most notably, in the *E*-mode polarization spectrum, where the signatures of features in the primordial spectrum are actually less washed out than in the temperature spectrum (Mortonson et al., 2009)—the forthcoming *Planck* polarization data will prove very useful in this regard. Additionally, since strong deviations from power law behavior typically indicate non-linear physics, these models generically also predict a non-Gaussian signal which is potentially observable in the bispectrum (Planck Collaboration XXIV, 2013). However, the best-fit wiggles and step-inflation models have oscillations with too high a frequency to be accessible to bispectrum analysis at present.

9. Combined analysis with *Planck* f_{NL} constraints for single field inflation

In the previous sections we have analysed inflationary models with a canonical kinetic term. This led to the tensor-to-scalar consistency condition, requiring $n_t = -r/8$. It is interesting to consider more general classes of inflationary models characterized by a non-standard kinetic term or more general higher derivative operators. An interesting subclass of these models are those in which the Lagrangian is a general function of the scalar inflaton field and its first derivative: $\mathcal{L} = P(\phi, X)$, where $X = -g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi/2$. A more general extension is provided by the so-called effective field theory of inflation (Cheung et al., 2008), which has a richer phenomenology.

We restrict our analysis to the first class of models (Garriga & Mukhanov, 1999; Chen et al., 2007), which includes k -inflation models (Armendáriz-Picón et al. 1999; Garriga &

Model	$-2\Delta \ln \mathcal{L}_{\text{max}}$	$\ln B_{0X}$	Parameter	Best fit value
Wiggles	-9.0	1.5	α_w	0.0294
			ω	28.90
			φ	0.075 π
Step-inflation	-11.7	0.3	\mathcal{A}_f	0.102
			$\ln(\eta_f/\text{Mpc})$	8.214
			$\ln x_d$	4.47
Cutoff	-2.9	0.3	$\ln(k_c/\text{Mpc}^{-1})$	-8.493
			λ_c	0.474

Table 11. Improvement in fit and logarithm of the Bayes factor with respect to power law Λ CDM and best fit parameter values for the wiggles, step-inflation, and cutoff models. The larger $\ln B_{0X}$, the greater the preference for a featureless power law spectrum.

Mukhanov 1999), and Dirac-Born-Infeld (DBI) models introduced in the context of brane inflation (Silverstein & Tong 2004; Alishahiha et al. 2004). In this class of models inflation can take place with a steep potential, or it can be driven by the kinetic term. One of the main features of inflationary models with a non-standard kinetic term is that the inflaton fluctuations can propagate at a sound speed $c_s < 1$. As shown in previous analyses (e.g., Peiris et al., 2007; Powell et al., 2009; Lorenz et al., 2008; Agarwal & Bean, 2009) there are strong degeneracies between the parameters determining the observable power spectra. Constraints on primordial non-Gaussianity can help break this degeneracy, and we show how *Planck*'s combined measurement of the power spectrum and the non-linearity parameter f_{NL} (Planck Collaboration XXIV, 2013) improves constraints on this class of models.

In models with a non-standard kinetic term the sound speed of the inflaton is given by $c_s^2 = P_{,X}/(P_{,X} + 2XP_{,XX})$ (Garriga & Mukhanov, 1999), so that in the canonical models, where $P(\phi, X) = V(\phi) - X$, one finds $c_s = 1$, while in general a non-trivial $c_s < 1$ corresponds to deviations from this standard case. Therefore, in these models, new parameters—like the sound speed and its running—enter the expressions of the inflationary observables. For the running of the sound speed it is useful to define an additional slow-roll parameter

$$s = \frac{\dot{c}_s}{c_s H}. \quad (55)$$

For values of the slow-roll parameters much less than unity, the leading order scalar power spectrum is modified (Garriga & Mukhanov, 1999) to

$$A_s \simeq \frac{1}{8\pi^2 M_{\text{pl}}^2} \frac{H^2}{c_s \epsilon_1}, \quad (56)$$

which is evaluated at $kc_s = aH$. The scalar spectral index gets an additional contribution from the running of the sound speed,

$$n_s - 1 = -2\epsilon_1 - \epsilon_2 - s. \quad (57)$$

The gravitational sector remains unaltered by the non-trivial inflaton sound speed, retaining the same form as for the standard slow-roll models. Therefore the usual consistency relation is modified to $r \simeq -8n_t c_s$ with $n_t = -2\epsilon_1$ as usual (Garriga &




Fig. 19. Marginalized posterior probability (red) and profile $-2\Delta \ln \mathcal{L}_{\max}$ (black) for selected parameters of the wiggles (top), step-inflation (middle), and cutoff model (bottom).

(Mukhanov, 1999). A more accurate relation that we employ in this analysis is given by

$$r = 16\epsilon_1 c_s^{\frac{1+\epsilon_1}{1-\epsilon_1}}. \quad (58)$$

This accounts for the difference in freeze-out between the scalar and tensor perturbations (Peiris et al., 2007; Powell et al., 2009; Lorenz et al., 2008; Agarwal & Bean, 2009), taking place at

sound-horizon crossing (at $kc_s = aH$) for the scalar fluctuations and at horizon-crossing ($k = aH$) for the tensor modes.

Limiting ourselves to the predictions at lowest-order in the slow-roll parameters, there are clearly degeneracies between the parameters ($A_s, c_s, \epsilon_1, \epsilon_2, s$), which make the constraints on the inflationary power spectra observables less stringent in terms of these microscopic parameters. However, for models where the inflaton field has a non-standard kinetic term with $c_s \ll 1$, a high level of primordial non-Gaussianity of the scalar perturbations is generated (see, e.g., Chen et al., 2007). In these models primordial non-Gaussianity is produced by the higher derivative interaction terms that arise when expanding the kinetic part of the Lagrangian, $P(\phi, X)$. The amplitude of the non-Gaussianity, defined by the nonlinearity parameter f_{NL} , receives two dominant contributions, arising from the inflaton interaction terms $(\delta\phi)^3$ and $\delta\phi(\nabla\delta\phi)^2$. Each of them produces non-Gaussianity shapes similar to the so-called “equilateral” type (Babich et al. 2004), i.e., a signal that peaks for equilateral triangles $k_1 = k_2 = k_3$. However, they are sufficiently different that the total signal can be very different from the equilateral one (Senatore et al., 2010). The non-linearity parameter of the second interaction term is $f_{\text{NL}} = (85/324)(1 - c_s^{-2})$, while the other is determined by a second independent amplitude (Chen et al., 2007; Senatore et al., 2010). Constraints on the primordial non-Gaussianity, presented in the companion paper Planck Collaboration XXIV (2013), thus allow us to construct a lower limit of the sound speed c_s . This helps reduce degeneracies in the parameter space of inflationary models with non-standard kinetic terms. In particular, without the limits on the sound speed coming from the constraints on primordial non-Gaussianity, it is not possible to derive an upper limit on the parameter ϵ_1 , because the relation between the tensor-to-scalar ratio and ϵ_1 also involves the sound speed (see, e.g., Eq. 58).

In this paper, we consider three cases. One is a general analysis as described above, where we focus on the simplest case of a constant speed of sound with $s = 0$. From the Planck limits on primordial non-Gaussianity in general single field models of inflation (Planck Collaboration XXIV, 2013), the most conservative constraint on the sound speed is

$$c_s \geq 0.02 \quad (95\% \text{ CL}). \quad (59)$$

In this large parameter space, we assume a uniform prior $0.02 \leq c_s \leq 1$ in Eq. 58 within the HFF formalism described in the Appendix. We show the joint constraints on ϵ_1 and ϵ_2 in Fig. 20. By including the 95% CL constraint on c_s from Eq. 59, Planck+WP constrain $\epsilon_1 < 0.053$. Such constraints can be compared with the restricted case of $c_s = 1$, also shown in Fig. 20, with $\epsilon_1 < 0.008$ at 95% CL.

The other two cases we analyse correspond to two specific models where the inflaton has a non-standard kinetic term. The degeneracy between the different slow-roll parameters is broken because these models specifically predict that $s = 0$, or $s \propto \epsilon_2$. As an example, we first consider the case where the action takes the Dirac-Born-Infeld (DBI) form

$$P(\phi, X) = -f(\phi)^{-1} \sqrt{1 - 2f(\phi)X} + f(\phi)^{-1} - V(\phi). \quad (60)$$

Here $V(\phi)$ is the potential and $f(\phi)$ is the warp factor determined by the geometry of the extra dimensions. For DBI models a stronger bound on c_s is derived (Planck Collaboration XXIV, 2013): $c_s > 0.07$ at 95% CL. With the uniform prior $0.07 < c_s < 1$ and $s = 0$, Planck + WP constrain $\epsilon_1 < 0.042$ at 95% CL.




Fig. 20. Marginalized joint 68% and 95% CL for (ϵ_1, ϵ_2) for *Planck* + WP data, comparing the canonical Lagrangian case with $c_s = 1$, to the case with varying c_s with a uniform prior $0.02 < c_s < 1$, derived from the *Planck* non-Gaussianity measurements.

An important case is $f(\phi) \simeq \lambda/\phi^4$ (for details, see [Silverstein & Tong 2004](#); [Alishahiha et al. 2004](#); [Chen et al. 2007](#), and references therein). There are two possibilities. First, in *ultraviolet* (UV) DBI models, the inflaton field moves under a quadratic potential $V(\phi) \simeq m^2\phi^2/2$ from the UV side of the warped space to the infrared side, with $m \gg M_{\text{pl}}/\sqrt{\lambda}$. It is known that this case is already at odds with observations, if theoretical internal consistency of the model and constraints on power spectra and primordial non-Gaussianity are taken into account ([Baumann & McAllister, 2007](#); [Lidsey & Huston, 2007](#); [Bean et al., 2007](#); [Peiris et al., 2007](#); [Bean et al., 2008](#)). It is therefore interesting to look at the other case, namely *infrared* (IR) DBI models ([Chen 2005b,a](#)) where the inflaton field moves from the IR to the UV side, and the inflaton potential is

$$V(\phi) = V_0 - \frac{1}{2}\beta H^2 \phi^2, \quad (61)$$

with a wide range of values allowed for β in principle, $0.1 < \beta < 10^9$ ([Bean et al., 2008](#)). Here, we focus on a minimal version of the IR DBI models, where string effects are neglected, so that the usual field theory computation of the primordial curvature perturbation holds. For IR DBI models accounting for such effects and a more involved treatment of the dynamics, see [Chen \(2005a,c\)](#); [Bean et al. \(2008\)](#). In this minimal IR DBI model, one finds ([Chen, 2005c](#); [Chen et al., 2007](#)) $c_s \simeq (\beta N_*/3)^{-1}$, $n_s - 1 = -4/N_*$, $dn_s/d \ln k = -4/N_*^2$ (in this model one can verify that $s \simeq 1/N_* \simeq \epsilon_2/3$). Here, primordial non-Gaussianity of the equilateral type is generated with an amplitude $f_{\text{NL}}^{\text{DBI}} = -(35/108)[(\beta^2 N_*^2/9) - 1]$.

If we consider $60 \leq N_* \leq 90$, then the predicted spectral index lies within the range $0.93 \leq n_s \leq 0.96$, which is consistent with the *Planck* measurement of the spectral index at the 3σ level, for $N_* \geq 60$. The constraints on non-Gaussianity give $f_{\text{NL}}^{\text{DBI}} = 11 \pm 69$ at 68% CL ([Planck Collaboration XXIV, 2013](#)). Combining these constraints with the power spectrum constraints, marginalizing over $60 \leq N_* \leq 90$, we obtain

$$\beta \leq 0.7 \quad (95\% \text{ CL}), \quad (62)$$

This strongly limits the allowed parameter space of these models.

As a final example, we consider a class of power-law k -inflation models characterized by the Lagrangian ([Armendáriz-Picón et al., 1999](#))

$$P(\phi, X) = \frac{4}{9} \frac{4 - 3\gamma}{\gamma^2} \frac{1}{\phi^2} (-X + X^2). \quad (63)$$

In this case, for small values of γ one finds: $c_s^2 \simeq \gamma/8$, $\mathcal{P}_R = 2H^2/(3\gamma c_s 8\pi^2 M_{\text{pl}}^2)(k/k_0)^{-3\gamma}$, $n_s - 1 = -3\gamma$. The sound speed is a constant ($s = 0$), with constant γ . The primordial non-Gaussianity in this model has an amplitude $f_{\text{NL}}^{\text{equil}} = -170/(81\gamma)$. Therefore, all the inflationary observables depend essentially on a single parameter, γ . Imposing a prior of $0 < \gamma < 2/3$, from the non-Gaussianity constraint $f_{\text{NL}}^{\text{equil}} = -42 \pm 75$ at 68% CL ([Planck Collaboration XXIV, 2013](#)) we obtain $\gamma \geq 0.05$ at 95% CL. At the same time, our measurement of the spectral index constrains $0.01 \leq \gamma \leq 0.02$ at 95% CL. This class of k -inflation models is therefore excluded by the combined constraints on primordial non-Gaussianity and the power spectrum.

10. Isocurvature modes

10.1. Theoretical background

In this section we explore the constraints imposed by *Planck* on scenarios where the primordial cosmological perturbations were not entirely *adiabatic*. These scenarios also include *isocurvature* modes, possibly correlated among themselves as well as with the adiabatic mode. The adiabatic mode is characterized by the property that at very early times the Universe obeyed a common, spatially-uniform equation of state, and all components initially shared a common velocity field. For the adiabatic mode the density perturbations in the various components (e.g., baryons, CDM, photons, neutrinos) are locked together. Here *baryons* include their accompanying leptons, assumed tightly coupled to maintain charge neutrality.

Isocurvature modes arise from spatial variations in the equation of state or from relative velocities between the components. To analyse how the CMB perturbations were imprinted, it is most convenient to define isocurvature modes at a sufficiently late time such that the relevant components, according to our present best understanding, consisted of baryons, photons, CDM, and neutrinos. Under this hypothesis, in addition to the adiabatic mode there are four possible non-decaying isocurvature modes: the baryon, CDM, and neutrino density isocurvature modes, and the neutrino velocity isocurvature mode (see, e.g., [Bucher et al. \(2000\)](#) for a discussion and further references).

The impact of isocurvature modes on the CMB was first studied in detail by [Peebles & Yu \(1970\)](#) and [Efstathiou & Bond \(1986, 1987\)](#), who contemplated the possibility that isocurvature perturbations rather than adiabatic perturbations were the sole source of cosmological perturbations. [Linde \(1985\)](#), [Polarski & Starobinsky \(1994\)](#), [Linde & Mukhanov \(1997\)](#); and [García-Bellido & Wands \(1996\)](#) pointed out various scenarios in which isocurvature perturbations could be generated within the context of inflation. [Bucher et al. \(2000\)](#) carried out a systematic study of isocurvature modes from a phenomenological perspective, pointing out the relevance of two additional modes (the neutrino density and velocity modes). [Lyth & Wands \(2002\)](#), [Moroi & Takahashi \(2001\)](#) and [Bartolo & Liddle \(2002\)](#) studied an interesting so-called curvaton scenario, in which adiabatic fluctuations from inflation contribute negligibly but quantum fluctuations in a transverse direction modulate the density of decaying

particles, leading to isocurvature perturbations correlated with the adiabatic mode.

Several authors have studied the constraints on isocurvature modes imposed by previous microwave background experiments, including Stompor et al. (1996), Langlois & Riazuelo (2000), Amendola et al. (2002), Peiris et al. (2003), Valiviita & Muhonen (2003), Bucher et al. (2004), Moodley et al. (2004), Beltran et al. (2004), Kurki-Suonio et al. (2005), Dunkley et al. (2005), Bean et al. (2006), Trotta (2007), Keskitalo et al. (2007), and Komatsu et al. (2009). A more complete set of references may be found in Valiviita et al. (2012).

Before proceeding we must define precisely how to characterize these isocurvature modes on superhorizon scales during the post-entropy-generation epoch, during which we assume that the stress-energy content of the Universe can be modelled as a multi-component fluid composed of baryons, CDM particles, photons, and neutrinos. If we assume that the evolution of the Universe during this epoch was adiabatic (used here in the sense of thermodynamically reversible), then the entropy per unit comoving volume is conserved and serves as a useful reference with respect to which the abundances of the other components can be expressed.

The baryon isocurvature mode may be expressed in terms of fractional fluctuations in the baryon-to-entropy ratio, which is conserved on superhorizon scales during this epoch. The CDM and neutrino density isocurvature (NDI) modes may be defined analogously. The neutrino velocity isocurvature (NVI) modes refer to fluctuations in the neutrino velocity relatively to the average bulk velocity of the cosmic fluid. For the CMB, the baryon and CDM isocurvature modes yield almost identical angular spectra, because the deficit of one is balanced by an excess of the other, so we do not consider them separately here. In this way the *primordial* isocurvature modes may be defined as dimensionless stochastic variables I_{CDI} , I_{NDI} , I_{NVI} , like the variable \mathcal{R} describing the adiabatic mode.¹⁵ In this basis, the CDI mode can be seen as an *effective* isocurvature mode, encoding both CDM and baryon isocurvature fluctuations through $I_{\text{CDI}}^{\text{effective}} = I_{\text{CDI}} + (\Omega_b/\Omega_c)I_{\text{BI}}$ (Gordon & Lewis, 2003).

Within this framework, Gaussian fluctuations for the most general cosmological perturbation are described by a 4×4 positive definite matrix-valued power spectrum of the form

$$\mathcal{P}(k) = \begin{pmatrix} \mathcal{P}_{\mathcal{R}} & \mathcal{P}_{\mathcal{R}}(k) & \mathcal{P}_{\mathcal{R}} & I_{\text{CDI}}(k) & \mathcal{P}_{\mathcal{R}} & I_{\text{NDI}}(k) & \mathcal{P}_{\mathcal{R}} & I_{\text{NVI}}(k) \\ \mathcal{P}_{I_{\text{CDI}}\mathcal{R}}(k) & \mathcal{P}_{I_{\text{CDI}}I_{\text{CDI}}}(k) & \mathcal{P}_{I_{\text{CDI}}I_{\text{NDI}}}(k) & \mathcal{P}_{I_{\text{CDI}}I_{\text{NVI}}}(k) \\ \mathcal{P}_{I_{\text{NDI}}\mathcal{R}}(k) & \mathcal{P}_{I_{\text{NDI}}I_{\text{CDI}}}(k) & \mathcal{P}_{I_{\text{NDI}}I_{\text{NDI}}}(k) & \mathcal{P}_{I_{\text{NDI}}I_{\text{NVI}}}(k) \\ \mathcal{P}_{I_{\text{NVI}}\mathcal{R}}(k) & \mathcal{P}_{I_{\text{NVI}}I_{\text{CDI}}}(k) & \mathcal{P}_{I_{\text{NVI}}I_{\text{NDI}}}(k) & \mathcal{P}_{I_{\text{NVI}}I_{\text{NVI}}}(k) \end{pmatrix}. \quad (64)$$

Following the conventions used in CAMB (Lewis & Bridle, 2002; Lewis, 2011) and CLASS (Lesgourgues, 2011; Blas et al., 2011), the *primordial* isocurvature modes are normalized as follows in the synchronous gauge: for the CDI mode, $\mathcal{P}_{II}(k)$ is the primordial power spectrum of the density contrast difference $\delta\rho_{\text{CDM}}/\rho_{\text{CDM}}$; for the NDI mode it is that of $\delta\rho_{\nu}/\rho_{\nu}$; and for the NVI mode, that of the neutrino velocity v_{ν} times $4/3$ ¹⁶.

If isocurvature modes are present, the most plausible mechanism for exciting them involves inflation with a multi-component inflaton field. To have an interesting spectrum on the large scales probed by the CMB, isocurvature modes require

¹⁵ The symbol \mathcal{S} is sometimes used in the literature to denote the isocurvature modes, also known as *entropy* perturbations. To prevent confusion we avoid this terminology because isocurvature modes are unrelated to any notion of thermodynamic entropy.

¹⁶ or in other words, of the neutrino perturbation dipole, $F_{\nu 1} = 4\theta_{\nu}/(3k)$ in the notation of Ma & Bertschinger (1995).




Fig. 21. C_{TT} anisotropy shape for the three isocurvature modes. *Top:* The shapes of the CDM isocurvature mode, neutrino density isocurvature mode and neutrino velocity isocurvature mode are shown together with the adiabatic mode. The modes have the same amplitude parameters ($\mathcal{P}_{\mathcal{R}\mathcal{R}}$ for the adiabatic mode and \mathcal{P}_{II} for each isocurvature mode). *Bottom:* the narrower multipole range illustrates the relative phases of the acoustic oscillations for these modes.

long-range correlations, and at present inflation with a multi-component inflaton provides a well-motivated scenario for establishing such correlations. Inflation with a single component scalar field can excite only the adiabatic mode. In models of inflation with light (compared to the Hubble expansion rate) transverse directions, the scalar field along these transverse directions becomes disordered in a way described by an approximately scale-invariant spectrum. If the inflaton has N light components, there are $(N-1)$ potential isocurvature modes during inflation. Whether or not the fluctuations along these transverse directions are subsequently transformed into the late-time isocurvature modes described above depends on the details of what happens after inflation, as described more formally below.

As explained for example in Gordon et al. (2001), Groot Nibbelink & van Tent (2000, 2002) and Byrnes & Wands (2006), for inflationary models where the inflaton follows a curved trajectory, correlations are generically established between the isocurvature and curvature degrees of freedom. To lowest order in the slow-roll approximation, this leads to a situa-

tion where the adiabatic perturbation is the sum of several components each of differing spectral index.

The post-inflationary evolution determines how the isocurvature fluctuations generated during inflation transmute into the three specific isocurvature modes studied here. Little is known about the details of what happens during the epoch of entropy generation, but to linear order we may express how the fields \mathcal{R}_{inf} (i.e., the curvature perturbation at the end of inflation) and the transverse components of the inflaton field $\sigma_1, \dots, \sigma_{N-1}$ (i.e., the components orthogonal to the slow-roll direction) transform into curvature perturbations and late-time isocurvature modes at the end of the epoch of entropy generation as the following linear transformation:

$$\begin{pmatrix} \mathcal{R}_{\text{out}} \\ \mathcal{I}_a \end{pmatrix} = \begin{pmatrix} 1 & \Sigma_A \\ \mathbf{0} & M_{aA} \end{pmatrix} \begin{pmatrix} \mathcal{R}_{\text{inf}} \\ \sigma_A \end{pmatrix}, \quad (65)$$

where $a = 1, \dots, 4 = \text{BI, CDI, NDI, NVI}$ and $A = 1, \dots, (N-1)$ labels the transverse components of the N component inflaton field. Physically, the fluctuations along the transverse directions modulate particle production during the epoch of entropy generation.

The neutrino density isocurvature can be excited in much the same way as the CDM and baryon isocurvature mode because at least within the standard electroweak model, in which there are no leptonic flavour changing processes, L_e , L_μ , and L_τ are separately conserved. Known non-perturbative processes such as the sphaleron can turn trade lepton and baryon asymmetries with each other and alter flavour asymmetries, but they cannot erase such asymmetries altogether. Plausible models generating the neutrino density mode are therefore possible (Bucher et al., 2000; Gordon & Malik, 2004), but as for the neutrino velocity isocurvature mode, to date no plausible generation mechanism has been put forth.

This extension of the adiabatic ΛCDM model to non-adiabatic initial conditions represents an important test of inflation. Single field inflation can produce only adiabatic perturbations, since exciting isocurvature perturbations requires additional degrees of freedom during inflation. Therefore a detection of primordial isocurvature perturbations would point to more complicated models of inflation.

10.2. Adiabatic with one isocurvature mode and free spectral indices

In this paper we investigate three of the four possible isocurvature modes of the ΛCDM scenario, since the baryon and CDM isocurvature perturbations are indistinguishable in the CMB angular power spectra. The CDM, neutrino density, and neutrino velocity isocurvature perturbations lead to different power spectra for CMB anisotropies, as shown in Fig. 21. We limit ourselves to studying one isocurvature mode at a time, in the presence of a curvature perturbation. More general combinations with two or three isocurvature modes may be contemplated, but without the *Planck* high frequency polarization likelihood, it is difficult to constrain this scenario, so we postpone a discussion of this case to the next release.

Theoretically, one expects the power spectra of the isocurvature modes and their correlations to exhibit near but not necessarily exact scale invariance. As a general test of adiabaticity, it is nevertheless interesting to compare a more general model to the *Planck* data, assuming that the adiabatic, isocurvature, and cross-correlation spectra obey a power law with free spectral indices. Blue values of the spectral indices are particularly interesting from the point of view of testing adiabaticity because

the acoustic peaks arising from two of the isocurvature modes are out of phase with the adiabatic peaks by roughly $(\pi/2)$ near the first acoustic peak. This is not true for the neutrino velocity isocurvature mode however.

In the literature, models with one isocurvature and the adiabatic mode (possibly correlated) are often parameterized by specifying the 2×2 correlation at a certain pivot scale k_0 through the components $\mathcal{P}_{\mathcal{R}\mathcal{R}}, \mathcal{P}_{\mathcal{R}\mathcal{I}}, \mathcal{P}_{\mathcal{I}\mathcal{I}}$ along with their respective spectral indices $n_{\mathcal{R}\mathcal{R}}, n_{\mathcal{R}\mathcal{I}}, n_{\mathcal{I}\mathcal{I}}$ (e.g., Amendola et al., 2002; Beltran et al., 2004). We do not follow this approach here because in the absence of a statistically significant detection, the posterior distributions for the spectral indices are difficult to interpret and sensitive to how the prior is chosen. We instead adopt a parameterization where \mathcal{P}_{ab} is specified at two scales $k = k_1$ and $k = k_2$ and interpolated geometrically according to the relation¹⁷

$$\mathcal{P}_{\text{ab}}(k) = \exp \left[\left(\frac{\ln(k) - \ln(k_2)}{\ln(k_1) - \ln(k_2)} \right) \ln(\mathcal{P}_{\text{ab}}^{(1)}) + \left(\frac{\ln(k) - \ln(k_1)}{\ln(k_2) - \ln(k_1)} \right) \ln(\mathcal{P}_{\text{ab}}^{(2)}) \right], \quad (66)$$

where $a, b = \mathcal{I}, \mathcal{R}$ and $\mathcal{I} = \mathcal{I}_{\text{CDI}}, \mathcal{I}_{\text{NDI}}, \text{ or } \mathcal{I}_{\text{NVI}}$. We set $k_1 = 2 \times 10^{-3} \text{ Mpc}^{-1}$ and $k_2 = 0.1 \text{ Mpc}^{-1}$, so that $[k_1, k_2]$ spans most of the range in k constrained by *Planck* data. A uniform prior for the components $\mathcal{P}_{\mathcal{R}\mathcal{R}}^{(1)}, \mathcal{P}_{\mathcal{I}\mathcal{I}}^{(1)}, \mathcal{P}_{\mathcal{R}\mathcal{I}}^{(1)}, \mathcal{P}_{\mathcal{R}\mathcal{R}}^{(2)}, \mathcal{P}_{\mathcal{I}\mathcal{I}}^{(2)}, \mathcal{P}_{\mathcal{R}\mathcal{I}}^{(2)}$ is assumed, where auto-correlation amplitudes $\mathcal{P}_{\mathcal{R}\mathcal{R}}^{(1)}, \mathcal{P}_{\mathcal{I}\mathcal{I}}^{(1)}, \mathcal{P}_{\mathcal{R}\mathcal{R}}^{(2)}, \mathcal{P}_{\mathcal{I}\mathcal{I}}^{(2)}$ are positive, although the cross-correlation amplitudes $\mathcal{P}_{\mathcal{R}\mathcal{I}}^{(1)}, \mathcal{P}_{\mathcal{R}\mathcal{I}}^{(2)}$ may take both signs subject to the constraints

$$(\mathcal{P}_{\mathcal{R}\mathcal{I}}^{(1)})^2 < \mathcal{P}_{\mathcal{R}\mathcal{R}}^{(1)} \mathcal{P}_{\mathcal{I}\mathcal{I}}^{(1)}, \quad (\mathcal{P}_{\mathcal{R}\mathcal{I}}^{(2)})^2 < \mathcal{P}_{\mathcal{R}\mathcal{R}}^{(2)} \mathcal{P}_{\mathcal{I}\mathcal{I}}^{(2)}, \quad (67)$$

to ensure positive definiteness. For the logarithm of the off-diagonal elements in Eq. 66 to be real, we must have $\mathcal{P}_{\mathcal{R}\mathcal{I}}^{(1)}, \mathcal{P}_{\mathcal{R}\mathcal{I}}^{(2)} > 0$. This Ansatz can be trivially modified to admit the case $\mathcal{P}_{\mathcal{R}\mathcal{I}}^{(1)}, \mathcal{P}_{\mathcal{R}\mathcal{I}}^{(2)} < 0$ by inserting appropriate minus signs, but this parameterization does not allow the case where the sign of the correlation changes. In practice we deal with this by assuming a uniform prior not on $\mathcal{P}_{\mathcal{R}\mathcal{I}}^{(2)}$, but on its absolute value, and then we impose $\mathcal{P}_{\mathcal{R}\mathcal{I}}^{(2)} = \text{sign}(\mathcal{P}_{\mathcal{R}\mathcal{I}}^{(1)}) \times |\mathcal{P}_{\mathcal{R}\mathcal{I}}^{(2)}|$. The constraints in Eq. 67 ensure that $\det(\mathcal{P}_{\text{ab}}(k))$ is positive definite within the interval $[k_1, k_2]$, but generically positive definiteness is violated sufficiently far outside this interval, either for very small or very large k . Where this happens we reduce the magnitude of $\mathcal{P}_{\mathcal{R}\mathcal{I}}$ so that there is either total correlation or anti-correlation. The kinks thus introduced lie outside the range $[k_1, k_2]$. Within this range, the spectral indices $n_{\mathcal{R}\mathcal{R}}, n_{\mathcal{R}\mathcal{I}}, n_{\mathcal{I}\mathcal{I}}$ are scale-independent. Finally, our sign conventions are such that positive values for $\mathcal{P}_{\mathcal{R}\mathcal{I}}^{(1,2)}$ correspond to a positive contribution of the cross-correlation term to the Sachs-Wolfe component of the total temperature spectrum.

When the constraining power of the data is weak, a crucial question is to what extent the posterior distribution results from the data rather than from the prior distribution. The parameterization above is not the only one that could have been adopted, and other possible priors are typically related by a non-constant Jacobian. For each model, we indicate the log-likelihood for the best fit model, in order to allow model comparison.

The *Planck*+WP results for the three isocurvature modes using this two-scale parameterization are shown in Fig. 22 and

¹⁷ Although the models spanned by the one-scale and two-scale parameterizations are the same, the priors for these parameterizations are related by a non-constant Jacobian and therefore do not coincide.

included in the summary Table 12. The power spectra $\mathcal{P}_{\mathcal{R}\mathcal{R}}(k)$, $\mathcal{P}_{\mathcal{R}\mathcal{I}}(k)$, and $\mathcal{P}_{\mathcal{I}\mathcal{I}}(k)$ are normalized according to the *primordial* values of the fields $\mathcal{R}(\mathbf{x})$ and $\mathcal{I}(\mathbf{x})$ defined above. It is interesting to consider how much isocurvature is allowed expressed as a fraction of the power in three bands spanning the CMB temperature spectrum observed by *Planck*. To this end, we define the following derived quantities

$$\alpha_{\mathcal{R}\mathcal{R}}(\ell_{\min}, \ell_{\max}) = \frac{(\Delta T)_{\mathcal{R}\mathcal{R}}^2(\ell_{\min}, \ell_{\max})}{(\Delta T)_{\text{tot}}^2(\ell_{\min}, \ell_{\max})}, \quad (68)$$

$$\alpha_{\mathcal{I}\mathcal{I}}(\ell_{\min}, \ell_{\max}) = \frac{(\Delta T)_{\mathcal{I}\mathcal{I}}^2(\ell_{\min}, \ell_{\max})}{(\Delta T)_{\text{tot}}^2(\ell_{\min}, \ell_{\max})}, \quad (69)$$

$$\alpha_{\mathcal{R}\mathcal{I}}(\ell_{\min}, \ell_{\max}) = \frac{(\Delta T)_{\mathcal{R}\mathcal{I}}^2(\ell_{\min}, \ell_{\max})}{(\Delta T)_{\text{tot}}^2(\ell_{\min}, \ell_{\max})}, \quad (70)$$

where

$$(\Delta T)_{\mathcal{X}}^2(\ell_{\min}, \ell_{\max}) = \sum_{\ell=\ell_{\min}}^{\ell_{\max}} (2\ell+1) C_{\mathcal{X},\ell}^{TT}. \quad (71)$$

The 95% confidence limits from the one-dimensional posterior distributions for these fractional contributions in the full range $(\ell_{\min}, \ell_{\max}) = (2, 2500)$ are shown in Table 12. The range of allowed values for $\alpha_{\mathcal{R}\mathcal{R}}(2, 2500)$ is a measure of the adiabaticity of fluctuations in the CMB. The posterior distributions of the fractions $\alpha_{\mathcal{I}\mathcal{I}}$, $\alpha_{\mathcal{R}\mathcal{I}}$ in three multipole ranges are shown in Fig. 23. We also report the primordial isocurvature fraction, defined as

$$\beta_{\text{iso}}(k) = \frac{\mathcal{P}_{\mathcal{I}\mathcal{I}}(k)}{\mathcal{P}_{\mathcal{R}\mathcal{R}}(k) + \mathcal{P}_{\mathcal{I}\mathcal{I}}(k)} \quad (72)$$

at three values of k . Table 12 also shows the effective $\chi^2 = -2 \ln \mathcal{L}_{\text{max}}$ for all models, compared to the minimal six-parameter Λ CDM model. In Fig. 24 we show the ratio of temperature spectra for the best-fit mixed model to the adiabatic model.




Fig. 22. Two dimensional distributions for power in isocurvature modes, using *Planck*+WP data.




Fig. 23. Fractional contribution of isocurvature modes to the spectrum. We show the distributions $\alpha_{\mathcal{I}\mathcal{I}}(2, 20)$, $\alpha_{\mathcal{R}\mathcal{I}}(2, 20)$, $\alpha_{\mathcal{I}\mathcal{I}}(21, 200)$, $\alpha_{\mathcal{R}\mathcal{I}}(21, 200)$, $\alpha_{\mathcal{I}\mathcal{I}}(201, 2500)$, $\alpha_{\mathcal{R}\mathcal{I}}(201, 2500)$ defined in Eq. 70 for the CDI, NDI, NVI modes, constrained with *Planck*+WP data.




Fig. 24. Temperature spectrum of best-fit models with a mixture of adiabatic and isocurvature modes. *Top:* spectrum of the best-fit mixed models relative to that of the pure adiabatic model. *Bottom:* zoom on the Sachs-Wolfe plateau of the best-fit temperature spectrum $D_\ell = [\ell(\ell+1)/2\pi] C_\ell^{TT}$, for each of the three cases plus the pure adiabatic model, shown together with *Planck* low- ℓ data points.

The results for $\alpha_{\mathcal{R}\mathcal{R}}(2, 2500)$ show that the nonadiabatic contribution to the temperature variance can be as large as 7% (9%, 5%) in the CDI (NDI, NVI) model (95% CL). These results are driven by the fact that on large scales, for $l \leq 40$, the *Planck* data points on average have a slightly smaller amplitude than the best-fitting Λ CDM model. Hence the data prefer a signifi-

Model	$\beta_{\text{iso}}(k_{\text{low}})$	$\beta_{\text{iso}}(k_{\text{mid}})$	$\beta_{\text{iso}}(k_{\text{high}})$	$\alpha_{\mathcal{R}\mathcal{R}}^{(2,2500)}$	$\alpha_{\mathcal{I}\mathcal{I}}^{(2,2500)}$	$\alpha_{\mathcal{R}\mathcal{I}}^{(2,2500)}$	Δn	$-2\Delta \ln \mathcal{L}_{\text{max}}$
General model:								
CDM isocurvature	0.075	0.39	0.60	[0.98:1.07]	0.039	[-0.093:0.014]	4	-4.6
ND isocurvature	0.27	0.27	0.32	[0.99:1.09]	0.093	[-0.18:0]	4	-4.2
NV isocurvature	0.18	0.14	0.17	[0.96:1.05]	0.068	[-0.090:0.026]	4	-2.5
Special CDM isocurvature cases:								
Uncorrelated, $n_{\mathcal{I}\mathcal{I}} = 1$, (“axion”)	0.036	0.039	0.040	[0.98:1]	0.016	–	1	0
Fully correlated, $n_{\mathcal{I}\mathcal{I}} = n_{\mathcal{R}\mathcal{R}}$, (“curvaton”)	0.0025	0.0025	0.0025	[0.97:1]	0.0011	[0:0.028]	1	0
Fully anti-correlated, $n_{\mathcal{I}\mathcal{I}} = n_{\mathcal{R}\mathcal{R}}$	0.0087	0.0087	0.0087	[1:1.06]	0.0046	[-0.067:0]	1	-1.3

Table 12. Isocurvature mode constraints. For each model, we report the 95% CL upper bound on the fractional primordial contribution of isocurvature modes at three co-moving wavenumbers ($k_{\text{low}} = 0.002 \text{ Mpc}^{-1}$, $k_{\text{mid}} = 0.05 \text{ Mpc}^{-1}$, and $k_{\text{high}} = 0.10 \text{ Mpc}^{-1}$), and the 95% CL bounds on the fractional contribution $\alpha_{\mathcal{I}\mathcal{I}}$ and $\alpha_{\mathcal{R}\mathcal{I}}$ to the total CMB temperature anisotropy in the range $2 \leq \ell \leq 2500$. We also report $-2\Delta \ln \mathcal{L}_{\text{max}}$ for the best-fitting model in each case, relative to the best-fit 6-parameter Λ CDM model, with the number of additional parameters Δn . In the Gaussian approximation, $-2\Delta \ln \mathcal{L}_{\text{max}}$ corresponds to $\Delta\chi^2$. The general models have six parameters that specify the primordial correlation matrix at two scales k_1 and k_2 , thus allowing all spectral indices to vary (so, four parameters more than the pure adiabatic model).

cant amount of anticorrelated isocurvature modes, leading to a reduction of amplitude of the Sachs-Wolfe plateau and to a decrease of the effective χ^2 by up to 4.6¹⁸. This situation explains the rather loose bounds on the derived parameter $\alpha_{\mathcal{I}\mathcal{I}}(2, 20)$, as shown in Fig. 23.

A comparison of $\mathcal{P}_{\mathcal{I}\mathcal{I}}^1$ and $\mathcal{P}_{\mathcal{I}\mathcal{I}}^2$ shows that best-fitting models have an isocurvature spectral index $n_{\mathcal{I}\mathcal{I}}$ close to 1.7 for CDI, 1.1 for NDI and 1.0 for NVI modes.

For CDI and NDI, the amplitude of acoustic peaks quickly decreases with increasing ℓ , so the constraints are entirely driven by small ℓ 's. Since the same value of the primordial amplitude $\mathcal{P}_{\mathcal{I}\mathcal{I}}^{(1)}$ leads to different plateau amplitudes for the two isocurvature models (see Fig. 21), the bounds on $\mathcal{P}_{\mathcal{I}\mathcal{I}}^{(1)}$ and $\mathcal{P}_{\mathcal{R}\mathcal{I}}^{(1)}$ are consistently stronger for CDI than for NDI. For NVI, the acoustic peak amplitude is larger than the plateau amplitude. In NVI models, the data cannot allow for a too large amplitude of correlated isocurvature modes at small ℓ , because the total spectrum would be distorted at larger ℓ . This possibility is strongly disfavoured by the data, which is consistent with the peak location predicted by a pure adiabatic model. Hence in the NVI case we obtain slightly stronger bounds and a smaller reduction of the effective χ^2 .

The fact that the data prefer models with a significant contribution from CDI or NDI modes should be interpreted with care. The detection of a shift in the phase of acoustic oscillations would bring unambiguous evidence in favour of isocurvature modes. With *Planck* data, we are not in this situation. The evidence is driven by a small deficit of amplitude in the Sachs-Wolfe plateau, that could have several different possible explanations (such as a deficit in the large-scale primordial power spectrum, as already seen in the previous sections). However, multi-field inflationary scenarios can produce the mixture of curvature and isocurvature fluctuations which we have found to provide a good fit to the *Planck* data.

¹⁸ For the three general models, the posterior distribution is actually multimodal. Here we are referring to models contributing to the main peak in the posterior, with the highest maximum likelihood. There is another peak with a smaller maximum likelihood, appearing in Fig. 23 as a small bump for positive values of the cross-correlation amplitude. In this paper, we do not carry out a separate investigation for models contributing to this secondary peak.

10.3. Special cases

The six-parameter models of the previous subsection, including one isocurvature mode and the adiabatic mode, make no assumptions about the spectral indices of each mode, or the degree of correlation between the isocurvature mode and the adiabatic mode. This leads to a large number of additional degrees of freedom. There are either theoretical and phenomenological motivations for limiting the values of the parameters to certain values, leading to specific cases with just one more degree of freedom with respect to the adiabatic case. The results are reported in Table 12, for uncorrelated perturbations with $n_{\mathcal{I}\mathcal{I}} = 1$, and fully correlated or anti-correlated perturbations with $n_{\mathcal{I}\mathcal{I}} = n_{\mathcal{R}\mathcal{R}}$. As for the general case, anti-correlated isocurvature perturbations slightly improve the fit to *Planck* data. In the following we consider the implications of our results for two important scenarios, the axion and curvaton scenarios.

10.3.1. Constraints on axion isocurvature

The axion field was proposed to solve the strong CP problem and constitutes a well-motivated dark matter candidate (see, e.g., Preskill et al. (1983), Turner (1990), Peccei (2008), Sikivie (2008), Raffelt (2008), and Kim & Carosi (2010)). The axion is the Goldstone boson of the broken Peccei-Quinn (PQ) symmetry. Under certain assumptions, the axion field may induce significant isocurvature perturbations (Turner et al., 1983; Axenides et al., 1983; Steinhardt & Turner, 1983; Linde, 1984, 1985; Seckel & Turner, 1985; Kofman, 1986; Lyth, 1990; Linde & Lyth, 1990; Turner & Wilczek, 1991; Linde, 1991; Lyth, 1992). If inflation takes place after PQ symmetry breaking, the quantum fluctuations of the inflaton are responsible for primordial curvature perturbations, while those of the axion field generate primordial entropy perturbations. After the QCD transition, when one of the vacua becomes preferred giving the axion field a mass, the axions behave as cold dark matter. This way of producing axionic dark matter is called the misalignment angle mechanism. In such a scenario, the CMB anisotropy may include significant power from CDM isocurvature fluctuations. In that case, the fraction $\beta_{\text{iso}} \equiv \mathcal{P}_{\mathcal{I}\mathcal{I}}/(\mathcal{P}_{\mathcal{R}\mathcal{R}} + \mathcal{P}_{\mathcal{I}\mathcal{I}})$ of CDM isocurvature modes is related to the energy scale of inflation, H_{inf} , through (Lyth, 1990;

Beltran et al., 2007; Bae et al., 2008; Hamann et al., 2009)

$$H_{\text{inf}} = \frac{0.96 \times 10^7 \text{ GeV}}{R_a} \left(\frac{\beta_{\text{iso}}}{0.04} \right)^{1/2} \left(\frac{\Omega_a}{0.120} \right)^{1/2} \left(\frac{f_a}{10^{11} \text{ GeV}} \right)^{0.408}, \quad (73)$$

where Ω_a is the relic axion density, R_a the fraction of CDM consisting of axions, and f_a the PQ symmetry breaking scale. In this model, CDM isocurvature perturbations should be totally uncorrelated with adiabatic perturbations, and have a spectral index n_{II} very close to one, since in the first-order slow-roll approximation the index reads $(1 - 2\epsilon_V)$. Since the sensitivity of the data to n_{II} is very limited (Beltran et al., 2007), we assume for simplicity that $n_{II} = 1$.

Within the general parametrization presented in Eq. 66, we can select the axion case by imposing $\mathcal{P}_{\mathcal{RI}}^{(1,2)} = 0$, as well as the condition

$$\mathcal{P}_{II}^{(2)} = \mathcal{P}_{II}^{(1)}, \quad (74)$$

corresponding to $n_{II} = 1$. We therefore have three independent parameters, $\mathcal{P}_{\mathcal{RR}}^{(1,2)}$, $\mathcal{P}_{II}^{(1)}$, and we sample these parameters with uniform prior distributions. The fraction $\beta_{\text{iso}}(k_*)$ with $k_* = 0.05 \text{ Mpc}^{-1}$ is then a derived parameter. Since the data constrain $\beta_{\text{iso}} \ll 1$, the relation between β_{iso} and $\mathcal{P}_{II}^{(1)}$ is nearly linear, so the primordial isocurvature fraction is sampled with a close-to-uniform prior.

Constraints on this model are shown in Table 12. We find

$$\beta_{\text{iso}} < 0.039 \quad (95\% \text{ CL, Planck+WP}), \quad (75)$$

at the scale $k_{\text{mid}} = 0.05 \text{ Mpc}^{-1}$, with a best-fit value of zero. Hence there is no evidence for axion-generated isocurvature perturbations. This limit significantly improves the latest CMB bounds: at the scale $k = 0.002 \text{ Mpc}^{-1}$, our result reads $\beta_{\text{iso}} < 0.036$, to be compared with $\beta_{\text{iso}} < 0.15$ for *WMAP* 9-year alone, or $\beta_{\text{iso}} < 0.061$ for *WMAP*+ACT+SPT at 95% CL (Hinshaw et al., 2012b). This bound can be used to exclude regions in the parameter space composed of f_a , R_a , and the energy scale of inflation, but cannot be used to obtain a model-independent bound on f_a . However, if we assume (i) that the PQ symmetry is broken during inflation, (ii) that it is not restored by the quantum fluctuations of the inflaton (which imposes $H_{\text{inf}}/(2\pi) < f_a$), nor by thermal fluctuations in case of a very efficient reheating stage, and (iii) that all of CDM consists in axions produced by the misalignment angle, then we can derive an upper bound on the energy scale of inflation as

$$H_{\text{inf}} \leq 0.87 \times 10^7 \text{ GeV} \left(\frac{f_a}{10^{11} \text{ GeV}} \right)^{0.408} \quad (95\% \text{ CL}). \quad (76)$$

10.3.2. Constraints on the curvaton scenario

In the simplest one-field inflationary models curvature perturbations arise from quantum fluctuations in the inflaton field, but this is not the only way to generate them. Entropy perturbations may seed curvature perturbations outside the horizon (Polarski & Starobinsky, 1994; Gordon et al., 2001), so it is possible that a significant component of the observed adiabatic mode could be strongly correlated with an isocurvature mode. This happens for instance in the curvaton scenario (Mollerach, 1990; Enqvist & Sloth, 2002; Moroi & Takahashi, 2001; Lyth & Wands, 2002; Lyth et al., 2003; Gordon & Lewis, 2003). The curvaton is an extra light scalar field acquiring a spectrum of fluctuations on cosmological scales during inflation. Depending on its density evolution and decay history, this field could be responsible for

part of the observed adiabatic perturbations, or all of them, or for a mixture of correlated adiabatic and isocurvature perturbations.

We focus here on the simplest viable version of this scenario, in which the curvaton decays into CDM particles while contributing to a non-negligible fraction

$$r_D = \frac{3\rho_{\text{curvaton}}}{3\rho_{\text{curvaton}} + 4\rho_{\text{radiation}}} \quad (77)$$

of the total energy density of the Universe. If the curvaton dominates at decay time ($r_D = 1$), its primordial fluctuations seed curvature perturbations equivalent to a pure adiabatic mode. If $r_D < 1$, curvaton fluctuations are only partially converted into adiabatic perturbations, while CDM particles carry CDI perturbations, that are fully correlated with the adiabatic ones since they share a common origin. We recall that with our conventions, “fully correlated” means that the cross-correlation term brings a positive contribution to the Sachs-Wolfe component of the total temperature spectrum; some authors define the correlation with the opposite sign and call this case “fully anti-correlated” (e.g., Komatsu et al., 2011; Hinshaw et al., 2012b). In this model, the CDI fraction is related to r_D by (Gordon & Lewis, 2003)

$$\frac{\mathcal{I}_{\text{CDI}}}{\mathcal{R}} = \frac{3(1 - r_D)}{r_D}. \quad (78)$$

In our notation this is equivalent to

$$\beta_{\text{iso}} = \frac{9(1 - r_D)^2}{r_D^2 + 9(1 - r_D)^2}. \quad (79)$$

Within the general parametrization presented in Eq. 66, we can satisfy this case by imposing

$$\frac{\mathcal{P}_{\mathcal{RI}}^{(1)}}{\sqrt{\mathcal{P}_{\mathcal{RR}}^{(1)}\mathcal{P}_{II}^{(1)}}} = \frac{\mathcal{P}_{\mathcal{RI}}^{(2)}}{\sqrt{\mathcal{P}_{\mathcal{RR}}^{(2)}\mathcal{P}_{II}^{(2)}}} = 1, \quad (80)$$

together with the condition

$$\mathcal{P}_{II}^{(2)} = \frac{\mathcal{P}_{II}^{(1)}\mathcal{P}_{\mathcal{RR}}^{(2)}}{\mathcal{P}_{\mathcal{RR}}^{(1)}}, \quad (81)$$

corresponding to $n_{II} = n_{\mathcal{RR}}$. Like in the axion case, this results in three independent parameters $\mathcal{P}_{\mathcal{RR}}^{(1,2)}$, $\mathcal{P}_{II}^{(1)}$ which we sample with uniform priors. The constraints for this model are shown in Table 12. The best-fit model is still the pure adiabatic case, and the upper bound

$$\beta_{\text{iso}} < 0.0025 \quad (95\% \text{ CL, Planck+WP}) \quad (82)$$

is scale independent, since the adiabatic and isocurvature tilts are assumed to be equal. This is a significant improvement over the *WMAP* 9-year bounds, $\beta_{\text{iso}} < 0.012$ for *WMAP* alone, or $\beta_{\text{iso}} < 0.0076$ for *WMAP*+ACT+SPT at 95% CL (Hinshaw et al., 2012b). We conclude that in this scenario, the curvaton should decay when it dominates the energy density of the Universe, with $r_D > 0.983$.

The non-linearity parameter in the curvaton model studied here is (Bartolo et al., 2004c,b)

$$f_{\text{NL}}^{\text{local}} = \frac{5}{4r_D} - \frac{5}{3} - \frac{5r_D}{6}, \quad (83)$$

assuming a quadratic potential for the curvaton field (Sasaki et al., 2006). In the pure adiabatic case ($r_D = 1$) this leads to

$f_{\text{NL}}^{\text{local}} = -5/4$. The constraint $0.98 < r_D < 1$ then corresponds to $-1.25 < f_{\text{NL}}^{\text{local}} < -1.21$. Taking into account the Planck result $f_{\text{NL}}^{\text{local}} = 2.7 \pm 5.8$ (Planck Collaboration XXIV, 2013), we conclude that the Planck data are consistent with the scenario where the curvaton decays into CDM when it dominates the energy density of the Universe, and its fluctuations are almost entirely converted into adiabatic ones.

11. Conclusions

This paper establishes the status of cosmic inflation in the context of the first release of the *Planck* cosmological results, which includes the temperature data from the first 2.6 sky surveys. CMB polarization as measured by *Planck* will be the subject of a future release. We find that standard slow-roll single field inflation is compatible with the *Planck* data. This result is confirmed by other papers of this series. *Planck* in combination with WMAP 9-year large angular scale polarization (WP) yields $\Omega_K = -0.006 \pm 0.018$ at 95% CL by combining temperature and lensing information (Planck Collaboration XVI, 2013; Planck Collaboration XVII, 2013). The bispectral non-Gaussianity parameter f_{NL} measured by *Planck* is consistent with zero (Planck Collaboration XXIV, 2013). These results are consistent with zero spatial curvature and a small value of f_{NL} , as predicted in the simplest slow-roll inflationary models.

A key *Planck* result is the measurement of the scalar perturbation spectral index. *Planck*+WP data give $n_s = 0.9603 \pm 0.0073$ (and $n_s = 0.9629 \pm 0.0057$ when also combined with BAO). This result disfavors the Harrison-Zeldovich (HZ) $n_s = 1$ model at more than 5σ . Even in extended cosmological models, the HZ spectrum cannot be reconciled with the data. Allowing a general reionisation scenario yields $\Delta\chi_{\text{eff}}^2 = 12.5$ with respect to Λ CDM for *Planck*+WP data. When the primordial Helium abundance or the effective number of neutrino species are allowed to vary, the best-fit of the HZ model to a combination of *Planck*+WP and BAO data is still worse by $\Delta\chi_{\text{eff}}^2 = 4.6$, and 8.0, respectively.

We find no evidence for *Planck* data preferring a generalization of a simple power law spectrum to include a running of the spectral index ($dn_s/d\ln k = -0.0134 \pm 0.0090$) or a running of the running ($d^2n_s/d\ln k^2 = 0.020^{+0.016}_{-0.015}$ with *Planck*+WP). In a model admitting tensor fluctuations, the 95% CL bound on the tensor-to-scalar ratio is $r_{0.002} < 0.12$ (< 0.11) using *Planck*+WP (plus high- ℓ). This bound on r implies an upper limit for the inflation energy scale of 1.9×10^{16} GeV or, equivalently, for the Hubble parameter $H_* < 3.7 \times 10^{-5} M_{\text{pl}}$, at 95% CL.

The degeneracy between n_s and r , which plagued previous CMB measurements, is now removed by the *Planck* precision in the determination of the highest acoustic peaks. Inflaton potentials with a concave shape are favoured and occupy most of the 95% confidence region allowed by *Planck*+WP in the $n_s - r$ plane. Models with an exponential potential, a monomial potential with a power larger than two, or hybrid models driven by a quadratic term are disfavoured at more than 95% confidence. The quadratic large field model, in the past often cited as the simplest inflationary model, is now at the edge of the 95% CL contours allowed by *Planck*+WP+high- ℓ CMB data.

A Bayesian parameter estimation and model comparison analysis of a representative sample of single-field slow-roll models shows that *Planck* is able to discriminate between these models with results that are robust even when a broad set of entropy generation scenarios are allowed. In addition to confirming the exclusion of the ϕ^4 potential, the Bayesian evidence computed from the *Planck* data provides significant odds (logarithms of the

Bayes factor of ~ -5 or lower relative to Λ CDM) against large-field models compatible with previous cosmological data, such as the ϕ^2 potential, and two-parameter potentials such as natural inflation and the hilltop potential. As presented in Sect. 5, *Planck* establishes strong constraints on the parameter values of specific inflationary scenarios. For example, the scale parameter of the natural inflation potential is constrained to be $\log(f/M_{\text{pl}}) \gtrsim 1.1$ (95% CL), improving upon the WMAP7 limit on f by a factor of two. The *Planck* data limit the possibilities for the unexplored physics between the end of inflation and the beginning of the radiation-dominated era. Data-driven constraints are obtained on w_{int} , the effective equation of state in the post-inflationary era. Particularly for the disfavoured models listed above, their parameters are pushed to unnatural values $w_{\text{int}} \gtrsim 1/3$ in order to become more compatible with the data.

Using an essentially exact numerical calculation of the predicted primordial spectrum, we reconstruct the observable window of the inflaton potential, expanding the potential as a Taylor series up to a fixed order. For an observable potential described by a polynomial of order three, the reconstruction agrees well with the slow-roll predictions. If a quartic term is allowed, the result deviate from the slow-roll prediction, because the *Planck* data favour a slightly smaller amplitude for the Sachs-Wolfe plateau relative to the $\ell > 40$ part of the power spectrum than the best-fitting minimal Λ CDM model with a power law primordial spectrum. A potential with an fourth order polynomial can fit this feature, thus reducing the effective χ_{eff}^2 by approximately four.

A penalized-likelihood reconstruction of the primordial power spectrum shows hints of structure at modest estimated statistical significance. Possible sources of unaccounted errors are discussed. Parameterized models producing superimposed oscillations (possibly motivated by deviations from the Bunch-Davies vacuum state, axion monodromy, or a sharp step in the inflaton potential) improve the χ_{eff}^2 by roughly 10, where three extra parameters have been added. However, a Bayesian model comparison analysis does not strongly favour the model with oscillations over the standard featureless power spectrum. With *Planck* polarization data, a more conclusive result on superimposed oscillations is expected.

We combine power-spectrum constraints with those on the non-linearity parameter f_{NL} (Planck Collaboration XXIV, 2013) to constrain single field inflation with generalized Lagrangians, in which non-Gaussianities are larger than those predicted by the simplest slow-roll inflationary models. We show how the limits on the inflation sound speed derived in Planck Collaboration XXIV (2013) are crucial to constrain slow-roll parameters for generalized Lagrangians. We also show how particular examples of DBI Inflation and k -inflation can be constrained by this combination of *Planck* data.

We test the hypothesis that the primordial cosmological perturbations were exclusively adiabatic. We analyse all non-singular (i.e., non-decaying) isocurvature modes arbitrarily correlated to the adiabatic mode, using a parameterization where the isocurvature contributions are specified at two scales. The oscillatory pattern in the *Planck* temperature spectrum is compatible with purely adiabatic perturbations, and therefore constrains any isocurvature contribution to be small at those multipoles. As a consequence, axion and curvaton scenarios, in which the CDM isocurvature mode is uncorrelated or fully correlated with the adiabatic mode, are not favoured by *Planck*. The upper bounds on the isocurvature fraction at $k = 0.05 \text{ Mpc}^{-1}$ are 0.039 for the axion, and 0.0025 for the curvaton, at 95% CL. However general models with an arbitrarily correlated mixture of adiabatic

and (CDM or neutrino) isocurvature modes have the freedom to lower the Sachs-Wolfe plateau relative to the high- ℓ spectrum, and reduce the effective χ^2_{eff} by more than four.

The simplest inflationary models have passed an exacting test with the *Planck* data. The full mission data including *Planck*'s polarization measurements will help answer further fundamental questions, including the possibilities for non-smooth power spectra, the energy scale of inflation, and extensions to more complex models.

Acknowledgements. The development of *Planck* has been supported by: ESA; CNES and CNRS/INSU-IN2P3-INP (France); ASI, CNR, and INAF (Italy); NASA and DoE (USA); STFC and UKSA (UK); CSIC, MICINN, JA and RES (Spain); Tekes, AoF and CSC (Finland); DLR and MPG (Germany); CSA (Canada); DTU Space (Denmark); SER/SSO (Switzerland); RCN (Norway); SFI (Ireland); FCT/MCTES (Portugal); and PRACE (EU). A description of the Planck Collaboration and a list of its members, including the technical or scientific activities in which they have been involved, can be found at http://www.sciops.esa.int/index.php?project=planck&page=Planck_Collaboration. We especially thank Julien Tréguer and James Zibin for invaluable help with perfecting the figures and a critical reading of the manuscript. We gratefully acknowledge IN2P3 Computer Center (<http://cc.in2p3.fr>) for providing a significant amount of the computing resources and services needed for this work.

References

- Abbott, L. & Wise, M. B. 1984, Nucl.Phys., B244, 541
- Achúcarro, A., Gong, J.-O., Hardeman, S., Palma, G. A., & Patil, S. P. 2011, JCAP, 1101, 030, [arXiv:1010.3693](#)
- Acquaviva, V., Bartolo, N., Matarrese, S., & Riotto, A. 2003, Nuclear Physics B, 667, 119, [arXiv:astro-ph/0209156](#)
- Adams, F. C., Bond, J. R., Freese, K., Frieman, J. A., & Olinto, A. V. 1993, Phys.Rev., D47, 426, [arXiv:hep-ph/9207245](#)
- Adams, J., Cresswell, B., & Easther, R. 2001, Phys. Rev. D, 64, 123514, [arXiv:astro-ph/0102236](#)
- Adams, J. A., Cresswell, B., & Easther, R. 2001, Phys.Rev., D64, 123514, [arXiv:astro-ph/0102236](#)
- Adshead, P., Dvorkin, C., Hu, W., & Lim, E. A. 2012, Phys.Rev., D85, 023531, [arXiv:1110.3050](#)
- Adshead, P., Easther, R., Pritchard, J., & Loeb, A. 2011, JCAP, 1102, 021, [arXiv:1007.3748](#)
- Agarwal, N. & Bean, R. 2009, Phys. Rev. D, 79, 023503, [arXiv:0809.2798](#)
- Akaike, H. 1974, IEEE Transactions on Automatic Control, 19, 716
- Albrecht, A. & Steinhardt, P. J. 1982, Phys.Rev.Lett., 48, 1220
- Alishahiha, M., Silverstein, E., & Tong, D. 2004, Phys. Rev. D, 70, 123505, [arXiv:hep-th/0404084](#)
- Allahverdi, R., Brandenberger, R., Cyr-Racine, F.-Y., & Mazumdar, A. 2010, Ann.Rev.Nucl.Part.Sci., 60, 27, [arXiv:1001.2600](#)
- Amendola, L., Gordon, C., Wands, D., & Sasaki, M. 2002, Phys.Rev.Lett., 88, 211302, [arXiv:astro-ph/0107089](#)
- Anantua, R., Easther, R., & Giblin, J. T. 2009, Phys.Rev.Lett., 103, 111303, [arXiv:0812.0825](#)
- Anderson, L., Aubourg, E., Bailey, S., et al. 2012, MNRAS, 427, 3435, [arXiv:1203.6594](#)
- Armendáriz-Picón, C., Damour, T., & Mukhanov, V. 1999, Physics Letters B, 458, 209, [arXiv:hep-th/9904075](#)
- Audren, B., Lesgourgues, J., Benabed, K., & Prunet, S. 2012, [arXiv:1210.7183](#)
- Aver, E., Olive, K. A., & Skillman, E. D. 2012, JCAP, 1204, 004, [arXiv:1112.3713](#)
- Axenides, M., Brandenberger, R. H., & Turner, M. S. 1983, Phys.Lett., B126, 178
- Babich, D., Creminelli, P., & Zaldarriaga, M. 2004, J. Cosmology Astropart. Phys., 8, 9, [arXiv:astro-ph/0405356](#)
- Bae, K. J., Huh, J.-H., & Kim, J. E. 2008, JCAP, 0809, 005, [arXiv:0806.0497](#)
- Banks, T., Kaplan, D. B., & Nelson, A. E. 1994, Phys. Rev., D49, 779, [arXiv:hep-ph/9308292](#)
- Bardeen, J. M., Steinhardt, P. J., & Turner, M. S. 1983, Phys.Rev., D28, 679
- Barrow, J. D. 1990, Phys.Lett., B235, 40
- Barrow, J. D. & Liddle, A. R. 1993, Phys.Rev., D47, 5219, [arXiv:astro-ph/9303011](#)
- Bartolo, N., Komatsu, E., Matarrese, S., & Riotto, A. 2004a, Phys.Rept., 402, 103, [arXiv:astro-ph/0406398](#)
- Bartolo, N. & Liddle, A. R. 2002, Phys.Rev., D65, 121301, [arXiv:astro-ph/0203076](#)
- Bartolo, N., Matarrese, S., & Riotto, A. 2004b, Phys.Rev.Lett., 93, 231301, [arXiv:astro-ph/0407505](#)
- Bartolo, N., Matarrese, S., & Riotto, A. 2004c, Phys.Rev., D69, 043503, [arXiv:hep-ph/0309033](#)
- Baumann, D. & McAllister, L. 2007, Phys. Rev. D, 75, 123508, [arXiv:hep-th/0610285](#)
- Bean, R., Chen, X., Peiris, H., & Xu, J. 2008, Phys. Rev. D, 77, 023527, [arXiv:0710.1812](#)
- Bean, R., Dunkley, J., & Pierpaoli, E. 2006, Phys.Rev., D74, 063503, [arXiv:astro-ph/0606685](#)
- Bean, R., Shandera, S. E., Tye, S.-H. H., & Xu, J. 2007, J. Cosmology Astropart. Phys., 5, 4, [arXiv:hep-th/0702107](#)
- Beltrán, M., García-Bellido, J., & Lesgourgues, J. 2007, Phys.Rev., D75, 103507, [arXiv:hep-ph/0606107](#)
- Beltrán, M., García-Bellido, J., Lesgourgues, J., & Riazuelo, A. 2004, Phys.Rev., D70, 103530, [arXiv:astro-ph/0409326](#)
- Benetti, M., Pandolfi, S., Lattanzi, M., Martinelli, M., & Melchiorri, A. 2013, Phys.Rev., D87, 023519, [arXiv:1210.3562](#)
- Bennett, C., Hill, R., Hinshaw, G., et al. 2011, Astrophys.J.Suppl., 192, 17, [arXiv:1001.4758](#)
- Bennett, C., Larson, D., Weiland, J., et al. 2012, [arXiv:1212.5225](#)
- Beutler, F., Blake, C., Colless, M., et al. 2011, Mon.Not.Roy.Astron.Soc., 416, 3017, [arXiv:1106.3366](#)
- Bezrukov, F. & Shaposhnikov, M. 2009, JHEP, 07, 089, [arXiv:0904.1537](#)
- Blas, D., Lesgourgues, J., & Tram, T. 2011, JCAP, 1107, 034, [arXiv:1104.2933](#)
- Bozza, V., Giovannini, M., & Veneziano, G. 2003, JCAP, 0305, 001, [arXiv:hep-th/0302184](#)
- Bridges, M., Feroz, F., Hobson, M. P., & Lasenby, A. N. 2009, MNRAS, 400, 1075, [arXiv:0812.3541](#)
- Bridle, S., Lewis, A., Weller, J., & Efstathiou, G. 2003, Mon.Not.Roy.Astron.Soc., 342, L72, [arXiv:astro-ph/0302306](#)
- Brout, R., Englert, F., & Gunzig, E. 1978, Annals Phys., 115, 78
- Bucher, M. & Cohn, J. 1997, Phys.Rev., D55, 7461, [arXiv:astro-ph/9701117](#)
- Bucher, M., Dunkley, J., Ferreira, P., Moodley, K., & Skordis, C. 2004, Phys.Rev.Lett., 93, 081301, [arXiv:astro-ph/0401417](#)
- Bucher, M., Goldhaber, A. S., & Turok, N. 1995, Phys.Rev., D52, 3314, [arXiv:hep-ph/9411206](#)
- Bucher, M., Moodley, K., & Turok, N. 2000, Phys.Rev., D62, 083508, [arXiv:astro-ph/9904231](#)
- Bucher, M. & Turok, N. 1995, Phys.Rev., D52, 5538, [arXiv:hep-ph/9503393](#)
- Bunch, T. & Davies, P. 1978, Proc.Roy.Soc.Lond., A360, 117
- Burgess, C., Easther, R., Mazumdar, A., Mota, D. F., & Multamaki, T. 2005, JHEP, 0505, 067, [arXiv:hep-th/0501125](#)
- Byrnes, C. T. & Wands, D. 2006, Phys.Rev., D74, 043529, [arXiv:astro-ph/0605679](#)
- Casadio, R., Finelli, F., Kamenshchik, A., Luzzi, M., & Venturi, G. 2006, JCAP, 0604, 011, [arXiv:gr-qc/0603026](#)
- Chen, X. 2005a, Journal of High Energy Physics, 8, 45, [arXiv:hep-th/0501184](#)
- Chen, X. 2005b, Phys. Rev. D, 71, 063506, [arXiv:hep-th/0408084](#)
- Chen, X. 2005c, Phys. Rev. D, 72, 123518, [arXiv:astro-ph/0507053](#)
- Chen, X. 2010, Advances in Astronomy, 2010, [arXiv:1002.1416](#)
- Chen, X., Huang, M.-x., Kachru, S., & Shiu, G. 2007, J. Cosmology Astropart. Phys., 1, 2, [arXiv:hep-th/0605045](#)
- Cheung, C., Creminelli, P., Fitzpatrick, A. L., Kaplan, J., & Senatore, L. 2008, JHEP, 0803, 014, [arXiv:0709.0293](#)
- Chung, D. J., Everett, L. L., & Matchev, K. T. 2007, Phys.Rev., D76, 103530, [arXiv:0704.3285](#)
- Chung, D. J., Kolb, E. W., Riotto, A., & Tkachev, I. I. 2000, Phys.Rev., D62, 043508, [arXiv:hep-ph/9910437](#)
- Coleman, S. R. & De Luccia, F. 1980, Phys.Rev., D21, 3305
- Contaldi, C. R., Peloso, M., Kofman, L., & Linde, A. D. 2003, JCAP, 0307, 002, [arXiv:astro-ph/0303636](#)
- Cortès, M. & Liddle, A. R. 2009, Phys.Rev., D80, 083524, [arXiv:0905.0289](#)
- Cortès, M., Liddle, A. R., & Mukherjee, P. 2007, Phys.Rev., D75, 083520, [arXiv:astro-ph/0702170](#)
- Covi, L., Hamann, J., Melchiorri, A., Slosar, A., & Sorbera, I. 2006, Phys.Rev., D74, 083509, [arXiv:astro-ph/0606452](#)
- Cox, R. T. 1946, Am. J. Phys., 14, 1
- Danielsson, U. H. 2002, Phys.Rev., D66, 023511, [arXiv:hep-th/0203198](#)
- Das, S., Louis, T., Nolte, M. R., et al. 2013, [arXiv:1301.1037](#)
- de Carlos, B., Casas, J. A., Quevedo, F., & Roulet, E. 1993, Phys. Lett., B318, 447, [arXiv:hep-ph/9308325](#)
- Dodelson, S., Kinney, W. H., & Kolb, E. W. 1997, Phys.Rev., D56, 3207, [arXiv:astro-ph/9702166](#)
- Dunkley, J., Bucher, M., Ferreira, P., Moodley, K., & Skordis, C. 2005, Phys.Rev.Lett., 95, 261303, [arXiv:astro-ph/0507473](#)
- Dunkley, J., Calabrese, E., Sievers, J., et al. 2013, [arXiv:1301.0776](#)

- Dunkley, J., Hlozek, R., Sievers, J., et al. 2011, *Astrophys.J.*, 739, 52, [arXiv:1009.0866](#)
- Dunkley, J., Komatsu, E., Nolte, M. R., et al. 2009, *ApJS*, 180, 306, [arXiv:0803.0586](#)
- Dvali, G. R., Shafi, Q., & Schaefer, R. K. 1994, *Phys. Rev. Lett.*, 73, 1886, [arXiv:hep-ph/9406319](#)
- Easther, R., Flauger, R., & Gilmore, J. B. 2011, *JCAP*, 1104, 027, [arXiv:1003.3011](#)
- Easther, R., Greene, B. R., Kinney, W. H., & Shiu, G. 2001, *Phys.Rev.*, D64, 103502, [arXiv:hep-th/0104102](#)
- Easther, R. & Peiris, H. 2006, *JCAP*, 0609, 010, [arXiv:astro-ph/0604214](#)
- Easther, R. & Peiris, H. V. 2012, *Phys. Rev. D*, 85, 103533, [arXiv:1112.0326](#)
- Efstathiou, G. & Bond, J. R. 1986, *MNRAS*, 218, 103
- Efstathiou, G. & Bond, J. R. 1987, *MNRAS*, 227, 33P
- Elgarøy, Ø., Hannestad, S., & Haugbølle, T. 2003, *JCAP*, 0309, 008, [arXiv:astro-ph/0306229](#)
- Enqvist, K. & Sloth, M. S. 2002, *Nucl.Phys.*, B626, 395, [arXiv:hep-ph/0109214](#)
- Eriksen, H. K. et al. 2007, *Astrophys.J.*, 656, 641, [arXiv:astro-ph/0606088](#)
- Fabbri, R. & Pollock, M. 1983, *Phys.Lett.*, B125, 445
- Fakir, R. & Unruh, W. 1990, *Phys.Rev.*, D41, 1783
- Feroz, F. & Hobson, M. 2008, *Mon.Not.Roy.Astron.Soc.*, 384, 449, [arXiv:0704.3704](#)
- Feroz, F., Hobson, M., & Bridges, M. 2009, *Mon.Not.Roy.Astron.Soc.*, 398, 1601, [arXiv:0809.3437](#)
- Finelli, F., Hamann, J., Leach, S. M., & Lesgourgues, J. 2010, *JCAP*, 1004, 011, [arXiv:0912.0522](#)
- Fixsen, D. 2009, *Astrophys.J.*, 707, 916, [arXiv:0911.1955](#)
- Flauger, R., McAllister, L., Pajer, E., Westphal, A., & Xu, G. 2010, *JCAP*, 1006, 009, [arXiv:0907.2916](#)
- Freese, K., Frieman, J. A., & Olinto, A. V. 1990, *Physical Review Letters*, 65, 3233
- Freivogel, B., Kleban, M., Rodríguez Martínez, M., & Susskind, L. 2006, *JHEP*, 0603, 039, [arXiv:hep-th/0505232](#)
- García-Bellido, J. & Wands, D. 1996, *Phys.Rev.*, D53, 5437, [arXiv:astro-ph/9511029](#)
- Garriga, J., Montes, X., Sasaki, M., & Tanaka, T. 1998, *Nucl.Phys.*, B513, 343, [arXiv:astro-ph/9706229](#)
- Garriga, J., Montes, X., Sasaki, M., & Tanaka, T. 1999, *Nucl.Phys.*, B551, 317, [arXiv:astro-ph/9811257](#)
- Garriga, J. & Mukhanov, V. F. 1999, *Physics Letters B*, 458, 219, [arXiv:hep-th/9904176](#)
- Gauthier, C. & Bucher, M. 2012, *JCAP*, 1210, 050, [arXiv:1209.2147](#)
- Gong, J.-O. & Stewart, E. D. 2001, *Phys.Lett.*, B510, 1, [arXiv:astro-ph/0101225](#)
- Gordon, C. & Lewis, A. 2003, *Phys.Rev.*, D67, 123513, [arXiv:astro-ph/0212248](#)
- Gordon, C. & Malik, K. A. 2004, *Phys.Rev.*, D69, 063508, [arXiv:astro-ph/0311102](#)
- Gordon, C., Wands, D., Bassett, B. A., & Maartens, R. 2001, *Phys.Rev.*, D63, 023506, [arXiv:astro-ph/0009131](#)
- Gott, J. 1982, *Nature*, 295, 304
- Gott, J. & Statler, T. 1984, *Phys.Lett.*, B136, 157
- Gratton, S., Hertog, T., & Turok, N. 2000, *Phys.Rev.*, D62, 063501, [arXiv:astro-ph/9907212](#)
- Gratton, S., Lewis, A., & Turok, N. 2002, *Phys.Rev.*, D65, 043513, [arXiv:astro-ph/0111012](#)
- Gratton, S. & Turok, N. 1999, *Phys.Rev.*, D60, 123507, [arXiv:astro-ph/9902265](#)
- Grishchuk, L. 1975, *Sov.Phys.JETP*, 40, 409
- Groot Nibbelink, S. & van Tent, B. 2000, [arXiv:hep-ph/0011325](#)
- Groot Nibbelink, S. & van Tent, B. 2002, *Class.Quant.Grav.*, 19, 613, [arXiv:hep-ph/0107272](#)
- Guth, A. H. 1981, *Phys.Rev.*, D23, 347
- Guth, A. H. & Nomura, Y. 2012, *Phys.Rev.*, D86, 023534, [arXiv:1203.6876](#)
- Guth, A. H. & Pi, S. 1982, *Phys.Rev.Lett.*, 49, 1110
- Habib, S., Heitmann, K., Jungman, G., & Molina-Paris, C. 2002, *Phys.Rev.Lett.*, 89, 281301, [arXiv:astro-ph/0208443](#)
- Hamann, J., Hannestad, S., Raffelt, G. G., & Wong, Y. Y. Y. 2009, *JCAP*, 0906, 022, [arXiv:0904.0647](#)
- Hamann, J., Hannestad, S., Sloth, M. S., & Wong, Y. Y. Y. 2008a, *JCAP*, 0809, 015, [arXiv:0807.4528](#)
- Hamann, J., Lesgourgues, J., & Mangano, G. 2008b, *JCAP*, 0803, 004, [arXiv:0712.2826](#)
- Hamann, J., Lesgourgues, J., & Valkenburg, W. 2008c, *JCAP*, 0804, 016, [arXiv:0802.0505](#)
- Hamann, J., Shafieloo, A., & Souradeep, T. 2010, *JCAP*, 1004, 010, [arXiv:0912.2728](#)
- Hamimeche, S. & Lewis, A. 2008, *Phys.Rev.*, D77, 103013, [arXiv:0801.0554](#)
- Hannestad, S. 2004, *JCAP*, 0404, 002, [arXiv:astro-ph/0311491](#)
- Harrison, E. R. 1970, *Phys.Rev.*, D1, 2726
- Hawking, S. 1982, *Phys. Lett.*, B115, 295
- Hawking, S. & Turok, N. 1998, *Phys.Lett.*, B425, 25, [arXiv:hep-th/9802030](#)
- Hertog, T. & Turok, N. 2000, *Phys.Rev.*, D62, 083514, [arXiv:astro-ph/9903075](#)
- Hinshaw, G., Larson, D., Komatsu, E., et al. 2012a, [arXiv:1212.5226](#)
- Hinshaw, G., Larson, D., Komatsu, E., et al. 2012b, [arXiv:1212.5226](#)
- Hou, Z., Keisler, R., Knox, L., Millea, M., & Reichardt, C. 2011, [arXiv:1104.2333](#)
- Hou, Z., Reichardt, C., Story, K., et al. 2012, [arXiv:1212.6267](#)
- Hunt, P. & Sarkar, S. 2004, *Phys.Rev.*, D70, 103518, [arXiv:astro-ph/0408138](#)
- Ichikawa, K. & Takahashi, T. 2006, *Phys.Rev.*, D73, 063528, [arXiv:astro-ph/0601099](#)
- Ichiki, K. & Nagata, R. 2009, *Phys.Rev.*, D80, 083002
- Jaynes, E. T. & Bretthorst, G. L. 2003, *Probability Theory* (Cambridge, UK: Cambridge University Press)
- Jeffreys, H. 1998, *Theory of Probability*, 3rd edn. (Oxford University Press)
- Keskitalo, R., Kurki-Suonio, H., Muhonen, V., & Valiviita, J. 2007, *JCAP*, 0709, 008, [arXiv:astro-ph/0611917](#)
- Kim, J. E. & Carosi, G. 2010, *Rev.Mod.Phys.*, 82, 557, [arXiv:0807.3125](#)
- Kinney, W. H. 2002, *Phys.Rev.*, D66, 083508, [arXiv:astro-ph/0206032](#)
- Kinney, W. H., Kolb, E. W., Melchiorri, A., & Riotto, A. 2006, *Phys.Rev.*, D74, 023502, [arXiv:astro-ph/0605338](#)
- Kinney, W. H. & Riotto, A. 2006, *J. Cosmology Astropart. Phys.*, 3, 11, [arXiv:astro-ph/0511127](#)
- Kleban, M. & Schillo, M. 2012, *JCAP*, 1206, 029, [arXiv:1202.5037](#)
- Knox, L. & Turner, M. S. 1994, *Phys.Rev.Lett.*, 73, 3347, [arXiv:astro-ph/9407037](#)
- Kobayashi, T. & Takahashi, F. 2011, *JCAP*, 1101, 026, [arXiv:1011.3988](#)
- Kofman, L. 1986, *Phys.Lett.*, B173, 400
- Kofman, L., Linde, A. D., & Starobinsky, A. A. 1997, *Phys.Rev.*, D56, 3258, [arXiv:hep-ph/9704452](#)
- Komatsu, E., Smith, K. M., Dunkley, J., et al. 2011, *ApJS*, 192, 18, [arXiv:1001.4538](#)
- Komatsu, E. et al. 2009, *Astrophys.J.Suppl.*, 180, 330, [arXiv:0803.0547](#)
- Komatsu, E. et al. 2011, *Astrophys.J.Suppl.*, 192, 18, [arXiv:1001.4538](#)
- Kosowsky, A. & Turner, M. S. 1995, *Phys.Rev.*, D52, 1739, [arXiv:astro-ph/9504071](#)
- Kurki-Suonio, H., Muhonen, V., & Valiviita, J. 2005, *Phys. Rev.*, D71, 063005, [arXiv:astro-ph/0412439](#)
- Langlois, D. & Riazuelo, A. 2000, *Phys.Rev.*, D62, 043504, [arXiv:astro-ph/9912497](#)
- Larson, D., Dunkley, J., Hinshaw, G., et al. 2011, *ApJS*, 192, 16, [arXiv:1001.4635](#)
- Leach, S. M., Liddle, A. R., Martin, J., & Schwarz, D. J. 2002, *Phys.Rev.*, D66, 023515, [arXiv:astro-ph/0202094](#)
- Lesgourgues, J. 2011, [arXiv:1104.2932](#)
- Lesgourgues, J., Starobinsky, A. A., & Valkenburg, W. 2008, *JCAP*, 0801, 010, [arXiv:0710.1630](#)
- Lesgourgues, J. & Valkenburg, W. 2007, *Phys.Rev.*, D75, 123519, [arXiv:astro-ph/0703625](#)
- Lewis, A. 2011, <http://cosmologist.info/notes/CAMB.pdf>
- Lewis, A. & Bridle, S. 2002, *Phys.Rev.*, D66, 103511, [arXiv:astro-ph/0205436](#)
- Liddle, A. R. 2007, *Mon.Not.Roy.Astron.Soc.*, 377, L74, [arXiv:astro-ph/0701113](#)
- Liddle, A. R. & Leach, S. M. 2003, *Phys. Rev.*, D68, 103503, [arXiv:astro-ph/0305263](#)
- Liddle, A. R. & Lyth, D. H. 1993, *Phys.Rept.*, 231, 1, [arXiv:astro-ph/9303019](#)
- Lidsey, J. E. & Huston, I. 2007, *J. Cosmology Astropart. Phys.*, 7, 2, [arXiv:0705.0240](#)
- Lifshitz, E. & Khalatnikov, I. 1963, *Adv.Phys.*, 12, 185
- Linde, A., Noorbala, M., & Westphal, A. 2011, *JCAP*, 1103, 013, [arXiv:1101.2652](#)
- Linde, A. D. 1982, *Phys.Lett.*, B108, 389
- Linde, A. D. 1983, *Phys.Lett.*, B129, 177
- Linde, A. D. 1984, *JETP Lett.*, 40, 1333
- Linde, A. D. 1985, *Phys.Lett.*, B158, 375
- Linde, A. D. 1990, *Particle physics and inflationary cosmology* (Harwood)
- Linde, A. D. 1991, *Phys.Lett.*, B259, 38
- Linde, A. D. 2003, *JCAP*, 0305, 002, [arXiv:astro-ph/0303245](#)
- Linde, A. D. & Lyth, D. H. 1990, *Phys.Lett.*, B246, 353
- Linde, A. D. & Mukhanov, V. F. 1997, *Phys.Rev.*, D56, 535, [arXiv:astro-ph/9610219](#)
- Lorenz, L., Martin, J., & Ringeval, C. 2008, *Phys. Rev. D*, 78, 083513, [arXiv:0807.3037](#)
- Lucchin, F. & Matarrese, S. 1985, *Phys.Rev.*, D32, 1316
- Lucy, L. B. 1974, *AJ*, 79, 745
- Lyth, D. 1992, *Phys.Rev.*, D45, 3394
- Lyth, D. H. 1990, *Phys.Lett.*, B236, 408
- Lyth, D. H. 1997, *Phys.Rev.Lett.*, 78, 1861, [arXiv:hep-ph/9606387](#)

- Lyth, D. H. & Stewart, E. D. 1990, *Phys.Lett.*, B252, 336
- Lyth, D. H. & Stewart, E. D. 1996, *Phys.Rev.*, D53, 1784, [arXiv:hep-ph/9510204](#)
- Lyth, D. H., Ungarelli, C., & Wands, D. 2003, *Phys.Rev.*, D67, 023503, [arXiv:astro-ph/0208055](#)
- Lyth, D. H. & Wands, D. 2002, *Phys.Lett.*, B524, 5, [arXiv:hep-ph/0110002](#)
- Ma, C.-P. & Bertschinger, E. 1995, *Astrophys.J.*, 455, 7, [arXiv:astro-ph/9506072](#)
- Maldacena, J. 2003, *Journal of High Energy Physics*, 5, 13, [arXiv:astro-ph/0210603](#)
- Martin, J. & Brandenberger, R. 2003, *Phys.Rev.*, D68, 063513, [arXiv:hep-th/0305161](#)
- Martin, J. & Ringeval, C. 2004, *Phys.Rev.*, D69, 083515, [arXiv:astro-ph/0310382](#)
- Martin, J. & Ringeval, C. 2010, *Phys.Rev.*, D82, 023511, [arXiv:1004.5525](#)
- Martin, J., Ringeval, C., & Trotta, R. 2011, *Phys.Rev.*, D83, 063524, [arXiv:1009.4157](#)
- Martin, J. & Schwarz, D. J. 2003, *Phys.Rev.*, D67, 083512, [arXiv:astro-ph/0210090](#)
- McAllister, L., Silverstein, E., & Westphal, A. 2010, *Phys.Rev.*, D82, 046003, [arXiv:0808.0706](#)
- Meerburg, P. D., Wijers, R. A. M. J., & van der Schaar, J. P. 2012, *MNRAS*, 421, 369, [arXiv:1109.5264](#)
- Mehta, K. T., Cuesta, A. J., Xu, X., Eisenstein, D. J., & Padmanabhan, N. 2012, *MNRAS*, 427, 2168, [arXiv:1202.0092](#)
- Mollerach, S. 1990, *Phys.Rev.*, D42, 313
- Moodley, K., Bucher, M., Dunkley, J., Ferreira, P., & Skordis, C. 2004, *Phys.Rev.*, D70, 103520, [arXiv:astro-ph/0407304](#)
- Moroi, T. & Takahashi, T. 2001, *Phys.Lett.*, B522, 215, [arXiv:hep-ph/0110096](#)
- Mortonson, M. J., Dvorkin, C., Peiris, H. V., & Hu, W. 2009, *Phys. Rev. D*, 79, 103519, [arXiv:0903.4920](#)
- Mortonson, M. J. & Hu, W. 2008a, *Astrophys.J.*, 672, 737, [arXiv:0705.1132](#)
- Mortonson, M. J. & Hu, W. 2008b, *Astrophys.J.*, 686, L53, [arXiv:0804.2631](#)
- Mortonson, M. J., Peiris, H. V., & Easther, R. 2011, *Phys.Rev.*, D83, 043505, [arXiv:1007.4205](#)
- Mukhanov, V. F. 1985, *JETP Lett.*, 41, 493
- Mukhanov, V. F. 1988, *Sov.Phys.JETP*, 67, 1297
- Mukhanov, V. F. & Chibisov, G. 1981, *JETP Lett.*, 33, 532
- Mukhanov, V. F. & Chibisov, G. 1982, *Sov.Phys.JETP*, 56, 258
- Mukhanov, V. F., Feldman, H., & Brandenberger, R. H. 1992, *Phys.Rept.*, 215, 203
- Muslimov, A. 1990, *Class.Quant.Grav.*, 7, 231
- Nagata, R. & Yokoyama, J. 2008, *Phys.Rev.*, D78, 123002, [arXiv:0809.4537](#)
- Nagata, R. & Yokoyama, J. 2009, *Phys.Rev.*, D79, 043010, [arXiv:0812.4585](#)
- Norena, J., Wagner, C., Verde, L., Peiris, H. V., & Easther, R. 2012, *Phys.Rev.*, D86, 023505, [arXiv:1202.0304](#)
- Okada, N., Rehman, M. U., & Shafi, Q. 2010, *Phys.Rev.*, D82, 043502, [arXiv:1005.5161](#)
- Okamoto, T. & Hu, W. 2003, *Phys.Rev.*, D67, 083002, [arXiv:astro-ph/0301031](#)
- Olive, K. A. 1990, *Phys.Rept.*, 190, 307
- Padmanabhan, N., Xu, X., Eisenstein, D. J., et al. 2012, *MNRAS*, 427, 2132, [arXiv:1202.0090](#)
- Page, L. et al. 2007, *Astrophys.J.Suppl.*, 170, 335, [arXiv:astro-ph/0603450](#)
- Pahud, C., Liddle, A. R., Mukherjee, P., & Parkinson, D. 2007, *Mon.Not.Roy.Astron.Soc.*, 381, 489, [arXiv:astro-ph/0701481](#)
- Pallis, C. 2006, *Nucl.Phys.*, B751, 129, [arXiv:hep-ph/0510234](#)
- Pandolfi, S., Giusarma, E., Kolb, E. W., et al. 2010, *Phys.Rev.*, D82, 123527, [arXiv:1009.5433](#)
- Peccei, R. 2008, *Lect.Notes Phys.*, 741, 3, [arXiv:hep-ph/0607268](#)
- Peebles, P. & Yu, J. 1970, *Astrophys.J.*, 162, 815
- Peebles, P. J. E. & Yu, J. T. 1970, *Astrophys. J.*, 162, 815
- Peiris, H., Baumann, D., Friedman, B., & Cooray, A. 2007, *Phys. Rev. D*, 76, 103517, [arXiv:0706.1240](#)
- Peiris, H. & Easther, R. 2006a, *JCAP*, 0607, 002, [arXiv:astro-ph/0603587](#)
- Peiris, H. & Easther, R. 2006b, *JCAP*, 0610, 017, [arXiv:astro-ph/0609003](#)
- Peiris, H., Easther, R., & Flauger, R. 2013, [arXiv:1303.2616](#)
- Peiris, H. et al. 2003, *Astrophys.J.Suppl.*, 148, 213, [arXiv:astro-ph/0302225](#)
- Peiris, H. V. & Easther, R. 2008, *JCAP*, 0807, 024, [arXiv:0805.2154](#)
- Peiris, H. V., Komatsu, E., Verde, L., et al. 2003, *ApJS*, 148, 213, [arXiv:astro-ph/0302225](#)
- Peiris, H. V. & Verde, L. 2010, *Phys.Rev.*, D81, 021302, [arXiv:0912.0268](#)
- Planck Collaboration I. 2013, In preparation
- Planck Collaboration VI. 2013, In preparation
- Planck Collaboration XV. 2013, In preparation
- Planck Collaboration XVI. 2013, In preparation
- Planck Collaboration XVII. 2013, In preparation
- Planck Collaboration XXIII. 2013, In preparation
- Planck Collaboration XXIV. 2013, In preparation
- Planck Collaboration XXV. 2013, In preparation
- Polarski, D. & Starobinsky, A. A. 1994, *Phys.Rev.*, D50, 6123, [arXiv:astro-ph/9404061](#)
- Powell, B. A. & Kinney, W. H. 2007, *JCAP*, 0708, 006, [arXiv:0706.1982](#)
- Powell, B. A., Tzirakis, K., & Kinney, W. H. 2009, *J. Cosmology Astropart. Phys.*, 4, 19, [arXiv:0812.1797](#)
- Preskill, J., Wise, M. B., & Wilczek, F. 1983, *Phys.Lett.*, B120, 127
- Raffelt, G. G. 2008, *Lect.Notes Phys.*, 741, 51, [arXiv:hep-ph/0611350](#)
- Ratra, B. & Peebles, P. 1994, *Astrophys.J.*, 432, L5
- Ratra, B. & Peebles, P. 1995, *Phys.Rev.*, D52, 1837
- Reichardt, C., Shaw, L., Zahn, O., et al. 2012, *Astrophys.J.*, 755, 70, [arXiv:1111.0932](#)
- Richardson, W. H. 1972, *J. Opt. Soc. Am.*, 62, 55
- Rubakov, V., Sazhin, M., & Veryaskin, A. 1982, *Phys.Lett.*, B115, 189
- Salopek, D., Bond, J., & Bardeen, J. M. 1989, *Phys.Rev.*, D40, 1753
- Sasaki, M. 1986, *Prog.Theor.Phys.*, 76, 1036
- Sasaki, M., Tanaka, T., & Yakushige, Y. 1997, *Phys.Rev.*, D56, 616, [arXiv:astro-ph/9702174](#)
- Sasaki, M., Valiviita, J., & Wands, D. 2006, *Phys.Rev.*, D74, 103003, [arXiv:astro-ph/0607627](#)
- Sato, K. 1981, *MNRAS*, 195, 467
- Savage, C., Freese, K., & Kinney, W. H. 2006, *Phys.Rev.*, D74, 123511, [arXiv:hep-ph/0609144](#)
- Schwarz, G. 1978, *The Annals of Statistics*, 6, 461
- Sealfon, C., Verde, L., & Jimenez, R. 2005, *Phys.Rev.*, D72, 103520, [arXiv:astro-ph/0506707](#)
- Seckel, D. & Turner, M. S. 1985, *Phys.Rev.*, D32, 3178
- Senatore, L., Smith, K. M., & Zaldarriaga, M. 2010, *J. Cosmology Astropart. Phys.*, 1, 28, [arXiv:0905.3746](#)
- Shafieloo, A. & Souradeep, T. 2004, *Phys.Rev.*, D70, 043523, [arXiv:astro-ph/0312174](#)
- Shafieloo, A. & Souradeep, T. 2008, *Phys.Rev.*, D78, 023511, [arXiv:0709.1944](#)
- Sievers, J. L., Hlozek, R. A., Nolta, M. R., et al. 2013, [arXiv:1301.0824](#)
- Sikivie, P. 2008, *Lect.Notes Phys.*, 741, 19, [arXiv:astro-ph/0610440](#)
- Silverstein, E. & Tong, D. 2004, *Phys. Rev. D*, 70, 103505, [arXiv:hep-th/0310221](#)
- Silverstein, E. & Westphal, A. 2008, *Phys.Rev.*, D78, 106003, [arXiv:0803.3085](#)
- Sinha, R. & Souradeep, T. 2006, *Phys.Rev.*, D74, 043518, [arXiv:astro-ph/0511808](#)
- Spergel, D. N., Bean, R., Doré, O., et al. 2007, *ApJS*, 170, 377, [arXiv:astro-ph/0603449](#)
- Spergel, D. N., Verde, L., Peiris, H. V., et al. 2003, *ApJS*, 148, 175
- Spokoiny, B. 1993, *Phys. Lett.*, B315, 40, [arXiv:gr-qc/9306008](#)
- Starobinsky, A. 1983, *Sov.Astron.Lett.*, 9, 302
- Starobinsky, A. A. 1979, *JETP Lett.*, 30, 682
- Starobinsky, A. A. 1980, *Phys.Lett.*, B91, 99
- Starobinsky, A. A. 1982, *Phys.Lett.*, B117, 175
- Starobinsky, A. A. 1998, *Grav.Cosmol.*, 4, 88, [arXiv:astro-ph/9811360](#)
- Starobinsky, A. A. 2005, *JETP Lett.*, 82, 169, [arXiv:astro-ph/0507193](#)
- Steinhardt, P. J. & Turner, M. S. 1983, *Phys.Lett.*, B129, 51
- Stewart, E. D. & Lyth, D. H. 1993, *Physics Letters B*, 302, 171, [arXiv:gr-qc/9302019](#)
- Stompor, R., Banday, A. J., & Gorski, Krzysztof, M. 1996, *Astrophys.J.*, 463, 8, [arXiv:astro-ph/9511087](#)
- Story, K., Reichardt, C., Hou, Z., et al. 2012, [arXiv:1210.7231](#)
- Tanaka, T. & Sasaki, M. 1994, *Phys.Rev.*, D50, 6444, [arXiv:gr-qc/9406020](#)
- Tocchini-Valentini, D., Douspis, M., & Silk, J. 2005, *Mon.Not.Roy.Astron.Soc.*, 359, 31, [arXiv:astro-ph/0402583](#)
- Tocchini-Valentini, D., Hoffman, Y., & Silk, J. 2006, *Mon.Not.Roy.Astron.Soc.*, 367, 1095, [arXiv:astro-ph/0509478](#)
- Traschen, J. H. & Brandenberger, R. H. 1990, *Phys.Rev.*, D42, 2491
- Trotta, R. 2007, *Mon.Not.Roy.Astron.Soc.*, 375, L26, [arXiv:astro-ph/0608116](#)
- Trotta, R. & Hansen, S. H. 2004, *Phys.Rev.*, D69, 023509, [arXiv:astro-ph/0306588](#)
- Turner, M. S. 1983, *Phys. Rev. D*, 28, 1243
- Turner, M. S. 1990, *Phys.Rept.*, 197, 67
- Turner, M. S. & Wilczek, F. 1991, *Phys.Rev.Lett.*, 66, 5
- Turner, M. S., Wilczek, F., & Zee, A. 1983, *Phys.Lett.*, B125, 35
- Valiviita, J. & Muhonen, V. 2003, *Phys.Rev.Lett.*, 91, 131302, [arXiv:astro-ph/0304175](#)
- Valiviita, J., Savelainen, M., Talvitie, M., Kurki-Suonio, H., & Rusak, S. 2012, *Astrophys.J.*, 753, 151, [arXiv:1202.2852](#)
- Vázquez, J. A., Bridges, M., Hobson, M., & Lasenby, A. 2012, [arXiv:1203.1252](#)
- Verde, L. & Peiris, H. V. 2008, *JCAP*, 0807, 009, [arXiv:0802.1219](#)
- Vilenkin, A. 2007, *J.Phys.*, A40, 6777, [arXiv:hep-th/0609193](#)
- Yamamoto, K., Sasaki, M., & Tanaka, T. 1995, *Astrophys.J.*, 455, 412, [arXiv:astro-ph/9501109](#)

Zeldovich, Y. 1972, *Mon.Not.Roy.Astron.Soc.*, 160, 1P

- ¹ APC, AstroParticule et Cosmologie, Université Paris Diderot, CNRS/IN2P3, CEA/Irfu, Observatoire de Paris, Sorbonne Paris Cité, 10, rue Alice Domon et Léonie Duquet, 75205 Paris Cedex 13, France
- ² Aalto University Metsähovi Radio Observatory, Metsähovintie 114, FIN-02540 Kylmäla, Finland
- ³ African Institute for Mathematical Sciences, 6-8 Melrose Road, Muizenberg, Cape Town, South Africa
- ⁴ Agenzia Spaziale Italiana Science Data Center, c/o ESRIN, via Galileo Galilei, Frascati, Italy
- ⁵ Agenzia Spaziale Italiana, Viale Liegi 26, Roma, Italy
- ⁶ Astrophysics Group, Cavendish Laboratory, University of Cambridge, J J Thomson Avenue, Cambridge CB3 0HE, U.K.
- ⁷ Astrophysics & Cosmology Research Unit, School of Mathematics, Statistics & Computer Science, University of KwaZulu-Natal, Westville Campus, Private Bag X54001, Durban 4000, South Africa
- ⁸ Atacama Large Millimeter/submillimeter Array, ALMA Santiago Central Offices, Alonso de Cordova 3107, Vitacura, Casilla 763 0355, Santiago, Chile
- ⁹ CITA, University of Toronto, 60 St. George St., Toronto, ON M5S 3H8, Canada
- ¹⁰ CNRS, IRAP, 9 Av. colonel Roche, BP 44346, F-31028 Toulouse cedex 4, France
- ¹¹ California Institute of Technology, Pasadena, California, U.S.A.
- ¹² Centre for Theoretical Cosmology, DAMTP, University of Cambridge, Wilberforce Road, Cambridge CB3 0WA U.K.
- ¹³ Centro de Estudios de Física del Cosmos de Aragón (CEFCA), Plaza San Juan, 1, planta 2, E-44001, Teruel, Spain
- ¹⁴ Computational Cosmology Center, Lawrence Berkeley National Laboratory, Berkeley, California, U.S.A.
- ¹⁵ Consejo Superior de Investigaciones Científicas (CSIC), Madrid, Spain
- ¹⁶ DSM/Irfu/SPP, CEA-Saclay, F-91191 Gif-sur-Yvette Cedex, France
- ¹⁷ DTU Space, National Space Institute, Technical University of Denmark, Elektrovej 327, DK-2800 Kgs. Lyngby, Denmark
- ¹⁸ Département de Physique Théorique, Université de Genève, 24, Quai E. Ansermet, 1211 Genève 4, Switzerland
- ¹⁹ Departamento de Física Fundamental, Facultad de Ciencias, Universidad de Salamanca, 37008 Salamanca, Spain
- ²⁰ Departamento de Física, Universidad de Oviedo, Avda. Calvo Sotelo s/n, Oviedo, Spain
- ²¹ Department of Astronomy and Astrophysics, University of Toronto, 50 Saint George Street, Toronto, Ontario, Canada
- ²² Department of Astrophysics/IMAPP, Radboud University Nijmegen, P.O. Box 9010, 6500 GL Nijmegen, The Netherlands
- ²³ Department of Electrical Engineering and Computer Sciences, University of California, Berkeley, California, U.S.A.
- ²⁴ Department of Physics & Astronomy, University of British Columbia, 6224 Agricultural Road, Vancouver, British Columbia, Canada
- ²⁵ Department of Physics and Astronomy, Dana and David Dornsife College of Letter, Arts and Sciences, University of Southern California, Los Angeles, CA 90089, U.S.A.
- ²⁶ Department of Physics and Astronomy, University College London, London WC1E 6BT, U.K.
- ²⁷ Department of Physics, Gustaf Hållströmin katu 2a, University of Helsinki, Helsinki, Finland
- ²⁸ Department of Physics, Princeton University, Princeton, New Jersey, U.S.A.
- ²⁹ Department of Physics, University of California, Berkeley, California, U.S.A.
- ³⁰ Department of Physics, University of California, One Shields Avenue, Davis, California, U.S.A.
- ³¹ Department of Physics, University of California, Santa Barbara, California, U.S.A.
- ³² Department of Physics, University of Illinois at Urbana-Champaign, 1110 West Green Street, Urbana, Illinois, U.S.A.
- ³³ Dipartimento di Fisica e Astronomia G. Galilei, Università degli Studi di Padova, via Marzolo 8, 35131 Padova, Italy
- ³⁴ Dipartimento di Fisica e Scienze della Terra, Università di Ferrara, Via Saragat 1, 44122 Ferrara, Italy
- ³⁵ Dipartimento di Fisica, Università La Sapienza, P. le A. Moro 2, Roma, Italy
- ³⁶ Dipartimento di Fisica, Università degli Studi di Milano, Via Celoria, 16, Milano, Italy
- ³⁷ Dipartimento di Fisica, Università degli Studi di Trieste, via A. Valerio 2, Trieste, Italy
- ³⁸ Dipartimento di Fisica, Università di Roma Tor Vergata, Via della Ricerca Scientifica, 1, Roma, Italy
- ³⁹ Discovery Center, Niels Bohr Institute, Blegdamsvej 17, Copenhagen, Denmark
- ⁴⁰ Dpto. Astrofísica, Universidad de La Laguna (ULL), E-38206 La Laguna, Tenerife, Spain
- ⁴¹ European Southern Observatory, ESO Vitacura, Alonso de Cordova 3107, Vitacura, Casilla 19001, Santiago, Chile
- ⁴² European Space Agency, ESAC, Planck Science Office, Camino bajo del Castillo, s/n, Urbanización Villafranca del Castillo, Villanueva de la Cañada, Madrid, Spain
- ⁴³ European Space Agency, ESTEC, Keplerlaan 1, 2201 AZ Noordwijk, The Netherlands
- ⁴⁴ Haverford College Astronomy Department, 370 Lancaster Avenue, Haverford, Pennsylvania, U.S.A.
- ⁴⁵ Helsinki Institute of Physics, Gustaf Hållströmin katu 2, University of Helsinki, Helsinki, Finland
- ⁴⁶ INAF - Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, Padova, Italy
- ⁴⁷ INAF - Osservatorio Astronomico di Roma, via di Frascati 33, Monte Porzio Catone, Italy
- ⁴⁸ INAF - Osservatorio Astronomico di Trieste, Via G.B. Tiepolo 11, Trieste, Italy
- ⁴⁹ INAF/IASF Bologna, Via Gobetti 101, Bologna, Italy
- ⁵⁰ INAF/IASF Milano, Via E. Bassini 15, Milano, Italy
- ⁵¹ INFN, Sezione di Bologna, Via Irnerio 46, I-40126, Bologna, Italy
- ⁵² INFN, Sezione di Roma 1, Università di Roma Sapienza, Piazzale Aldo Moro 2, 00185, Roma, Italy
- ⁵³ IPAG: Institut de Planétologie et d'Astrophysique de Grenoble, Université Joseph Fourier, Grenoble 1 / CNRS-INSU, UMR 5274, Grenoble, F-38041, France
- ⁵⁴ IUCAA, Post Bag 4, Ganeshkhind, Pune University Campus, Pune 411 007, India
- ⁵⁵ Imperial College London, Astrophysics group, Blackett Laboratory, Prince Consort Road, London, SW7 2AZ, U.K.
- ⁵⁶ Infrared Processing and Analysis Center, California Institute of Technology, Pasadena, CA 91125, U.S.A.
- ⁵⁷ Institut Néel, CNRS, Université Joseph Fourier Grenoble I, 25 rue des Martyrs, Grenoble, France
- ⁵⁸ Institut Universitaire de France, 103, bd Saint-Michel, 75005, Paris, France
- ⁵⁹ Institut d'Astrophysique Spatiale, CNRS (UMR8617) Université Paris-Sud 11, Bâtiment 121, Orsay, France
- ⁶⁰ Institut d'Astrophysique de Paris, CNRS (UMR7095), 98 bis Boulevard Arago, F-75014, Paris, France
- ⁶¹ Institute for Space Sciences, Bucharest-Magurele, Romania
- ⁶² Institute of Astronomy and Astrophysics, Academia Sinica, Taipei, Taiwan
- ⁶³ Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, U.K.
- ⁶⁴ Institute of Theoretical Astrophysics, University of Oslo, Blindern, Oslo, Norway
- ⁶⁵ Instituto de Astrofísica de Canarias, C/Vía Láctea s/n, La Laguna, Tenerife, Spain
- ⁶⁶ Instituto de Física de Cantabria (CSIC-Universidad de Cantabria), Avda. de los Castros s/n, Santander, Spain

- ⁶⁷ Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, California, U.S.A.
- ⁶⁸ Jodrell Bank Centre for Astrophysics, Alan Turing Building, School of Physics and Astronomy, The University of Manchester, Oxford Road, Manchester, M13 9PL, U.K.
- ⁶⁹ Kavli Institute for Cosmology Cambridge, Madingley Road, Cambridge, CB3 0HA, U.K.
- ⁷⁰ LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France
- ⁷¹ LERMA, CNRS, Observatoire de Paris, 61 Avenue de l'Observatoire, Paris, France
- ⁷² Laboratoire AIM, IRFU/Service d'Astrophysique - CEA/DSM - CNRS - Université Paris Diderot, Bât. 709, CEA-Saclay, F-91191 Gif-sur-Yvette Cedex, France
- ⁷³ Laboratoire Traitement et Communication de l'Information, CNRS (UMR 5141) and Télécom ParisTech, 46 rue Barrault F-75634 Paris Cedex 13, France
- ⁷⁴ Laboratoire de Physique Subatomique et de Cosmologie, Université Joseph Fourier Grenoble I, CNRS/IN2P3, Institut National Polytechnique de Grenoble, 53 rue des Martyrs, 38026 Grenoble cedex, France
- ⁷⁵ Laboratoire de Physique Théorique, Université Paris-Sud 11 & CNRS, Bâtiment 210, 91405 Orsay, France
- ⁷⁶ Lawrence Berkeley National Laboratory, Berkeley, California, U.S.A.
- ⁷⁷ Leung Center for Cosmology and Particle Astrophysics, National Taiwan University, Taipei 10617, Taiwan
- ⁷⁸ Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str. 1, 85741 Garching, Germany
- ⁷⁹ McGill Physics, Ernest Rutherford Physics Building, McGill University, 3600 rue University, Montréal, QC, H3A 2T8, Canada
- ⁸⁰ MilliLab, VTT Technical Research Centre of Finland, Tietotie 3, Espoo, Finland
- ⁸¹ Niels Bohr Institute, Blegdamsvej 17, Copenhagen, Denmark
- ⁸² Observational Cosmology, Mail Stop 367-17, California Institute of Technology, Pasadena, CA, 91125, U.S.A.
- ⁸³ Optical Science Laboratory, University College London, Gower Street, London, U.K.
- ⁸⁴ SB-ITP-LPPC, EPFL, CH-1015, Lausanne, Switzerland
- ⁸⁵ SISSA, Astrophysics Sector, via Bonomea 265, 34136, Trieste, Italy
- ⁸⁶ School of Physics and Astronomy, Cardiff University, Queens Buildings, The Parade, Cardiff, CF24 3AA, U.K.
- ⁸⁷ School of Physics and Astronomy, University of Nottingham, Nottingham NG7 2RD, U.K.
- ⁸⁸ Space Research Institute (IKI), Russian Academy of Sciences, Profsoyuznaya Str, 84/32, Moscow, 117997, Russia
- ⁸⁹ Space Sciences Laboratory, University of California, Berkeley, California, U.S.A.
- ⁹⁰ Special Astrophysical Observatory, Russian Academy of Sciences, Nizhnij Arkhyz, Zelenchukskiy region, Karachai-Cherkessian Republic, 369167, Russia
- ⁹¹ Stanford University, Dept of Physics, Varian Physics Bldg, 382 Via Pueblo Mall, Stanford, California, U.S.A.
- ⁹² Sub-Department of Astrophysics, University of Oxford, Keble Road, Oxford OX1 3RH, U.K.
- ⁹³ Theory Division, PH-TH, CERN, CH-1211, Geneva 23, Switzerland
- ⁹⁴ UPMC Univ Paris 06, UMR7095, 98 bis Boulevard Arago, F-75014, Paris, France
- ⁹⁵ Université de Toulouse, UPS-OMP, IRAP, F-31028 Toulouse cedex 4, France
- ⁹⁶ University of Granada, Departamento de Física Teórica y del Cosmos, Facultad de Ciencias, Granada, Spain
- ⁹⁷ University of Miami, Knight Physics Building, 1320 Campo Sano Dr., Coral Gables, Florida, U.S.A.
- ⁹⁸ Warsaw University Observatory, Aleje Ujazdowskie 4, 00-478 Warszawa, Poland

Appendix A: Sampling the Hubble flow functions

In this appendix we briefly review how to constrain slow-roll inflation by sampling the Hubble flow functions (HFFs), and discuss how well the results agree with those derived by sampling directly the parameters $\ln A_s$, n_s , r , and $dn_s/d\ln k$. This method fully exploits an analytic perturbative expansion in terms of the HFF for the primordial spectra of cosmological fluctuations during slow-roll inflation (Stewart & Lyth, 1993; Gong & Stewart, 2001; Leach et al., 2002), which self-consistently extends to highest order the first terms presented in Eqs. 13-19. Since z''/z in Eq. 7 and a''/a in Eq. 9 can be rewritten exactly in terms of the Hubble flow functions, the HFF hierarchy, rather than the potential one, is best suited for this purpose. The slow-roll analytic power spectra have been calculated up to second order using the Green's function method (Gong & Stewart, 2001; Leach et al., 2002). Other approximations are available in the literature, including WKB (Martin & Schwarz, 2003), uniform approximation (Habib et al., 2002), or the method of comparison equations (Casadio et al., 2006).

The dependence of the amplitudes in Eqs. 13 and 14 in terms of HFF is given by

$$A_X = A_{X0} e^{b_X}, \quad (\text{A.1})$$

where $X = s, t$, and b_{s0}, b_{t0} are:

$$\begin{aligned} b_{s0} = & -2(C+1)\epsilon_1 - C\epsilon_2 + \left(-2C + \frac{\pi^2}{2} - 7\right)\epsilon_1^2 \\ & + \left(\frac{\pi^2}{8} - 1\right)\epsilon_2^2 + \left(-X^2 - 3X + \frac{7\pi^2}{12} - 7 + \Delta_{s0}\right)\epsilon_1\epsilon_2 \\ & + \left(-\frac{1}{2}X^2 + \frac{\pi^2}{24} + \Delta_{s0}\right)\epsilon_2\epsilon_3, \end{aligned} \quad (\text{A.2})$$

$$\begin{aligned} b_{t0} = & -2(C+1)\epsilon_1 + \left(-2C + \frac{\pi^2}{2} - 7\right)\epsilon_1^2 \\ & + \left(-C^2 - 2C + \frac{\pi^2}{12} - 2 + \Delta_{t0}\right)\epsilon_1\epsilon_2, \end{aligned} \quad (\text{A.3})$$

and $C \equiv \ln 2 + \gamma_E - 2 \approx -0.7296$ (γ_E is the Euler-Mascheroni constant). At second order the coefficients of the expansion depend on the approximation scheme: $X = C$ and $\Delta_{s0} = \Delta_{t0} = 0$ apply for the Green's function method (Gong & Stewart, 2001; Leach et al., 2002); $X = D = 1/3 - \ln 3$, $\Delta_{s0} = (D - C)(D + \ln 2) - 1/18$, $\Delta_{t0} = 2D(D - C) - 1/9$ apply for the method of comparison equations (Casadio et al., 2006). As predicted by the consistency relation, $A_{h0} = 16\epsilon_1 A_{R0}$.

The full perturbative expressions up to second order in HFF for the spectral indices and the running of the indices are

$$n_s - 1 = -2\epsilon_1 - \epsilon_2 - 2\epsilon_1^2 - (2C + 3)\epsilon_1\epsilon_2 - C\epsilon_2\epsilon_3, \quad (\text{A.4})$$

$$dn_s/d\ln k = -2\epsilon_1\epsilon_2 - \epsilon_2\epsilon_3, \quad (\text{A.5})$$

$$n_t = -2\epsilon_1 - 2\epsilon_1^2 - 2(C + 1)\epsilon_1\epsilon_2, \quad (\text{A.6})$$

$$dn_t/d\ln k = -2\epsilon_1\epsilon_2. \quad (\text{A.7})$$

We now constrain these parameters using the *Planck*+WP data. If we restrict ourselves to first order, we obtain $\epsilon_1 < 0.0074$ at 95% CL and $\epsilon_2 = 0.030^{+0.010}_{-0.009}$ at 68% CL. At second order with the Green's function method, we obtain $\epsilon_1 < 0.013$ at 95% CL, $\epsilon_2 = 0.043^{+0.013}_{-0.014}$ and $\epsilon_3 = 0.36^{+0.19}_{-0.22}$ at 68% CL. The comparison of results at first and second order is shown in Fig. A.1. Different approximation schemes lead to small differences, as Fig. A.2 shows for the Green's function method versus the method of comparison equations. Fig. A.3 shows the agreement between the physical parameters reconstructed from the HFF method and those directly sampled as in Sect. 4.3. We conclude that *Planck* sensitivity has carried the slow-roll inflationary analysis to a regime in which differences among analytic predictions and/or different priors are sub-leading effects up to second order.




Fig. A.1. *Planck* constraints on the HFFs (ϵ_1 , ϵ_2 , ϵ_3) assuming either $\epsilon_3 = 0$ and the first-order slow-roll approximation for the computation of the primordial spectra, or $\epsilon_3 \neq 0$ and the second-order slow-roll approximation (HFF GFM).




Fig. A.2. Comparison of the *Planck* constraints on the HFFs (ϵ_1 , ϵ_2 , ϵ_3) using the Green's function method (GFM) and the method of comparison equations (MCE).




Fig. A.3. *Planck* constraints on the spectral parameters $n_s, dn_s/d \ln k, r$. We compare constraints computed with the second-order slow-roll approximation, starting from flat priors on the HFF parameters at the pivot scale ($\epsilon_1, \epsilon_2, \epsilon_3$), with those obtained directly from $n_s, dn_s/d \ln k, r$. In the latter case we enforce the second-order consistency conditions for the tensor-to-scalar ratio and for the running of the tensor spectral index.