



Lecture 2

- Power spectrum of gravitational anisotropy
- Temperature anisotropy from sound waves

COBE 4-year Power Spectrum



The SW formula
allows us to determine
the **3d power
spectrum of ϕ** at
the last scattering
surface from C_l .

But how?

Sachs & Wolfe (1967) $\frac{\Delta T(\hat{n})}{T_0} = \frac{1}{3} \Phi(t_L, \hat{r}_L)$


SW Power Spectrum

$$a_{\ell m}^{\text{SW}} = \frac{4\pi T_0 i^\ell}{3} \int \frac{d^3 q}{(2\pi)^3} \Phi_{\mathbf{q}} j_\ell(q r_L) Y_\ell^{m*}(\hat{\mathbf{q}})$$

$$C_\ell \equiv \frac{1}{2\ell + 1} \sum_{m=-\ell}^{\ell} a_{\ell m} a_{\ell m}^*$$

gives...

$$C_{\ell, \text{SW}} = \frac{4\pi T_0^2}{9} \int \frac{d^3 q}{(2\pi)^3} \int \frac{d^3 q'}{(2\pi)^3} \Phi_{\mathbf{q}} \Phi_{\mathbf{q}'}^* j_\ell(q r_L) j_\ell(q' r_L) P_\ell(\hat{\mathbf{q}} \cdot \hat{\mathbf{q}}')$$



- But this is not exactly what we want. We want the **statistical average** of this quantity.

Power Spectrum of ϕ

- Statistical average of the right hand side contains

$$\langle \Phi_{\mathbf{q}} \Phi_{\mathbf{q}'}^* \rangle = \int d^3x \int d^3r \langle \Phi(\mathbf{x}) \Phi(\mathbf{x} + \mathbf{r}) \rangle \exp [i(\mathbf{q} - \mathbf{q}') \cdot \mathbf{x} - i\mathbf{q}' \cdot \mathbf{r}]$$

two-point correlation function

If $\langle \Phi(\mathbf{x}) \Phi(\mathbf{x} + \mathbf{r}) \rangle$ does not depend on locations ($\mathbf{x}$) but only on separations between two points ($\mathbf{r}$), then

$$\langle \Phi_{\mathbf{q}} \Phi_{\mathbf{q}'}^* \rangle = (2\pi)^3 \delta_D^{(3)}(\mathbf{q} - \mathbf{q}') \int d^3r \xi_{\phi}(\mathbf{r}) \exp(-i\mathbf{q} \cdot \mathbf{r})$$

consequence of “statistical homogeneity”

where we defined $\xi_{\phi}(\mathbf{r}) \equiv \langle \Phi(\mathbf{x}) \Phi(\mathbf{x} + \mathbf{r}) \rangle$

and used $\int d^3x \exp(i\mathbf{q} \cdot \mathbf{x}) = (2\pi)^3 \delta_D^{(3)}(\mathbf{q})$

Power Spectrum of ϕ

- In addition, if $\xi_\phi(\mathbf{r}) \equiv \langle \Phi(\mathbf{x})\Phi(\mathbf{x} + \mathbf{r}) \rangle$ depends only on the magnitude of the separation r and not on the directions, then

$$\langle \Phi_{\mathbf{q}} \Phi_{\mathbf{q}'}^* \rangle = (2\pi)^3 \delta_D^{(3)}(\mathbf{q} - \mathbf{q}') \int 4\pi r^2 dr \xi_\phi(r) \frac{\sin(qr)}{qr}$$

$$= (2\pi)^3 \delta_D^{(3)}(\mathbf{q} - \mathbf{q}') P_\phi(q)$$

Power spectrum!

Generic definition of the power spectrum for statistically homogeneous and isotropic fluctuations

SW Power Spectrum


- Thus, the power spectrum of the CMB in the SW limit is

$$\langle C_{\ell, \text{SW}} \rangle = \frac{16\pi^2 T_0^2}{9} \int_0^\infty \frac{q^2 dq}{(2\pi)^3} P_\phi(q) j_\ell^2(qr_L)$$

- In the flat-sky approximation,

$$\langle C_{\ell, \text{SW}} \rangle = \frac{T_0^2}{9r_L^2} \int_{-\infty}^\infty \frac{dq_{\parallel}}{2\pi} P_\phi \left(\sqrt{\frac{\ell^2}{r_L^2} + q_{\parallel}^2} \right)$$

Perpendicular
wavenumber, $(q_{\text{perp}})^2$



SW Power Spectrum

- Thus, the power spectrum of the CMB in the SW limit is

$$\langle C_{\ell, \text{SW}} \rangle = \frac{16\pi^2 T_0^2}{9} \int_0^\infty \frac{q^2 dq}{(2\pi)^3} P_\phi(q) j_\ell^2(qr_L)$$

- In the flat-sky approximation,

$$\langle C_{\ell, \text{SW}} \rangle = \frac{T_0^2}{9r_L^2} \int_{-\infty}^\infty \frac{dq_{\parallel}}{2\pi} P_\phi \left(\sqrt{\frac{\ell^2}{r_L^2} + q_{\parallel}^2} \right)$$

For a power-law form, $P_\phi(q) = (2\pi)^3 N_\phi^2 q^{n-4}$, we get

$$\langle C_{\ell, \text{SW}} \rangle = \frac{8\pi^2 N_\phi^2 T_0^2}{9\ell^2} \left(\frac{\ell}{r_L} \right)^{n-1} \frac{\sqrt{\pi}}{2} \frac{\Gamma[(3-n)/2]}{\Gamma[(4-n)/2]}$$

SW Power Spectrum

- Thus, the power spectrum of the CMB in the SW limit is

$$\langle C_{\ell, \text{SW}} \rangle = \frac{16\pi^2 T_0^2}{9} \int_0^\infty \frac{q^2 dq}{(2\pi)^3} P_\phi(q) j_\ell^2(qr_L)$$


- In the flat-sky approximation,

$$\langle C_{\ell, \text{SW}} \rangle = \frac{T_0^2}{9r_L^2} \int_{-\infty}^\infty \frac{dq_{\parallel}}{2\pi} P_\phi \left(\sqrt{\frac{\ell^2}{r_L^2} + q_{\parallel}^2} \right)$$



For a power-law form, $P_\phi(q) = (2\pi)^3 N_\phi^2 q^{n-4}$, we get

$$\langle C_{\ell, \text{SW}} \rangle = \frac{8\pi^2 N_\phi^2 T_0^2}{9\ell^2} \left(\frac{\ell}{r_L} \right)^{n-1} \frac{\sqrt{\pi}}{2} \frac{\Gamma[(3-n)/2]}{\Gamma[(4-n)/2]} \xrightarrow{n=1} \frac{8\pi^2 N_\phi^2 T_0^2}{9\ell(\ell+1)}$$



full-sky correction




COBE 4-year Power Spectrum



WMAP 9-year Power Spectrum



Planck 29-mo Power Spectrum






Planck 29-mo Power Spectrum



Clearly, the SW prediction does not fit!

$$\langle C_{\ell, \text{SW}} \rangle = \frac{8\pi^2 N_\phi^2 T_0^2}{9\ell^2} \left(\frac{\ell}{r_L} \right)^{n-1} \frac{\sqrt{\pi}}{2} \frac{\Gamma[(3-n)/2]}{\Gamma[(4-n)/2]}$$


Missing physics:
Hydrodynamics
(sound waves)







La Soupe Miso Cosmique

- When matter and radiation were hotter than 3000 K, matter was completely ionised. The Universe was filled with plasma, which behaves just like a soup
- Think about a Miso soup (if you know what it is). Imagine throwing Tofus into a Miso soup, while changing the density of Miso
- And imagine watching how ripples are created and propagate throughout the soup





This is a **viscous** fluid,
in which the amplitude of
sound waves **damps**
at shorter wavelength




When do sound waves become important?

- In other words, when would the Sachs-Wolfe approximation (purely gravitational effects) become invalid?
- The key to the answer: **Sound-crossing Time**
- Sound waves cannot alter temperature anisotropy at a given angular scale if there was not enough time for sound waves to propagate to the corresponding distance at the last-scattering surface
 - The distance traveled by sound waves within a given time = **The Sound Horizon**

Comoving Photon Horizon


- First, the comoving distance traveled by photons is given by setting the space-time distance to be null:

$$ds^2 = -c^2 dt^2 + a^2(t) dr^2 = 0$$


$$r_{\text{photon}} = c \int_0^t \frac{dt'}{a(t')}$$

Comoving Sound Horizon

- Then, we replace the speed of light with a time-dependent speed of sound:


$$r_s = \int_0^t \frac{dt'}{a(t')} c_s(t')$$

- We cannot ignore the effects of sound waves if $qr_s > 1$

Sound Speed

- Sound speed of an adiabatic fluid is given by

$$c_s^2 = \delta P / \delta \rho$$

- δP : pressure perturbation
- $\delta \rho$: density perturbation

- For a baryon-photon system:

$$c_s^2 = \delta P_\gamma / (\delta \rho_\gamma + \delta \rho_B)$$

We can ignore the baryon pressure because it is much smaller than the photon pressure

Sound Speed

- Using the adiabatic relationship between photons and baryons:

$$\delta\rho_B/\bar{\rho}_B = \delta\rho_\gamma/(\bar{\rho}_\gamma + \bar{P}_\gamma) = 3\delta\rho_\gamma/4\bar{\rho}_\gamma$$

[i.e., the ratio of the number densities of baryons and photons is equal everywhere]

- and pressure-density relation of a relativistic fluid, $\delta P_\gamma = \delta\rho_\gamma/3$,
We obtain

$$c_s^2 = \delta P_\gamma/(\delta\rho_\gamma + \delta\rho_B) = 1/3(1 + 3\bar{\rho}_B/4\bar{\rho}_\gamma)$$

- Or equivalently

$$c_s = \frac{1}{\sqrt{3(1 + R)}}$$

sound speed is reduced!

where

$$R \equiv 3\bar{\rho}_B/4\bar{\rho}_\gamma$$

Value of R?

- The baryon mass density goes like a^{-3} , whereas the photon energy density goes like a^{-4} . Thus, the ratio of the two, R , goes like a .
- The proportionality constant is:

$$R = \frac{3\Omega_B}{4\Omega_\gamma} \frac{a}{a_0} = 0.6120 \left(\frac{\Omega_B h^2}{0.022} \right) \frac{1091}{1+z}$$

where we used

$$\Omega_\gamma \equiv \frac{8\pi G \rho_{\gamma 0}}{3H_0^2} = 2.471 \times 10^{-5} h^{-2} \quad \text{for } T_0 = 2.725 \text{ K}$$

For the last-scattering redshift of $z_L=1090$
(or last-scattering temperature of $T_L=2974$ K),

$$r_s = 145.3 \text{ Mpc}$$

We cannot ignore the effects of sound waves
if $qr_s > 1$. Since $l \sim qr_L$, this means

$$l > r_L/r_s = 96$$

where we used $r_L=13.95$ Gpc

Creation of Sound Waves: Basic Equations

1. Conservation equations (energy and momentum)

2. Equation of state, relating pressure to energy density

$$P = P(\rho)$$

3. General relativistic version of the “Poisson equation”,
relating gravitational potential to energy density

$$\nabla^2 \Phi(t, \mathbf{x}) = 4\pi G a^2(t) \delta \rho_M(t, \mathbf{x})$$

4. Evolution of the “anisotropic stress” (viscosity)

Energy Conservation

- Total energy conservation:

$$\sum_{\alpha} \left\{ \delta \dot{\rho}_{\alpha} + \frac{\dot{a}}{a} (3\delta \rho_{\alpha} + 3\delta P_{\alpha} + \nabla^2 \pi_{\alpha}) - 3(\bar{\rho}_{\alpha} + \bar{P}_{\alpha}) \dot{\Psi} + \frac{1}{a^2} (\bar{\rho}_{\alpha} + \bar{P}_{\alpha}) \nabla^2 \delta u_{\alpha} \right\} = 0,$$

anisotropic stress:
[or, viscosity]
 $\Delta T_{ij} = a^2 \partial_i \partial_j \pi$

velocity potential
 $\mathbf{v}_{\alpha} = \frac{1}{a} \nabla \delta u_{\alpha}$

- C.f., Total energy conservation [unperturbed]

$$\sum_{\alpha} \left[\dot{\bar{\rho}}_{\alpha} + \frac{3\dot{a}}{a} (\bar{\rho}_{\alpha} + \bar{P}_{\alpha}) \right] = 0$$

Energy Conservation

- **Total energy conservation:**

$$\sum_{\alpha} \left\{ \delta \dot{\rho}_{\alpha} + \frac{\dot{a}}{a} (3\delta \rho_{\alpha} + 3\delta P_{\alpha} + \nabla^2 \pi_{\alpha}) - 3(\bar{\rho}_{\alpha} + \bar{P}_{\alpha}) \dot{\Psi} + \frac{1}{a^2} (\bar{\rho}_{\alpha} + \bar{P}_{\alpha}) \nabla^2 \delta u_{\alpha} \right\} = 0,$$

- **Again, this is the effect of locally-defined inhomogeneous scale factor, i.e.,**

- The spatial metric is given by $ds^2 = a^2(t) \exp(-2\Psi) d\mathbf{x}^2$

- Thus, locally we can define a new scale factor:

$$\tilde{a}(t, \mathbf{x}) = a(t) \exp(-\Psi)$$

Energy Conservation

- **Total energy conservation:**

$$\sum_{\alpha} \left\{ \delta \dot{\rho}_{\alpha} + \frac{\dot{a}}{a} (3\delta \rho_{\alpha} + 3\delta P_{\alpha} + \nabla^2 \pi_{\alpha}) - 3(\bar{\rho}_{\alpha} + \bar{P}_{\alpha}) \dot{\Psi} + \frac{1}{a^2} (\bar{\rho}_{\alpha} + \bar{P}_{\alpha}) \nabla^2 \delta u_{\alpha} \right\} = 0,$$

- **Momentum flux going outward (inward) -> reduction (increase) in the energy density**

$$\left(\begin{array}{l} \text{C.f., for a non-expanding medium:} \\ \dot{\rho} + \nabla \cdot (\rho \mathbf{v}) = 0 \end{array} \right)$$

Momentum Conservation

- **Total momentum conservation** $\mathbf{v}_\alpha = \frac{1}{a} \nabla \delta u_\alpha$

$$\sum_\alpha \left\{ \frac{\partial}{\partial t} [(\bar{\rho}_\alpha + \bar{P}_\alpha) \delta u_\alpha] + \frac{3\dot{a}}{a} (\bar{\rho}_\alpha + \bar{P}_\alpha) \delta u_\alpha + (\bar{\rho}_\alpha + \bar{P}_\alpha) \Phi + \delta P_\alpha + \nabla^2 \pi_\alpha \right\} = 0,$$

- **Cosmological redshift of the momentum**
- **Gravitational force given by potential gradient**
- **Force given by pressure gradient**
- **Force given by gradient of anisotropic stress**

Equation of State

- Pressure of non-relativistic species (i.e., baryons and cold dark matter) can be ignored relative to the energy density. Thus, we set them to zero: $\mathbf{P_B=0=P_D}$ and $\mathbf{\delta P_B=0=\delta P_D}$
- Unperturbed pressure of relativistic species (i.e., photons and relativistic neutrinos) is given by the third of the energy density, i.e., $\mathbf{P_\gamma=\rho_\gamma/3}$ and $\mathbf{P_\nu=\rho_\nu/3}$
- Perturbed pressure involves contributions from the **bulk viscosity**:
$$\delta P_\gamma = \frac{(\delta \rho_\gamma - \nabla^2 \pi_\gamma)}{3}$$
$$\delta P_\nu = \frac{(\delta \rho_\nu - \nabla^2 \pi_\nu)}{3}$$

The reason for this is that
trace of the stress-energy
of relativistic species

vanishes: $\sum_{\mu=0,1,2,3} T_{\mu}^{\mu} = 0$

- Pressure of dark matter and cold dark matter vanishes. Thus $T_{\mu}^{\mu} = 0$
- Unperturbed pressure and energy density of photons $T_0^0 + \sum_{i=1}^3 T_i^i = -\rho + 3P + \nabla^2 \pi = 0$

- Perturbed pressure involves contributions from the **bulk**

viscosity: $\delta P_{\gamma} = \frac{(\delta \rho_{\gamma} - \nabla^2 \pi_{\gamma})}{3}$

$$\delta P_{\nu} = \frac{(\delta \rho_{\nu} - \nabla^2 \pi_{\nu})}{3}$$

Two Remarks

- In the standard scenario:
 - Energy densities are conserved separately; thus we do not need to sum over all species
 - Momentum densities of photons and baryons are NOT conserved separately but they are coupled via Thomson scattering. This must be taken into account when writing down separate conservation equations

Conservation Equations for Photons and Baryons

- Fourier transformation replaces $\nabla^2 \rightarrow -q^2$

$$X(t, \mathbf{x}) = (2\pi)^{-3} \int d^3q X_{\mathbf{q}}(t) \exp(i\mathbf{q} \cdot \mathbf{x})$$

$$\frac{\partial}{\partial t}(\delta\rho_\gamma/\bar{\rho}_\gamma) - \frac{4q^2}{3a^2}\delta u_\gamma = 4\dot{\Psi}$$

$$\frac{\partial}{\partial t}(\delta\rho_B/\bar{\rho}_B) - \frac{q^2}{a^2}\delta u_B = 3\dot{\Psi}$$

$$a\frac{\partial}{\partial t}(\delta u_\gamma/a) + \Phi + \frac{\delta\rho_\gamma}{4\bar{\rho}_\gamma} - \frac{q^2\pi_\gamma}{2\bar{\rho}_\gamma} = \sigma_T\bar{n}_e(\delta u_B - \delta u_\gamma)$$

$$\delta\dot{u}_B + \Phi = -\frac{\sigma_T\bar{n}_e}{R}(\delta u_B - \delta u_\gamma)$$

momentum transfer via scattering

$$R \equiv 3\bar{\rho}_B/4\bar{\rho}_\gamma$$

Conservation Equations for Photons and Baryons

- Fourier transformation replaces $\nabla^2 \rightarrow -q^2$

$$X(t, \mathbf{x}) = (2\pi)^{-3} \int d^3q X_{\mathbf{q}}(t) \exp(i\mathbf{q} \cdot \mathbf{x})$$

$$\frac{\partial}{\partial t} (\delta\rho_\gamma / \bar{\rho}_\gamma) - \frac{4q^2}{3a^2} \delta u_\gamma = 4\dot{\Psi}$$

$$\frac{\partial}{\partial t} (\delta\rho_B / \bar{\rho}_B) - \frac{q^2}{a^2} \delta u_B = 3\dot{\Psi}$$

$$a \frac{\partial}{\partial t} (\delta u_\gamma / a) + \Phi + \frac{\delta\rho_\gamma}{4\bar{\rho}_\gamma} - \frac{q^2 \pi_\gamma}{2\bar{\rho}_\gamma} = \sigma_T \bar{n}_e (\delta u_B - \delta u_\gamma)$$

what about
photon's viscosity?

$$\delta\dot{u}_B + \Phi = -\frac{\sigma_T \bar{n}_e}{R} (\delta u_B - \delta u_\gamma)$$

$$R \equiv 3\bar{\rho}_B / 4\bar{\rho}_\gamma$$

Formation of a Photon-baryon Fluid

- Photons are not a fluid. Photons free-stream at the speed of light
 - The conservation equations are not enough because we need to specify the evolution of viscosity
 - Solving for viscosity requires information of the phase-space distribution function of photons: **Boltzmann equation**
- However, frequent scattering of photons with baryons* can make photons behave as a fluid: **Photon-baryon fluid**

**Photons scatter with electrons via Thomson scattering. Protons scatter with electrons via Coulomb scattering. Thus we can say, effectively, photons scatter with baryons*

Let's solve them!

- Fourier transformation replaces $\nabla^2 \rightarrow -q^2$

$$X(t, \mathbf{x}) = (2\pi)^{-3} \int d^3q X_{\mathbf{q}}(t) \exp(i\mathbf{q} \cdot \mathbf{x})$$

$$\frac{\partial}{\partial t}(\delta\rho_\gamma/\bar{\rho}_\gamma) - \frac{4q^2}{3a^2}\delta u_\gamma = 4\dot{\Psi}$$

$$\frac{\partial}{\partial t}(\delta\rho_B/\bar{\rho}_B) - \frac{q^2}{a^2}\delta u_B = 3\dot{\Psi}$$

$$a\frac{\partial}{\partial t}(\delta u_\gamma/a) + \Phi + \frac{\delta\rho_\gamma}{4\bar{\rho}_\gamma} - \frac{q^2\pi_\gamma}{2\bar{\rho}_\gamma} = \sigma_T\bar{n}_e(\delta u_B - \delta u_\gamma)$$

$$\delta\dot{u}_B + \Phi = -\frac{\sigma_T\bar{n}_e}{R}(\delta u_B - \delta u_\gamma)$$

$$R \equiv 3\bar{\rho}_B/4\bar{\rho}_\gamma$$

Tight-coupling Approximation

- When Thomson scattering is efficient, the relative velocity between photons and baryons is small. We write

$$\delta u_B - \delta u_\gamma = d / \sigma_T \bar{n}_e$$

[d is an arbitrary dimensionless variable]

- And take $\sigma_T \bar{n}_e \rightarrow \infty$ *. We obtain

$$a \frac{\partial}{\partial t} (\delta u_\gamma / a) + \Phi + \frac{\delta \rho_\gamma}{4 \bar{\rho}_\gamma} = d, \quad \delta \dot{u}_\gamma + \Phi = -\frac{d}{R}$$

**In this limit, viscosity π_γ is exponentially suppressed. This result comes from the Boltzmann equation but we do not derive it here. It makes sense physically.*

Tight-coupling Approximation

- Eliminating d and using the fact that R is proportional to the scale factor, we obtain

$$a \frac{\partial}{\partial t} [(1 + R) \delta u_\gamma / a] + (1 + R) \Phi + \frac{\delta \rho_\gamma}{4 \bar{\rho}_\gamma} = 0$$

- Using the energy conservation to replace δu_γ with $\delta \rho_\gamma / \rho_\gamma$, we obtain

$$\frac{1}{a(1 + R)} \frac{\partial}{\partial t} \left[a(1 + R) \frac{\partial}{\partial t} (\delta \rho_\gamma / \bar{\rho}_\gamma - 4\Psi) \right] + \frac{4q^2}{3a^2} \Phi + \frac{q^2}{a^2} \frac{\delta \rho_\gamma / \bar{\rho}_\gamma}{3(1 + R)} = 0$$

Wave Equation, with the speed of sound of $c_s^2 = 1/3(1+R)$!

Sound Wave!

- To simplify the equation, let's first look at the high-frequency solution
- Specifically, we take $q \gg aH$ (the wavelength of fluctuations is much shorter than the Hubble length). Then we can ignore time derivatives of R and Ψ because they evolve in the Hubble time scale:

$$\frac{1}{a} \frac{\partial}{\partial t} \left[a \frac{\partial}{\partial t} (\delta \rho_\gamma / \bar{\rho}_\gamma) \right] + \frac{q^2 c_s^2}{a^2} [\delta \rho_\gamma / \bar{\rho}_\gamma + 4(1 + R)\Phi] = 0$$


Solution: SOUND WAVE!

$$\frac{\delta \rho_\gamma}{4\bar{\rho}_\gamma} + \Phi = A \cos(qr_s) + B \sin(qr_s) - R\Phi$$


Recap


- Photons are not a fluid; but Thomson scattering couples photons to baryons, forming a **photon-baryon fluid**
- The reduced sound speed, $c_s^2 = 1/3(1+R)$, emerges automatically
- $\delta\rho_\gamma/4\rho_\gamma$ is the temperature anisotropy at the bottom of the potential well. Adding gravitational redshift, the observed temperature anisotropy is $\delta\rho_\gamma/4\rho_\gamma + \Phi$, which is given by

$$\frac{\delta\rho_\gamma}{4\bar{\rho}_\gamma} + \Phi = A \cos(qr_s) + B \sin(qr_s) - R\Phi$$



Effect of Baryon-Density





Stone: Fluctuations “entering the horizon”

- This is a tricky concept, but it is important
- Suppose that there are fluctuations at all wavelengths, including the ones that exceed the Hubble length (which we loosely call our “horizon”)
 - Let’s not ask the origin of these “super-horizon fluctuations”, but just assume their existence
- As the Universe expands, our horizon grows and we can see longer and longer wavelengths
 - **Fluctuations “entering the horizon”**

Physical Length [h^{-1} Mpc]

Last scattering



Radiation Era

Matter Era

10 Gpc/h today

1 Gpc/h today

100 Mpc/h today

10 Mpc/h today

1 Mpc/h today

“enter the horizon”


— Horizon

- - - Hubble length

a_{EQ}

a_L

Scale Factor, a/a_0



Three Regimes

- **Super-horizon scales [$q < aH$]**
 - Only gravity is important
 - Evolution differs from Newtonian
- **Sub-horizon but super-sound-horizon [$aH < q < aH/c_s$]**
 - Only gravity is important
 - Evolution similar to Newtonian
- **Sub-sound-horizon scales [$q > aH/c_s$]**
 - Hydrodynamics important -> Sound waves

ζ :

Conserved on large scales

- For the adiabatic initial condition, there exists a useful quantity, ζ , which **remains constant on large scales** (*super-horizon scales*, $q \ll aH$) regardless of the contents of the Universe
 - ζ is conserved regardless of whether the Universe is radiation-dominated, matter-dominated, or whatever
- Energy conservation for $q \ll aH$:

$$\delta \dot{\rho}_\alpha + \frac{3\dot{a}}{a}(\delta \rho_\alpha + \delta P_\alpha) - 3(\bar{\rho}_\alpha + \bar{P}_\alpha)\dot{\Psi} = 0$$


ζ :

Conserved on large scales

- If pressure is a function of the energy density only, i.e., $P_\alpha = P_\alpha(\rho_\alpha)$, then

$$\frac{1}{3} \frac{\delta \rho_\alpha(t, \mathbf{x})}{\bar{\rho}_\alpha(t) + \bar{P}_\alpha(t)} - \Psi(t, \mathbf{x}) = \zeta_\alpha(\mathbf{x})$$

integration constant



Integrate

$$\delta \dot{\rho}_\alpha + \frac{3\dot{a}}{a}(\delta \rho_\alpha + \delta P_\alpha) - 3(\bar{\rho}_\alpha + \bar{P}_\alpha)\dot{\Psi} = 0$$

ζ :

Conserved on large scales

- If pressure is a function of the energy density only, i.e., $P_\alpha = P_\alpha(\rho_\alpha)$, then

$$\frac{1}{3} \frac{\delta \rho_\alpha(t, \mathbf{x})}{\bar{\rho}_\alpha(t) + \bar{P}_\alpha(t)} - \Psi(t, \mathbf{x}) = \zeta_\alpha(\mathbf{x})$$


integration constant

For the adiabatic initial condition, all species share the same value of ζ_α , i.e., $\zeta_\alpha = \zeta$


$$q_{\text{EQ}}$$

- Which fluctuation entered the horizon before the matter-radiation equality?
- $q_{\text{EQ}} = a_{\text{EQ}} H_{\text{EQ}} \sim 0.01 (\Omega_{\text{M}} h^2 / 0.14) \text{ Mpc}^{-1}$
- At the last scattering surface, this subtends the multipole of $l_{\text{EQ}} = q_{\text{EQ}} r_{\text{L}} \sim 140$

Effect of Baryon-Density



Effect of Baryon-Density



Peak Locations?

High-frequency solution, for $q \gg aH$

$$\frac{\delta\rho_\gamma}{4\bar{\rho}_\gamma} + \Phi = A \cos(qr_s) + B \sin(qr_s) - R\Phi$$

- VERY roughly speaking, the angular power spectrum C_l is given by $\left[\frac{\delta\rho_\gamma}{4\bar{\rho}_\gamma} + \Phi \right]^2$ with $q \rightarrow l/r_L$
 - Question: What are the integration constants, **A** and **B**?
 - Answer: They depend on the initial conditions; namely, adiabatic or not?
 - For adiabatic initial condition, **$A \gg B$ when q is large**
[We will show it later.]

Peak Locations?


High-frequency solution, for $q \gg aH$

$$\frac{\delta\rho_\gamma}{4\bar{\rho}_\gamma} + \Phi = A \cos(qr_s) + B \sin(qr_s) - R\Phi$$

- VERY roughly speaking, the angular power spectrum C_l is given by $\left[\frac{\delta\rho_\gamma}{4\bar{\rho}_\gamma} + \Phi \right]^2$ with $q \rightarrow l/r_L$
- If $A \gg B$, the locations of peaks are

$$l = (1, 2, \dots) \pi r_L / r_s(t_L) = (1, 2, \dots) \times 302$$

Effect of Baryon-Density



$\Delta T_1 [\mu K]$ 110
100
90
80
70
60
50
40
30
20
10
0

The simple estimates do
not match!

This is simply because
these angular scales do
not satisfy $q \gg aH$, i.e, the
oscillations are not pure
cosine even for the
adiabatic initial condition.

We need a better solution!

10

Better Solution in Radiation-dominated Era

Going back to the original tight-coupling equation..

$$\frac{1}{a(1+R)} \frac{\partial}{\partial t} \left[a(1+R) \frac{\partial}{\partial t} (\delta\rho_\gamma/\bar{\rho}_\gamma - 4\Psi) \right] + \frac{4q^2}{3a^2} \Phi + \frac{q^2}{a^2} \frac{\delta\rho_\gamma/\bar{\rho}_\gamma}{3(1+R)} = 0$$

- In the radiation-dominated era, $R \ll 1$
- Change the independent variable from the time (t) to

$$\varphi \equiv qr_s = 2qt/\sqrt{3}a$$

Better Solution in Radiation-dominated Era

Then the equation simplifies to

$$\partial^2 X / \partial \varphi^2 + X + \Phi + \Psi = 0$$

where $X \equiv \delta \rho_\gamma / 4\bar{\rho}_\gamma - \Psi$

- In the radiation-dominated era, $R \ll 1$
- Change the independent variable from the time (t) to

$$\varphi \equiv q r_s = 2qt / \sqrt{3}a$$

Better Solution in Radiation-dominated Era

Then the equation simplifies to

$$\partial^2 X / \partial \varphi^2 + X + \Phi + \Psi = 0$$

where $X \equiv \delta \rho_\gamma / 4 \bar{\rho}_\gamma - \Psi$

The solution is

$$X = \tilde{A} \cos \varphi + \tilde{B} \sin \varphi - \int_0^\varphi d\varphi' \sin(\varphi - \varphi') (\Phi + \Psi)(\varphi')$$

We rewrite this using the formula for trigonometry:

$$\sin(\varphi - \varphi') = \sin(\varphi) \cos(\varphi') - \cos(\varphi) \sin(\varphi')$$

Better Solution in Radiation-dominated Era

Then the equation simplifies to

$$\partial^2 X / \partial \varphi^2 + X + \Phi + \Psi = 0$$

where $X \equiv \delta \rho_\gamma / 4 \bar{\rho}_\gamma - \Psi$

The solution is

$$X = (\tilde{A} + \Delta A) \cos \varphi + (\tilde{B} + \Delta B) \sin \varphi$$

where

$$\Delta A(\varphi) \equiv \int_0^\varphi d\varphi' \sin \varphi' (\Phi + \Psi)(\varphi'),$$
$$\Delta B(\varphi) \equiv - \int_0^\varphi d\varphi' \cos \varphi' (\Phi + \Psi)(\varphi')$$

Einstein's Equations

- Now we need to know Newton's gravitational potential, ϕ , and the scalar curvature perturbation, ψ .
- Einstein's equations - let's look up any text books:

$$\nabla^2 \Psi = 4\pi G a^2 \sum_{\alpha} \left[\delta \rho_{\alpha} - \frac{3\dot{a}}{a} (\bar{\rho}_{\alpha} + \bar{P}_{\alpha}) \delta u_{\alpha} \right]$$

$$\dot{\Psi} + \frac{\dot{a}}{a} \Phi = -4\pi G \sum_{\alpha} (\bar{\rho}_{\alpha} + \bar{P}_{\alpha}) \delta u_{\alpha}$$

$$\partial_i \partial_j (\Phi - \Psi) = -8\pi G a^2 \partial_i \partial_j \sum_{\alpha} \pi_{\alpha}$$

Einstein's Equations

- Now we need to know Newton's gravitational potential, ϕ , and the scalar curvature perturbation, ψ .
- Einstein's equations - let's look up any text books:

$$\nabla^2 \Psi = 4\pi G a^2 \sum_{\alpha} \left[\delta \rho_{\alpha} - \frac{3\dot{a}}{a} (\bar{\rho}_{\alpha} + \bar{P}_{\alpha}) \delta u_{\alpha} \right]$$

$$\dot{\Psi} + \frac{\dot{a}}{a} \Phi = -4\pi G \sum_{\alpha} (\bar{\rho}_{\alpha} + \bar{P}_{\alpha}) \delta u_{\alpha}$$

$$\partial_i \partial_j (\Phi - \Psi) = -8\pi G a^2 \partial_i \partial_j \left(\sum_{\alpha} \pi_{\alpha} \right)$$

Einstein's Equations

- Now we need to know Newton's gravitational potential, ϕ , and the scalar curvature perturbation, ψ .
- Einstein's equations - let's look up any text books:

$$\nabla^2 \Psi = 4\pi G a^2 \sum_{\alpha} \left[\delta \rho_{\alpha} - \frac{3\dot{a}}{a} (\bar{\rho}_{\alpha} + \bar{P}_{\alpha}) \delta u_{\alpha} \right]$$

$$\dot{\Psi} + \frac{\dot{a}}{a} \Phi = -4\pi G \sum_{\alpha} (\bar{\rho}_{\alpha} + \bar{P}_{\alpha}) \delta u_{\alpha}$$

$$\partial_i \partial_j (\Phi - \Psi) = -8\pi G a^2 \partial_i \partial_j \pi_{\nu}$$

**Will come back to
this later.
For now, let's ignore
any viscosity.**

Einstein's Equations

- Now we need to know Newton's gravitational potential, ϕ , and the scalar curvature perturbation, ψ .
- Einstein's equations - let's look up any text books:

$$\nabla^2 \Psi = 4\pi G a^2 \sum_{\alpha} \left[\delta \rho_{\alpha} - \frac{3\dot{a}}{a} (\bar{\rho}_{\alpha} + \bar{P}_{\alpha}) \delta u_{\alpha} \right]$$

$$\dot{\Psi} + \frac{\dot{a}}{a} \Phi = -4\pi G \sum_{\alpha} (\bar{\rho}_{\alpha} + \bar{P}_{\alpha}) \delta u_{\alpha}$$

$$\Phi = \Psi$$

**Will come back to
this later.
For now, let's ignore
any viscosity.**

Einstein's Equations in Radiation-dominated Era

- Now we need to know Newton's gravitational potential, Φ , and the scalar curvature perturbation, ψ .
- Einstein's equations - let's look up any text books:

$$\frac{\partial^2 \Phi}{\partial \varphi^2} + \frac{4}{\varphi} \frac{\partial \Phi}{\partial \varphi} + \Phi = \frac{3}{2\varphi^2} \frac{\delta \mathcal{P}}{\bar{\rho}_R}$$

$$\sum_{\alpha} \delta P_{\alpha}(t, \mathbf{x}) = \frac{\sum_{\alpha} \dot{\bar{P}}_{\alpha}(t)}{\sum_{\alpha} \dot{\bar{\rho}}_{\alpha}(t)} \sum_{\alpha} \delta \rho_{\alpha}(t, \mathbf{x}) + \delta \mathcal{P}(t, \mathbf{x})$$

“non-adiabatic” pressure

Einstein's Equations in Radiation-dominated Era

- Now we need to know Newton's gravitational potential, ϕ , and the scalar curvature perturbation, ψ .
- Einstein's equations - let's look up any text books:

$$\frac{\partial^2 \Phi}{\partial \varphi^2} + \frac{4}{\varphi} \frac{\partial \Phi}{\partial \varphi} + \Phi = \frac{3}{2\varphi^2} \frac{\delta \mathcal{P}}{\bar{\rho}_R}$$

We shall ignore this

$$\sum_{\alpha} \delta P_{\alpha}(t, \mathbf{x}) = \frac{\sum_{\alpha} \dot{P}_{\alpha}(t)}{\sum_{\alpha} \dot{\bar{\rho}}_{\alpha}(t)} \sum_{\alpha} \delta \rho_{\alpha}(t, \mathbf{x}) + \delta \mathcal{P}(t, \mathbf{x})$$

“non-adiabatic” pressure

Solution (Adiabatic) in Radiation-dominated Era

$$\Phi_{\text{ADI}} = -2\zeta(\sin \varphi - \varphi \cos \varphi) / \varphi^3$$

where

$$\varphi \equiv qr_s = 2qt / \sqrt{3}a$$

- Low-frequency limit (*super-sound-horizon scales*, $qr_s \ll 1$)
 - $\Phi_{\text{ADI}} \rightarrow -2\zeta/3 = \text{constant}$
- High-frequency limit (*sub-sound-horizon scales*, $qr_s \gg 1$)
 - $\Phi_{\text{ADI}} \rightarrow 2\zeta \cos \varphi / \varphi^2 \propto a^{-2}$ **damp**

Solution (Adiabatic) in Radiation-dominated Era

$$\Phi_{\text{ADI}} = -2\zeta(\sin \varphi - \varphi \cos \varphi) / \varphi^3$$

where

Poisson Equation

$$-q^2 \Phi = 4\pi G a^2 \delta \rho$$

- Low-frequency (sub sound horizon scales, $qr_s \ll 1$)

& oscillation solution for radiation

- $\Phi_{\text{ADI}} \rightarrow -2\zeta/3$

$$\delta \rho_R / \bar{\rho}_R \propto \cos \varphi$$

- High-frequency in (super sound horizon scales, $qr_s \gg 1$)

- $\Phi_{\text{ADI}} \rightarrow 2\zeta \cos \varphi / \varphi^2 \propto a^{-2}$ **damp**

Solution (Adiabatic) in Radiation-dominated Era

$$\Phi_{\text{ADI}} = -2\zeta(\sin \varphi - \varphi \cos \varphi) / \varphi^3$$

where

$$\varphi \equiv qr_s = 2qt / \sqrt{3}a$$

- Low-frequency limit (*super-sound-horizon scales*, $qr_s \ll 1$)
 - $\Phi_{\text{ADI}} \rightarrow -2\zeta/3 = \text{constant}$
- High-frequency limit (*sub-sound-horizon scales*, $qr_s \gg 1$)
 - $\Phi_{\text{ADI}} \rightarrow 2\zeta \cos \varphi / \varphi^2 \propto a^{-2}$ **damp**

Sound Wave Solution in the Radiation-dominated Era

The solution is

$$X = (\tilde{A} + \Delta A) \cos \varphi + (\tilde{B} + \Delta B) \sin \varphi$$

where

$$X \equiv \delta \rho_\gamma / 4 \bar{\rho}_\gamma - \Psi$$

$$\Delta A(\varphi) \equiv \int_0^\varphi d\varphi' \sin \varphi' (\Phi + \Psi)(\varphi'),$$

$$\Delta B(\varphi) \equiv - \int_0^\varphi d\varphi' \cos \varphi' (\Phi + \Psi)(\varphi')$$

Sound Wave Solution in the Radiation-dominated Era

The solution is

$$X = (\tilde{A} + \Delta A) \cos \varphi + (\tilde{B} + \Delta B) \sin \varphi$$

where

$$X \equiv \delta \rho_\gamma / 4 \bar{\rho}_\gamma - \Psi$$

$$\Delta A(\varphi) \equiv \int_0^\varphi d\varphi' \sin \varphi' (\Phi + \Psi)(\varphi') = -2\zeta(1 - \sin^2 \varphi / \varphi^2)$$

$$\Delta B(\varphi) \equiv - \int_0^\varphi d\varphi' \cos \varphi' (\Phi + \Psi)(\varphi')$$

$$= 2\zeta(\varphi - \cos \varphi \sin \varphi) / \varphi^2$$

Sound Wave Solution in the Radiation-dominated Era

The solution is

$$X = (\tilde{A} + \Delta A) \cos \varphi + (\tilde{B} + \Delta B) \sin \varphi$$

where

$$X \equiv \delta\rho_\gamma/4\bar{\rho}_\gamma - \Psi \xrightarrow{\varphi \ll 1} \zeta \quad \text{i.e., } \underline{\tilde{A}_{\text{ADI}} = \zeta, \tilde{B}_{\text{ADI}} = 0}$$

$$\Delta A(\varphi) \equiv \int_0^\varphi d\varphi' \sin \varphi' (\Phi + \Psi)(\varphi') = \underline{-2\zeta(1 - \sin^2 \varphi/\varphi^2)}$$

$$\Delta B(\varphi) \equiv -\int_0^\varphi d\varphi' \cos \varphi' (\Phi + \Psi)(\varphi')$$

$$= \underline{2\zeta(\varphi - \cos \varphi \sin \varphi)/\varphi^2}$$

Sound Wave Solution in the Radiation-dominated Era

The adiabatic solution is


$$X = \frac{\delta\rho_\gamma}{4\bar{\rho}_\gamma} - \Psi = \zeta \left(-\cos\varphi + \frac{2}{\varphi} \sin\varphi \right)$$

with

$$\Phi = \Psi = -2\zeta(\sin\varphi - \varphi\cos\varphi)/\varphi^3$$

Therefore, the solution is a **pure cosine**
only in the **high-frequency** limit!

Effect of Baryon-Density



Roles of viscosity

- **Neutrino viscosity**

- Modify potentials: $\partial_i \partial_j (\Phi - \Psi) = -8\pi G a^2 \partial_i \partial_j \pi_\nu$

- **Photon viscosity**

- Viscous photon-baryon fluid: **damping of sound waves**

Silk (1968) "Silk damping"

$$a \frac{\partial}{\partial t} (\delta u_\gamma / a) + \Phi + \frac{\delta \rho_\gamma}{4\bar{\rho}_\gamma} - \frac{q^2 \pi_\gamma}{2\bar{\rho}_\gamma} = \sigma_T \bar{n}_e (\delta u_B - \delta u_\gamma)$$

High-frequency solution without neutrino viscosity

The solution is

$$\underline{X = (\zeta + \Delta A) \cos \varphi + (\Delta B) \sin \varphi}$$

where

$$X \equiv \delta\rho_\gamma / 4\bar{\rho}_\gamma - \Psi$$

$$\Delta A(\varphi) \equiv \int_0^\varphi d\varphi' \sin \varphi' (\Phi + \Psi)(\varphi') = -2\zeta (1 - \sin^2 \varphi / \varphi^2)$$

$$\Delta B(\varphi) \equiv - \int_0^\varphi d\varphi' \cos \varphi' (\Phi + \Psi)(\varphi') \xrightarrow{\varphi \gg 1} -2\zeta$$

$$= 2\zeta (\varphi - \cos \varphi \sin \varphi) / \varphi^2 \xrightarrow{\varphi \gg 1} \underline{0}$$

High-frequency solution with neutrino viscosity

The solution is

$$\underline{X} = (-\zeta + \Delta A_\nu) \cos \varphi + \Delta B_\nu \sin \varphi$$

where

$$X \equiv \delta\rho_\gamma / 4\bar{\rho}_\gamma - \Psi$$

$$\Delta A_\nu \xrightarrow{\varphi \gg 1} 0.338 R_\nu \zeta$$

$$\Delta B_\nu \xrightarrow{\varphi \gg 1} 0.418 R_\nu \zeta$$

non-zero value!

$$R_\nu \equiv \bar{\rho}_\nu / (\bar{\rho}_\gamma + \bar{\rho}_\nu) \approx 0.409$$

High-frequency solution **with** neutrino viscosity

Using the formula for trigonometry, we write

$$X = -C \cos(\varphi + \theta)$$

where

$$C \equiv \sqrt{(-\zeta + \Delta A_\nu)^2 + \Delta B_\nu^2}$$

$$\approx \zeta (1 + 4R_\nu/15)^{-1} \quad \text{Hu \& Sugiyama (1996)}$$

$$\tan \theta = -\frac{\Delta B_\nu}{-\zeta + \Delta A_\nu} \approx 0.063\pi \quad \text{Phase shift!}$$

Bashinsky & Seljak (2004)

High-frequency solution

with neutrino viscosity

Thus, the neutrino viscosity will:

The s

X

where

C

- (1) Reduce the amplitude of sound waves at large multipoles
- (2) Shift the peak positions of the temperature power spectrum

$$\tan \theta = -\frac{\Delta B_\nu}{-\zeta + \Delta A_\nu} \approx 0.063\pi \quad \text{Phase shift!}$$

Bashinsky & Seljak (2004)

Photon Viscosity

- In the tight-coupling approximation, the photon viscosity damps exponentially
- To take into account a non-zero photon viscosity, we go to a higher order in the tight-coupling approximation

Yesterday: Tight-coupling Approximation (1st-order)

- When Thomson scattering is efficient, the relative velocity between photons and baryons is small. We write

$$\delta u_B - \delta u_\gamma = d / \sigma_T \bar{n}_e$$

[d is an arbitrary dimensionless variable]

- And take $\sigma_T \bar{n}_e \rightarrow \infty$ *. We obtain

$$a \frac{\partial}{\partial t} (\delta u_\gamma / a) + \Phi + \frac{\delta \rho_\gamma}{4 \bar{\rho}_\gamma} = d, \quad \delta \dot{u}_\gamma + \Phi = -\frac{d}{R}$$

**In this limit, viscosity π_γ is exponentially suppressed. This result comes from the Boltzmann equation but we do not derive it here. It makes sense physically.*

Today: Tight-coupling Approximation (2nd-order)

- When Thomson scattering is efficient, the relative velocity between photons and baryons is small. We write

$$\delta u_B - \delta u_\gamma = d_1 / \sigma_T \bar{n}_e + \underline{q d_2 / (\sigma_T \bar{n}_e)^2}$$

where

$$d_1 = -R(\delta \dot{u}_\gamma + \Phi) \quad [\mathbf{d_2 \text{ is an arbitrary dimensionless variables}}]$$

- And take $\sigma_T \bar{n}_e \rightarrow \infty$. We obtain

$$a \frac{\partial}{\partial t} (\delta u_\gamma / a) + \Phi + \frac{\delta \rho_\gamma}{4 \bar{\rho}_\gamma} - \frac{q^2 \pi_\gamma}{2 \bar{\rho}_\gamma} = -R(\delta \dot{u}_\gamma + \Phi) + \frac{q}{\sigma_T \bar{n}_e} d_2$$

$$\frac{\partial}{\partial t} \left[\frac{R(\delta \dot{u}_\gamma + \Phi)}{\sigma_T \bar{n}_e} \right] = \frac{q}{R \sigma_T \bar{n}_e} d_2$$

Tight-coupling Approximation (2nd-order)

- Eliminating d_2 and using the fact that R is proportional to the scale factor, we obtain

$$a \frac{\partial}{\partial t} [(1 + R) \delta u_\gamma / a] + (1 + R) \Phi + \frac{\delta \rho_\gamma}{4 \bar{\rho}_\gamma} - \frac{q^2 \pi_\gamma}{2 \bar{\rho}_\gamma} + R \frac{\partial}{\partial t} \left[\frac{R(\delta \dot{u}_\gamma + \Phi)}{\sigma_T \bar{n}_e} \right] = 0$$

- Getting π_γ requires an approximate solution of the Boltzmann equation in the 2nd-order tight coupling. We do not derive it here. The answer is

$$\Delta T_{ij} = a^2 \partial_i \partial_j \pi_\gamma = - \frac{32}{45} \frac{\bar{\rho}_\gamma}{\sigma_T \bar{n}_e} \partial_i \partial_j \delta u_\gamma$$

Tight-coupling Approximation (2nd-order)

- Eliminating d_2 and using the fact that R is proportional to the scale factor, we obtain

$$a \frac{\partial}{\partial t} [(1 + R) \delta u_\gamma / a] + (1 + R) \Phi + \frac{\delta \rho_\gamma}{4 \bar{\rho}_\gamma} - \frac{q^2 \pi_\gamma}{2 \bar{\rho}_\gamma} + R \frac{\partial}{\partial t} \left[\frac{R(\delta \dot{u}_\gamma + \Phi)}{\sigma_T \bar{n}_e} \right] = 0$$

- Getting π_γ requires an approximation equation in the 2nd-order tight coupling here. The answer is

Given by the **spatial gradient of the velocity field**

- a well-known result in fluid dynamics

$$\Delta T_{ij} = a^2 \partial_i \partial_j \pi_\gamma = - \frac{32}{45} \frac{\bar{\rho}_\gamma}{\sigma_T \bar{n}_e} \partial_i \partial_j \delta u_\gamma$$

Damped Oscillator

- Using the energy conservation to replace δu_γ with $\delta \rho_\gamma / \rho_\gamma$, we obtain, for $q \gg aH$,

$$\frac{1}{a} \frac{\partial}{\partial t} \left[a \frac{\partial}{\partial t} (\delta \rho_\gamma / \bar{\rho}_\gamma) \right] + 2\Gamma \frac{\partial}{\partial t} (\delta \rho_\gamma / \bar{\rho}_\gamma) + \frac{q^2 c_s^2}{a^2} [\delta \rho_\gamma / \bar{\rho}_\gamma + 4(1 + R)\Phi] = 0$$

New term, giving damping!

where

$$\Gamma(q, t) \equiv \frac{q^2}{6a^2 \sigma_T \bar{n}_e} \left[\frac{16}{15(1 + R)} + \frac{R^2}{(1 + R)^2} \right]$$

Damped Oscillator

- Using the energy conservation to replace δu_γ with $\delta \rho_\gamma / \rho_\gamma$, we obtain, for $q \gg aH$,

$$\frac{1}{a} \frac{\partial}{\partial t} \left[a \frac{\partial}{\partial t} (\delta \rho_\gamma / \bar{\rho}_\gamma) \right] + 2\Gamma \frac{\partial}{\partial t} (\delta \rho_\gamma / \bar{\rho}_\gamma) + \frac{q^2 c_s^2}{a^2} [\delta \rho_\gamma / \bar{\rho}_\gamma + 4(1 + R)\Phi] = 0$$

New term, giving damping!

where

Important for high frequencies
(large multipoles)

$$\Gamma(q, t) \equiv \frac{q^2}{6a^2 \sigma_T \bar{n}_e} \left[\frac{16}{15(1 + R)} + \frac{R^2}{(1 + R)^2} \right]$$

Damped Oscillator

- Using the energy conservation to replace δu_γ with $\delta \rho_\gamma / \rho_\gamma$, we obtain, for $q \gg aH$,

$$\frac{1}{a} \frac{\partial}{\partial t} \left[a \frac{\partial}{\partial t} (\delta \rho_\gamma / \bar{\rho}_\gamma) \right] + 2\Gamma \frac{\partial}{\partial t} (\delta \rho_\gamma / \bar{\rho}_\gamma) + \frac{q^2 c_s^2}{a^2} [\delta \rho_\gamma / \bar{\rho}_\gamma + 4(1 + R)\Phi] = 0$$

New term, giving damping!

SOLUTION:

$$\frac{\delta \rho_\gamma}{4\bar{\rho}_\gamma} + \Phi = [A \cos(qr_s) + B \sin(qr_s)] \exp \left[- \int_0^t dt' \Gamma(q, t') \right] - R\Phi$$

Exponential damping!

$$\approx \exp \left(-q^2 / \sigma_\tau \bar{n}_e H \right)$$

Damped Oscillator

- Using the energy conservation to replace δu_γ with $\delta \rho_\gamma / \rho_\gamma$, we obtain, for $q \gg aH$,

$$\frac{1}{a} \frac{\partial}{\partial t} \left[a \frac{\partial}{\partial t} (\delta \rho_\gamma / \bar{\rho}_\gamma) \right] + 2\Gamma \frac{\partial}{\partial t} (\delta \rho_\gamma / \bar{\rho}_\gamma) + \frac{q^2 c_s^2}{a^2} [\delta \rho_\gamma / \bar{\rho}_\gamma + 4(1 + R)\Phi] = 0$$

New term, giving damping!

SOLUTION:

$$\frac{\delta \rho_\gamma}{4\bar{\rho}_\gamma} + \Phi = [A \cos(qr_s) + B \sin(qr_s)] \exp(-q^2 / q_{\text{Silk}}^2) - R\Phi$$


Exponential damping!

$$a/q_{\text{Silk}} \approx (\sigma \tau \bar{n}_e H)^{-1/2} \quad \begin{array}{l} \text{“diffusion length”} \\ \text{= length traveled by photon's random walks} \end{array}$$


Diffusion Length

- The mean free path of the photon between scatterings is $(\sigma_T n_e)^{-1}$
 - Below this scale, you do not have a photon-baryon fluid
- The number of scatterings per Hubble time is $N_{\text{scattering}} = \sigma_T n_e / H$
- Then, the length traveled by photons by random walks within the Hubble time is $(\sigma_T n_e)^{-1}$ times $\sqrt{N_{\text{scatterings}}}$
 - The diffusion length is thus $(\sigma_T n_e)^{-1}$ times $\sqrt{N_{\text{scatterings}}} = (\sigma_T n_e H)^{-1/2}$

Diffusion Damping




- Diffusion mixes hot and cold photons -> Damping of anisotropies



Additional Damping

- The power spectrum is $\left[\frac{\delta\rho_\gamma}{4\bar{\rho}_\gamma} + \Phi \right]^2$ with $q \rightarrow l/r_L$. The damping factor is thus $\exp(-2q^2/q_{\text{silk}}^2)$
- $q_{\text{silk}}(t_L) = 0.139 \text{ Mpc}^{-1}$. This corresponds to a multipole of $l_{\text{silk}} \sim q_{\text{silk}} r_L / \sqrt{2} = 1370$. Seems too large, compared to the exact calculation
- There is an additional damping due to a finite width of the last scattering surface, $\sigma \sim 250 \text{ K}$
 - “Fuzziness damping” – Bond (1996)
 - “Landau damping” - Weinberg (2001)

$$q_{\text{fuzziness}}^{-2} = \frac{3\sigma^2 t_L^2}{8a_0^2 T_0^2 (1 + R_L)} \approx (0.20 \text{ Mpc}^{-1})^{-2}$$



Recap

- The basic structure of the temperature power spectrum is
 - The Sachs-Wolfe “plateau” at low multipoles
 - Sound waves at intermediate multipoles
 - 1st-order tight-coupling
 - Silk damping and Fuzziness damping at high multipoles
 - 2nd-order tight-coupling